

## **High risk of infection caused Posttraumatic Stress symptoms in individuals with poor sleep quality: A study on influence of Coronavirus disease (COVID-19) in China**

Fan Zhang<sup>1,2,+</sup>, Zhilei Shang<sup>1,2,+</sup>, Haiying Ma<sup>3,+</sup>, Yanpu Jia<sup>1,2,+</sup>, Luna Sun<sup>1,2</sup>, Xin Guo<sup>4</sup>, Lili Wu<sup>1,2</sup>, Zhuoer Sun<sup>1,2</sup>, Yaoguang Zhou<sup>1,2</sup>, Yan Wang<sup>1,2</sup>, Nianqi Liu<sup>1,2</sup>, Weizhi Liu<sup>1,2,\*</sup>

<sup>1</sup> Lab for Post-traumatic Stress Disorder, Faculty of Psychology and Mental Health, Naval Medical University, Shanghai 200433, China

<sup>2</sup> The Emotion & Cognition Lab, Faculty of Psychology and Mental Health, Naval Medical University, Shanghai 200433, China

<sup>3</sup> Counselling and Psychological Services Center, Faculty of Psychology and Mental Health, Naval Medical University, Shanghai 200433, China

<sup>4</sup> The Battalion 5 of Cadet Brigade, School of Basic Medicine, Naval Medical University, Shanghai 200433, China

\*Corresponding. 13024141970@163.com

+ these authors contributed equally to this work.

### **Abstract**

The influence of the outbreak of coronavirus disease (COVID-19) on mental health was poorly understood. The present study aimed to exam sleep problems and posttraumatic stress symptoms (PTSS) in Chinese immediately after the massive outbreak of COVID-19. A total of 2027 Chinese participated in the present study. Wuhan-expose history, sleep quality and PTSS were measured with self-rating scales. Results showed that there were significant differences of PCL-5 and of sleep quality scores in different data-collection dates ( $ps < 0.05$ ). There were significant differences of PCL-5 scores ( $t = -2.93$ ,  $p < 0.05$ ) and latency onset of sleep ( $\chi^2 = 9.77$ ,  $p < 0.05$ ) between participants with and without Wuhan-expose history. The interaction effect of Wuhan exposure history  $\times$  sleep quality significantly influenced PCL-5 ( $ps < 0.05$ ). These results indicate that keeping good sleep quality in individuals with high infectious risk is a way to prevent PTSS.

### **Key words**

posttraumatic stress disorder (PTSS); infectious disease; novel coronavirus; COVID-19; sleep quality

### **Introduction**

On Dec 31, 2019, the World Health Organization (WHO) China Office was informed of cases of an unknown etiology detected in Wuhan City, Hubei province of China (WHO, 2020a). On Jan 7, 2020, a novel coronavirus, which had not been documented before was identified by the Chinese Center for Disease Control and Prevention (CDC) from the throat swab sample of a patient, and was confirmed as the cause of the Coronavirus disease (COVID-19) by WHO (2020a). On Jan 30, 2020 the outbreak of COVID-19 has been determined by the WHO as a Public Health Emergency of International Concern (PHEIC) (WHO, 2020b). Thus far (up to Feb 4, 2020), more than 20,000 firmed cases have been confirmed in China, and more than 20 countries reported confirmed cases (WHO, 2020c).

According to the Diagnostic and Statistical Manual of Mental Disorders-5th edition (DSM-5) (APA, 2013), Posttraumatic Stress Disorder (PTSD) is characterized by four symptom groups: (a) involuntary memories of the trauma such as intrusions or nightmares; (b) persistent avoidance of stimuli associated with the traumatic event; (c) negative alterations in cognitions and mood that are associated with the trauma; and (d) alterations in arousal and reactivity that are associated with the trauma. Also in DSM-5, exposure to traumatic events is the immediate causes of PTSD and an essential condition of diagnosis. Being infected with a life-threatening physical illness is a traumatic event and can lead to posttraumatic stress symptoms. For example, 12%-15% survivors of SARS showed one type of PTSD symptom (PTSS) at one month after discharge (Wu et al, 2005); the morbidity of PTSD in patients with traumatic injury reached 9.6% at months (Bryant et al., 2017). In addition, not only being attacked by traumatic illness could cause PTSD symptoms, being contacts and carers of a deadly virus could also lead to symptoms like alternations in mood and arousal (Tine et al., 2016). Living in the endemic area, every Chinese is a carer and a (possible) contact of COVID-19. Although having experienced the outbreak of Severe Acute Respiratory Syndrome (SARS), a severe infectious disease which had caused hundreds of death in China in 2003 (WHO, 2003), the traumatic course of the infection and the characteristic of human transmission of novel coronavirus (Li et al., 2020) were beyond expectation, causing fear, helplessness and PTSS among Chinese people. Hence in the present study, we focused on the acute PTSS in Chinese under the background of massive COVID-19 outbreak.

Previous studies have proved that massive outbreaks of epidemics like SARS and Ebola virus disease (EVD) could cause sleep problems in relevant individuals. For example, Yu et al. (2005) reported that midlife women felt “sleep was restless” during SARS and this problem was closely associated with their overall negative emotions. Another study showed that chronic post-SARS symptoms were characterized by sleep-related problems (Moldofsky and Patcai, 2011). Survivors of EVD also reported various sleep quality problems (Keita et al., 2017). More importantly, Sleep and PTSS are closely related, and sleep disturbances are core features of PTSD. According to the model raised by Richard et al. (2020), sleep disturbances (short sleep duration, frequent arousals, and et al.) could lead to maladaptive sleep-related compensatory behaviors and cause hyperarousal and anxiety-related disorders like PTSD. Sleep disruption also promoted anxious emotions by disrupt emotion regulation and adaptive memory functions. Thus, it was reasonable to hypothesize that wide-spread transmission of COVID-19 could causes sleep problems in Chinese and result in PTSS.

In early December 2019, the first pneumonia case was identified in Wuhan, the capital city of Hubei province (Huang et al., 2020). With the spreading of the epidemic, Wuhan became the worst influenced area, where the mortality and incidence rate of COVID-19 were above the national level according to the National Health Commission of China (2020). Moreover, not only Wuhan residents were more likely to be infected, 72.3% nonresidents of Wuhan had contact with individuals lived in Wuhan (Guan et al., 2020). Wuhan expose history had become an important

diagnosis standard and a source of fear. As a result, we collected data of Wuhan exposure to measure the risk of infection. We hypothesized that Wuhan residents and individuals who had been to Wuhan or contacted with Wuhan residents, would show more sleep problems and PTSS for they were faced with high possibility of infection of COVID-19. We also explored the relationship among the infection risk, sleep and PTSS to identify the role of sleep problems in the development of PTSD. We hope the current study could reflect the influence of COVID-19 on psychological well-being in Chinses and shed light on post-epidemic psychological intervention.

## **Methods**

The methods and reporting procedures were in accordance with the STROBE (STrengthening the Reporting of OBServational studies in Epidemiology) checklist.

### **Participants**

A total of 2032 subjects from 31 provinces in the mainland of China, Hong Kong, Macao and Taiwan participated in the present study. Criteria for inclusion was that subjects should be Chinese race and age  $\geq 18$  years at the time of the COVID-19 infection. Table 1 demonstrated demographic statistics of subjects. The average age of all subjects was  $35.47 \pm 11.32$ .

All the study participants were provided with written informed consent. This project was approved by the Ethics Committee of the Naval Military Medical University.

### **Wuhan expose history**

Subjects who were Wuhan residents, or had been to Wuhan, or had contacted with anyone from Wuhan during the outbreak of COVID-19 scored 1 point and those not scored 0. A total of 230 subjects showed a history of exposure to Wuhan.

### **Sleep quality**

According to previous findings, certain types of sleep quality disturbances were more common across anxiety-related disorders than others (Leskin et al., 2002; Neylan, 1998). As a result, we only selected the most related items from the Pittsburgh Sleep Quality Index (PSQI) (Buysse et al., 1989) to investigate sleep quality. 4 items from PSQI were used to measure subject's sleep quality. The first was the subjective sleep quality measuring participants' satisfaction with sleep. Subject scored 0 if they had very good subjective sleep quality and scored 3 for very bad subjective sleep quality. The second item collected data on latency onset of sleep. Scores varied from 0~3 with 0 standing for having no problems falling asleep and 3 standing for hard to fall asleep for 3 times or more within a week. The third item was the sleep fragment. Subjects score 1 for no occurrence of the problem and scored 3 if they had the problem for 3 times or more within a week. The last item collected data on everyday sleep duration and the scores varied from 0 (more than 7 hours) to 3 (less than 5 hours). In summary, 0 and 1 point of the 1<sup>st</sup> item indicated very good and good subjective sleep quality and 2 and 3 indicated sleep quality problems. For the remaining items, 0 point indicated good sleep quality and 1 to 3 points indicated increasingly severe sleep quality problems.

## PTSS

PTSD Checklist for DSM-5 (PCL-5) (Weathers et al., 2013) was used to measure PTSS. PCL-5 is a self-report 5-point Likert-Type scale (1=not at all, and 5=extremely) with 20 items measuring PTSS of instruction, avoidance, numbing and hyperarousal. The Chinese version of PCL-5 showed excellent consistency and convergent validity in previous studies (Li et al., 2018; Liu et al., 2014). Internal consistency (Cronbach's  $\alpha$ ) of PCL-5 in the present study was 0.87.

## Data collection

The data was collected during Jan 30, 2020 to Feb 3, 2020 through an online survey platform <https://www.wjx.cn/>. To complete the survey, participants had to answer all the listed questions in a time limitation of 2 to 30 minutes. Answers with response time below 2 mins or above 30mins were invalid and would be deleted in the following analysis.

## Data analysis

5 participants were deleted because of illegal response time. Data from 2027 participants were analyzed. One-way variance analysis (ANOVA) and Least-Significant Difference (LSD) test were used to analyze the differences of PTSS during each day of survey. Chi-square test was applied to analyzed differences of sleep quality by time and Wuhan-exposed history. Scores of PTSS were compared between subjects with and without Wuhan exposure history using *t*-test. Interaction effects of sleep and Wuhan exposure history on PTSS was analyzed using two-way ANOVAs and simple effect analyses.

## Results

### General results of sleep quality and PTSS

Figure 1 showed trends of PCL-5 scores and sleep quality during the 5-day survey. Both PCL-5 scores and sleep quality scores showed raising tendency from Jan 30, 2020 to Feb 2, 2020 and decreased in Feb 3, 2020. One-way ANOVA showed that there was a significant difference of PCL-5 scores on different dates,  $F=4.38$ ,  $p<0.05$ . LSD test showed significant increases was illustrated from Jan 31 to Feb 2 ( $p<0.05$ ); however, the PCL-5 score in Feb 3, 2020 were significantly lower than that in Feb 2, 2020 ( $p<0.05$ ).

Chi-square test showed significant differences of sleep quality in the 5-days period,  $\chi^2=12.33$ ,  $p<0.05$  for subjective sleep quality,  $\chi^2=34.32$ ,  $p<0.05$  for sleep fragment and  $\chi^2=20.03$ ,  $p<0.05$  for sleep duration. However, latency onset of sleep did not illustrate significant change during the period.

### Influence of Wuhan exposure history on sleep quality and PTSS

Results demonstrated that participants with Wuhan exposure history generally showed more latency onset of sleep ( $\chi^2=9.77$ ,  $p<0.05$ ) than those not. Wuhan exposure history also significantly influence PCL-5 scores ( $t=-2.93$ ,  $p<0.05$ ). However, there no significant differences of other sleep problems between participants with and without Wuhan-exposure history. See table 2 for detailed information.

### **Interaction of exposure history and sleep quality on PTSS**

Table 3 showed results of ANOVAs. To test the exposure history and sleep quality on PTSS, the mean values of PCL-5 entered into  $2 \times 2$  ANOVAs with exposure history and sleep quality as independent variables. Results showed a significant interaction effect of Wuhan exposure history  $\times$  subjective sleep quality on PCL-5 ( $F=4.01, p<0.01, \eta_p^2=0.006$ ). Simple effect tests indicated that there was a significant main effect of Wuhan exposure history on PCL-5 in subjects with relatively poorer subject sleep quality (scored 2 and 3 points in subject sleep quality), but not in subjects with better subject sleep quality. Results also demonstrated interaction of Wuhan exposure history  $\times$  latency onset of sleep ( $F=3.67, p<0.05, \eta_p^2=0.006$ ), but Wuhan-exposure only significantly affected PTSS in those showed severe problems falling asleep (score 3 points in latency onset of sleep). Interactions of Wuhan exposure history  $\times$  sleep fragment ( $F=3.14, p<0.05, \eta_p^2=0.006$ ) and Wuhan exposure history  $\times$  sleep duration ( $F=3.65, p<0.05, \eta_p^2=0.005$ ) were significant. Results of simple effect tests were similar: there were only a main effect of Wuhan exposure history in subjects who were easier to wake up and had shorter sleep duration (scored 3 and 4 points in easily waking during the night or too early in the morning, and in sleep duration). These results indicated that sleep quality regulated the relationship between Wuhan exposure history with sleep and PTSS.

### **Discussion**

According to National Health Commission of China (NHC), until Feb 4, 2020, prevalence rate and fatality rate of COVID-19 in Wuhan were much high above the national level (NHC, 2020), making exposure to Wuhan a risk factor of infection and a cause of mental health problem. The current study aimed to investigate the influence of potential danger of infection on sleep and PTSS. We also explored regulating effects of sleep quality on the relationship between Wuhan exposure history during outbreak of COVID-19 and PTSS. To this end, we conducted ANOVAs and Chi-square tests with the above variables.

Results indicated that data of PTSS growth rapidly in the first 4 days in our study, and decreased in the last day. Interestingly, scores of sleep quality showed the same tendency. This might indicate the close relationship between sleep and PTSD. Moreover, up to the end of our survey, the firm cases and susceptible cases were still frantically increased, showing no sign of decline (WHO, 2020e). We assumed that results of the present study suggested that emergence of psychological “inflection point” which meant a turning from deterioration of mental health to growth of resilience showed on Feb 3, 2020. It was a date before the number of confirmed cases of COVID-19 showed any sign of downward trends in China. We supposed that the change in psychological response might be a result of Chinese government’s efforts in disease management, psychoeducation and prevention education.

Results illustrated that Wuhan exposure history affected PTSS. For those having Wuhan-exposure experiences, PTSS were more obvious. This result was in line with previous findings that subjective perception of dangerousness during SARS related with severity of PTSD

symptoms (Mak et al., 2010). Results also highlighted the influence of trauma exposure on latency onset of sleep, which was different in subjects with and without Wuhan exposure history. Difficulty in falling asleep might be a result of increased presleep cognitions. According to Fast (2003), traumatic experiences caused reactions of fear, increased arousal and re-experience, and these reactions affected presleep cognitions and led to changes of sleep onset latency. Increased sleep onset latency then influenced mood, cognition and physiological responses.

We confirmed that subjective sleep quality, delayed sleep onset, sleep fragment and sleep duration all regulated the effect of Wuhan exposure history on PTSS. Wuhan-exposure history influenced only those with relatively poor sleep quality, but if individuals showed few or none sleep problems, Wuhan-exposure was not a risk factor of PTSS. These results indicated that traumatic events provide bases for PTSD but one's ability to keep sleep quality was a protective factor of onset and development of PTSD. This was line with Germain et al.'s (2016) model of sleep and PTSD which indicated that sleep related more closely with PTSS than trauma events. These results were also consistent with previous studies which showed alters of REM sleep after uncontrolled stress accounted for the long term physiological and behavioral impairments in an animal model (Vanderheyden et al., 2015). The mechanisms regulating the interactions of traumatic stress and sleep were poorly understood, and we extended previous findings by showing regulation effect of sleep quality on the relationship between trauma exposure and PTSD in human samples. To prevent mental illness and protect mental health in the peak rush of COVID-19 and other infectious fatal diseases, health authorities should be mindful of the influence of infectious risks on psychological reactions, propagate the importance of keeping good sleep quality like enough sleep time, keeping early hours etc.

Limitations of the current study included the use of a health example and the fact that we used online survey in data collection. Only a few subjects were confirmed or suspected COVID-19 patients in our study. And the mean levels of sleep quality and PTSS were in the moderate range. For those being infected with COVID-19, sleep and mental health problems might be far more severe. Considering the risk of infection, we used online survey to investigate relationship among Wuhan exposure history, sleep and PTSS. However, it was hard to guarantee authenticity with this method and the validity was limited. We suggested that future studies could examine PTSD and related factors among COVID-19 survivors with face-to-face investigation or interview.

## **Conclusion**

The present study suggested that sleep quality level decreased and posttraumatic stress symptoms increased with time and change of this tendency was shown on 3<sup>rd</sup> Feb, 2020. Individuals with and without Wuhan-exposure history showed different levels of PTSS and sleep onset latency under the influence of COVID-19. Wuhan-exposure history influenced PTSS severity in individuals with poor sleep quality, but not those with high-quality sleep.

## Reference

American Psychiatric Association, 2013. Diagnostic and Statistical Manual of Mental Disorders, 5th ed. American Psychiatric Publishing, Arlington, VA.

Bryant, R.A., Creamer, M., O'Donnell, M., et al. Acute and Chronic Posttraumatic Stress Symptoms in the Emergence of Posttraumatic Stress Disorder A Network Analysis. *Jama Psychiat*. 2017;74(2):135-142. DOI: 10.1001/jamapsychiatry.2016.3470.

Buysse, D.J., III Reynolds, C.F., Monk ,T.H., Berman, S.R., Kupfer, D.J. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiat Res*. 1989;28(2):193-213. DOI: DOI: 10.1016/0165-1781(89)90047-4.

Fast, K.M. The relationship between presleep cognitions and delayed sleep-onset in a sample of women seeking treatment for sexual assault. 2003, University of Hawai'i; Hilo.

Germain, A., McKeon, A.B., Campbell, R.L. Sleep in PTSD: Conceptual model and novel directions in brain-based research and interventions. *Curr Opin Psychol*. 2016;14:84-89. DOI: 10.1016/j.copsyc.2016.12.004.

Gibson, C.J., Richards, A., Villanueva, C., Barrientos, M., Neylan, T.C., Inslicht, S.S. Subjective Sleep Related to Post Traumatic Stress Disorder Symptoms among Trauma-Exposed Men and Women. *Behav Sleep Med*. 2019, 17(4):492-501. DOI: 10.1080/15402002.2017.1409223.

Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*, 2020. DOI: 10.1056/NEJMoa2002032.

Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497-506. DOI: 10.1016/S0140-6736(20)30183-5.

Keita, M.M , Taverne, B. , Sy Savané, S., et al. Depressive symptoms among survivors of Ebola virus disease in Conakry (Guinea): preliminary results of the PostEboGui cohort. *BMC Psychiatry*, 2017, 17(1):127. DOI: 10.1186/s12888-017-1280-8.

Leskin, G.A., Woodward, S.H., Young, H.E., Sheikh, J.I. Effects of comorbid diagnoses on sleep disturbance in PTSD. *J Psychiatr Res*. 2002;36(6):449-452. DOI: 10.1016/s0022-3956(02)00025-0.

Li, G., Li, W., Cao, C., et al. DSM-5 posttraumatic stress symptom dimensions and health-related quality of life among Chinese earthquake survivors. *Euro J Psychotraumatol*, 2018, 9(1), 1468710. DOI: 10.1080/20008198.2018.1468710.

Li, Q., Guan, X., Wu, P., et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl j med*. 2020, in press. DOI: 10.1056/NEJMoa2001316.

Liu, P., Wang, L. , Cao, C., et al. The underlying dimensions of DSM-5 posttraumatic stress disorder symptoms in an epidemiological sample of Chinese earthquake survivors. *J anxiety disorders*, 2014, 28(4):345-351. DOI: 10.1056/NEJMoa2001316.

Mak, I.W.C., Chu, C.M., Pan, P.C., Yiu, M.G.C., Ho, S.C., Chan, V.L.. Risk factors for

chronic post-traumatic stress disorder (PTSD) in SARS survivors. *Gen Hosp Psychiatry*. 2010;32(6):590-598. DOI: 10.1016/j.janxdis.2014.03.008.

Moldofsky H, Pstcai J. Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome; a case-controlled study. *BMC Neurology*, 2011,11:37. DOI: 10.1186/1471-2377-11-37.

National health commission of the People's Republic of China. National health commission press conference on February 4, 2020, (<http://www.nhc.gov.cn/xwzb/webcontroller.do?titleSeq=11220&gecstype=1>). Accessed on Feb 4, 2020.

National Health Commission of the People's Republic of China. Home page (<http://www.nhc.gov.cn>). Accessed on Mar 5, 2020.

Neylan, T.C. Marmar, C.R., Metzler, T.J., et al. Sleep disturbances in the Vietnam generation: findings from a nationally representative sample of male Vietnam Veterans. *Am J Psychiatry*. 1998;155:929-933. DOI: 10.1176/ajp.155.7.929.

Richards, A., Kanady, J.C., Neylan, T.C. Sleep disturbance in PTSD and other anxiety-related disorders: an updated review of clinical features, physiological characteristics, and psychological and neurobiological mechanisms. *Neuropsychopharmacol*. 2020;45(1):55-73. DOI: 10.1038/s41386-019-0486-5.

Bortel, T.V., Basnayake, A., Wurie, F., Jambai, M., Nellums, L.B. Psychosocial effects of an Ebola outbreak at individual, community and international levels. *Bull World Health Organ*. 2016;94(3):210-214. DOI: 10.2471/BLT.15.158543.

Weathers, F.W., Litz, B.T., Keane, T.M., Palmieri, P.A., Marx, B.P., Schnurr, P.P. (2013). The PTSD checklist for DSM-5 (PCL-5) Scale. Available from the National Center for PTSD. <http://www.ptsd.va.gov>.

WHO. (2003). Cumulative number of reported probable cases of SARS: From 1 Nov 2002 to 11 July 2003. Geneva, Switzerland: World Health Organization.

WHO. Novel Coronavirus (2019-nCov) situation report-1. 2020a. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-n-cov.pdf?sfvrsn=20a99c10\\_4](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-n-cov.pdf?sfvrsn=20a99c10_4).

WHO. Statement on the second meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). 2020b. [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)). Accessed on Feb 4, 2020.

WHO. Novel Coronavirus (2019-nCov) situation report-15. 2020c. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200204-sitrep-15-ncov.pdf?sfvrsn=88fe8ad6\\_4](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200204-sitrep-15-ncov.pdf?sfvrsn=88fe8ad6_4).


WHO. Novel Coronavirus (2019-nCov) situation report-14. 2020d.

[https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200204-sitrep-14-ncov.pdf?sfvrsn=88fe8ad6\\_4](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200204-sitrep-14-ncov.pdf?sfvrsn=88fe8ad6_4).

Wu, K.K., Chan, S.K., Ma, T.M. Posttraumatic Stress after SARS. *Emerg Infect Dis.* 2005;11(8):1297-1300. DOI: 10.3201/eid1108.041083

Vanderheyden, W.M., George, S.A., Urpa, L., Kehoe, M., Liberzon, I., Poe, G.R. Sleep alterations following exposure to stress predict fear-associated memory impairments in a rodent model of PTSD. *ExpBrain Res.*, 2015, 233(8): 2335-46. DOI: 10.1007/s00221-015-4302-0.

Yu, H.Y.R., Ho, S.C., So, K.F.E., Yin, L.L. The psychological burden experienced by Hong Kong midlife women during the SARS epidemic. *Stress Health.* 2005;21(3):177-184. DOI: 10.1002/smi.1051.



**Figure 1** Changes of PCL-5 scores and participants with sleep quality problems in the 5-day survey

**Table 1** Demographic Statistics,exposure history and sleep status of all subjects

	N	%
<b>Gender</b>		
Male	786	38.8
Female	1241	61.2
<b>Education Level</b>		
High school or below	254	12.5
University or college	1310	64.6
Postgraduate or above	463	22.8
<b>Exposure history of Wuhan</b>		
No	1797	88.7
Yes	230	11.3
<b>Subjective sleep quality</b>		
Very good	816	40.3
Good	859	42.4
Poor	320	15.8
Very poor	32	1.6
<b>Latency onset of sleep</b>		
No	1358	67.0
<Once a week	312	15.4
once or twice a week	251	12.4
>=Three times a week	106	5.2
<b>Sleep fragment</b>		
No	1133	55.9
<Once a week	312	15.4
once or twice a week	368	18.2
>=Three times a week	214	10.6
<b>Sleep duration</b>		
>7h	996	49.1
6-7h	645	31.8
5-6h	314	15.5
<5h	72	3.6
<b>Total</b>	2027	100.0

**Table 2 Sleep quality and PCL -5 score by Wuhan exposure history**

	Exposure history of Wuhan				$\chi^2$	p-value
	No(n=1997)		Yes(n=230)			
	N	%	N	%		
<b>Subjective sleep quality</b>						
very good	724	40.3%	92	40.0%	6.643	0.084
good	768	42.7%	91	39.6%		
poor	281	15.6%	39	17.0%		
very poor	24	1.3%	8	3.5%		
<b>Latency onset of sleep</b>						
No	1212	67.4%	146	63.5%	9.768	0.021
<Once a week	282	15.7%	30	13.0%		
once or twice a week	218	12.1%	33	14.3%		
>=Three times a week	85	4.7%	21	9.1%		
<b>Sleep fragment</b>						
No	1001	55.7%	132	57.4%	3.996	0.262
<Once a week	286	15.9%	26	11.3%		
once or twice a week	325	18.1%	43	18.7%		
>=Three times a week	185	10.3%	29	12.6%		
<b>Sleep duration</b>						
>7h	872	48.5%	124	53.9%	3.861	0.277
6-7h	584	32.5%	61	26.5%		
5-6h	276	15.4%	38	16.5%		
<5h	65	3.6%	7	3.0%		
	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>t</b>	<b>p-value</b>
<b>PCL-5 Score</b>	11.49	9.981	14.00	12.546	-2.925	0.004

Note. PCL-5=PTSD Checklist-Civilian Version.

**Table 3 Interaction of Wuhan exposure history and sleep quality on PCL-5**

Source	Type III Sum of Squares	df	Mean Square	F	p-value
Exposure in Wuhan	1193.48	1	1193.48	14.321	<0.001
Subjective sleep quality	29974.03	3	9991.34	119.891	<0.001
Exposure in Wuhan × Subjective sleep quality	1003.67	3	334.56	4.014	0.007
Error	168257.11	2019	83.34		
Total	497244.00	2027			
Exposure in Wuhan	1007.27	1	1007.27	11.762	0.001
Unable to fall asleep within 30 minutes	26266.76	3	8755.59	102.239	<0.001
Exposure in Wuhan × Latency onset of sleep	942.36	3	314.12	3.668	0.012
Error	172903.73	2019	85.64		
Total	497244.00	2027			
Exposure in Wuhan	1072.77	1	1072.77	12.035	0.001
Easily waking during the night or too early in the morning	19371.99	3	6457.33	72.441	<0.001
Exposure in Wuhan × Easily waking during the night or too early in the morning	840.49	3	280.16	3.143	0.024
Error	179972.74	2019	89.14		
Total	497244.00	2027			
Exposure in Wuhan	1547.88	1	1547.88	15.855	<0.001
Sleep duration	10966.04	3	3655.35	37.443	<0.001
Exposure in Wuhan × Sleep duration	1070.30	3	356.77	3.654	0.012
Error	197106.05	2019	97.63		
Total	497244.00	2027			