


Journal Pre-proof

Wuhan and Hubei COVID-19 mortality analysis reveals the critical role of timely supply of medical resources

Zuqin Zhang , Wei Yao , Yan Wang , Cheng Long , Xinmiao Fu

PII: S0163-4453(20)30145-6
DOI: <https://doi.org/10.1016/j.jinf.2020.03.018>
Reference: YJINF 4492



To appear in: *Journal of Infection*

Accepted date: 14 March 2020

Please cite this article as: Zuqin Zhang , Wei Yao , Yan Wang , Cheng Long , Xinmiao Fu , Wuhan and Hubei COVID-19 mortality analysis reveals the critical role of timely supply of medical resources, *Journal of Infection* (2020), doi: <https://doi.org/10.1016/j.jinf.2020.03.018>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd on behalf of The British Infection Association.

Highlights

- COVID-19 mortality rates in Wuhan and Hubei decay exponentially
- COVID-19 mortality rates outside Hubei and Wuhan are nearly constant.
- COVID-19 recovery rates in Wuhan and Hubei grow exponentially.
- Over 40000 aided health workers help Hubei to effectively treat patients.
- Newly supplied beds allow over 38000 patients in Wuhan to be treated in hospitals.

Journal Pre-proof

Letter to Editor

Wuhan and Hubei COVID-19 mortality analysis reveals the critical role of timely supply of medical resources

Title Page

Wuhan and Hubei COVID-19 mortality analysis reveals the critical role of timely supply of medical resources

Zuqin Zhang^{1,#}, Wei Yao^{1,#}, Yan Wang¹, Cheng Long^{2,*} and Xinmiao Fu^{1,*}

¹ Provincial University Key Laboratory of Cellular Stress Response and Metabolic Regulation, College of Life Sciences, Fujian Normal University, Fuzhou City, Fujian Province 350117, China

² Department of Orthopaedic, Sichuan University West China Hospital, Chengdu City, Sichuan Province, China

Running title: Resource-driven COVID-19 treatment in China

These authors contributed to this work equally.

* To whom correspondence should be addressed to Professors Xinmiao Fu (xmfu@fjnu.edu.cn) and Dr. Cheng Long (longchenghx@126.com)

Keywords: COVID-19, SARS-CoV-2, coronavirus, epidemic, mortality rate

The 2019 novel coronavirus diseases (COVID-19) outbreak caused by SARS-CoV-2 is on-going in China and has hit many countries¹⁻³. As of 3 March 2020, there have been 80270 confirmed cases and 2981 deaths in China, most of which are from the epicenter of the outbreak, Wuhan City, the capital of Hubei Province. New COVID-19 cases have been steadily declining in China and more than 60000 patients have been recovered⁴, largely due to the effective implementation of comprehensive control measures in China^{5,6}. Here we report that some of these measures, such as a dramatic and timely increase of medical supplies, may play a critical role such that the mortality and recovery rates of COVID-19 in Wuhan follow exponential

decay and growth modes, respectively.

We collected data for analysis on the officially released cumulative numbers of confirmed, dead and recovered cases (from 23 Jan to 3 Mar 2020) in five geographic regions, i.e., mainland China, Hubei Province, outside Hubei (in China), Wuhan City and outside Wuhan (in Hubei). As of 3 Mar 2020, crude fatality ratios (CFRs) in the above regions are 0.027 ± 0.006 , 0.035 ± 0.007 , 0.005 ± 0.002 , 0.045 ± 0.012 and 0.021 ± 0.008 , respectively, in line with earlier reports^{5,6}. While the mortality rates of COVID-19 outside Hubei and outside Wuhan appear constant over time, the mortality rates in Hubei and Wuhan decline continuously (**Fig. 1A**). Strikingly, the mortality rates in Hubei and Wuhan are well-fitted with the exponential decay mode (R^2 being 0.93 and 0.82, respectively; **Fig. 1A** and **Table S1**), and it is the same forth with that in China (R^2 being 0.86) but not with that outside Hubei and outside Wuhan (R^2 being 0.39 and 0.32, respectively). Remarkably, we found that the recovery rates of COVID-19 patients in the above regions were all well-fitted with the exponential growth mode (R^2 being 0.96, 0.95, 0.95, 0.88 and 0.95, respectively; **Fig. 1B** and **Table S1**). Such intriguing pattern for the COVID-19 mortality and recovery rates in Wuhan (or Hubei) somehow contradicts traditional epidemiological models wherein both are assumed as constants⁷.

The above unique pattern may reflect the fact that COVID-19 patients in Wuhan (or Hubei) have been treated more effectively day by day. Here we focused on two components essential for effective treatments, i.e., the supply of health workers and hospital beds. As a matter of fact, a great number (up to 42 000, as of 1 March 2020) of health workers have been aided by other provinces in China (**Table S2**) and they are working in different cities of Hubei (**Table S3**). This extraordinary aid keeps the ratio of the health workers to patients in Hubei at above 0.6 despite the number of remaining confirmed cases has ten-fold increased up to 50 000 on 18 Feb (**Fig. 1C**). Results also show that the number of acute care beds from more than 45 designated hospitals plus two newly built ones in Wuhan has been consecutively increasing up to 23 532 (as of Feb 24) under the government-directed re-allocation (**Fig. 1D**). This supply thus enabled the severe and critical patients to be treated timely and effectively⁶. More importantly, there have been over 10 temporary hospitals (named Fangcang hospitals) reconstructed from gymnasium and exhibition centers, which provide more than 26 000 makeshift beds for mild patients (**Fig. 1D**). These combinations guaranteed nearly 100% percent of COVID-19 patients to be treated in hospitals even if the number of remaining confirmed cases has ten-fold increased up to 38 000 on 18 Feb (**Fig. 1D**). In contrast, a lot of patients had to stay at home in the early stage of the outbreak in Wuhan due to the shortage of beds such that many

transmissions in households occurred ⁶.

Accordingly, the effective implementation of comprehensive control measures and infection-treatment practices is critical for combatting any new pathogens, not only interrupting the transmissions but also saving the patients. Timely supplied medical resources, including re-allocation of acute care beds, rapid construction of new hospitals and generous aid of health workers by other less-severe areas, apparently help the epicenter of the outbreak Hubei (Wuhan) to accomplish a unique and also encouraging outcome for life-saving such that the mortality and recovery rates of nearly 50 000 COVID-19 patients exponentially decays and grows, respectively. Other crucial factors contributing to this success may include the improved and optimized diagnosis and treatment strategies ⁸, which are critical for saving severe and critical patients ^{5, 6, 9}. This speculation appears to be supported by the exponential growth of the COVID-19 recovery rate outside Hubei (**Fig. 1B**) where medical resources are relatively sufficient ¹⁰. Collectively, the achievement made in Hubei (or Wuhan) may provide useful guidance for many countries to be better prepared for the potential pandemic ² that may overwhelm local health care systems.

Acknowledgments

This work is support by the National Natural Science Foundation of China (No. 31972918 and 31770830 to XF). All authors report no conflicts of interest relevant to this article.

References

1. Tang, J.W., P.A. Tambyah, and D.S.C. Hui, Emergence of a novel coronavirus causing respiratory illness from Wuhan, China. *Journal of Infection*, 2020; <https://doi.org/10.1016/j.jinf.2020.01.014>.
2. WHO: *Coronavirus disease (COVID-19) outbreak*. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
3. Zhou, P.e.a., A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, 2020; <https://doi.org/10.1038/s41586-020-2012-7> (2020).
4. *the National Health Commission of China, Real-time Data of the COVID-19 Outbreaks in China (in Chinese)*, http://www.nhc.gov.cn/xcs/yqfkdt/qzbd_index.shtml.
5. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. Vital surveillances: the epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19)—China, 2020, <http://weekly.chinacdc.cn/en/article/id/e53946e2-c6c4-41e9-9a9b-fea8db1a8f51>, in *China CDC Weekly*.
6. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19), <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>. 2020.
7. Wang, X.S., J. Wu, and Y. Yang, Richards model revisited: Validation by and application to infection dynamics. *Journal of Theoretical Biology*, 2012. 313 12–19.
8. *the National Health Commission of China, Coronavirus Disease 2019 (COVID-19) diagnosis and treatment plan (in Chinese, 7th version)*. Available from: <http://www.nhc.gov.cn/yzygj/s7653p/202003/46c9294a7dfe4cef80dc7f5912eb1989.shtml>.

9. Yang, X., et al., Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*, 2020; [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
10. Ji, Y., et al., Potential association between COVID-19 mortality and healthcare resource availability. *Lancet Global Health*, 2020; [https://doi.org/10.1016/S2214-109X\(20\)30068-1](https://doi.org/10.1016/S2214-109X(20)30068-1).

Figure 1




Fig. 1 Fitting the COVID-19 mortality and recovery rates with exponential decay and growth functions, respectively.

(A, B) Mortality rate (panel A) and recovery rate (panel B) for COVID-19 in China over time and by location, with exponential decay- and growth-based regression analyses being performed, respectively (as shown by colored solid lines). Parameters from the regression analyses are shown in **Table S1**, with R^2 being shown here. (C) Numbers of the aided health workers in Hubei over time. Ratio of the aided health workers to patients was also plotted (note: most of the aided health workers are working in Wuhan; refer to **Table S3**). (D) Numbers of the remaining confirmed cases of COVID-19, and acute care beds, makeshift beds from Fangcang hospitals and total beds in Wuhan over time. Ratio of beds to patients was also plotted. Here the data of newly supplied beds in Hubei are not available and thus the data for Wuhan were analyzed.

Graphic abstract

