

RECOMMENDATIONS

- Close monitoring of all PICU patients especially ventilated cases for detection of early signs of pneumothorax before progression to life threatening tension pneumothorax.
- Strict application of protective lung strategies in order to minimize occurrence of pneumothorax in mechanically ventilated cases.
- Further use of bedside ultrasonography in diagnosis of pneumothorax due to its high sensitivity and specificity.
- Use of bedside ultrasonography guidance during insertion of central lines to minimize pneumothorax and other complications secondary to CVC insertion.

REFERENCES

1. Kliegman RM, Behrman RE, Stanton BF, Jenson HB. Pneumothorax. In: Winnie GB, editors. Nelson textbook of paediatrics. 19th edition. Philadelphia :Saunders Elsevier 2011:1835-37.
2. Parthasarathy A , Borker AS, Nair MKC. Pneumothorax. In: Raichur DV, editors. Partha's Fundamentals of paediatrics. 2nd edition. New Delhi :Jaypee Brothers Medical Publishers 2013 :278-9
3. Jantz MA, Anthony VB. Pathophysiology of the pleura. Respiration 2008;75(2):121-133.
4. Noppen M, Schramel F. Pneumothorax . Eur Respir Mon 2002;22:279-96
5. Baumann MH, Noppen M . Pneumothorax. Respirology 2004 ;9: 157-64.
6. Rimensberger PC. Indications of non- conventional ventilation modes. IN: Bollen C, Rimensberger PC, editors. Pediatric and neonatal mechanical ventilation. From basics to clinical practise. New York: Springer 2015:562.
7. Eckstein M, Henderson S, Markovchick V .Chapter 38- Thorax. In: Marx JA , editor. Rosen's Emergency Medicine : Concepts and clinical practice, 5th edition. 2002,St Louis: Mosby, pp386-91.
8. Shaw KS, Prasil P, Nguyen LT, Laberge J. Pediatric spontaneous pneumothorax. Semin Pediatr Surg 2003;12:55-61.
9. Poenaru D, Yazbeck S, Murphy S. Primary spontaneous pneumothorax in children. J Pediatr Surg 1994;9:1183-85.
10. Robinson PD, Cooper P, Ranganthan SC. Evidence-based management of pediatric primary spontaneous pneumothorax . Pediatr Respir Rev 2009;10:110-7.
11. Wilcox DT, Glick PL, Karamanoukian HL, Allen JE, Azizkhan RG . Spontaneous pneumothorax: a single institution, a 12 year experience in pneumothorax under 16 years of age. J Pediatr Surg 1995;30(10):1451-4.
12. Damore DT, Dayan PS. Medical causes of pneumomediastinum in children. Clin Pediatr (Phila) 2001;40(2):87-91.
13. Hafen GM, Unkomunne OC, Robinson PJ. Pneumothorax in cystic fibrosis: a retrospective case series. Arch Dis Child 2006;91(11):924-25.
14. Sahn Sa, Heffner JE, .Spontaneous pneumothorax. N Engl J Med 2000;342(12):868-74.
15. Posner K , Needleman JP. Pneumothorax. Pediatr Rev 2008;29(2);69-70.
16. Bliss D, Silen M. Pediatric thoracic trauma. Crit Care Med 2002; 30(supp11):s409-s415.

References

17. Celik B, Sahin E, Nadir A , Kaptanoglu M. Iatrogenic pneumothorax: etiology, incidence and risk factors. *Thorac Cardiovasc Surg* 2009;57(5):286-90.
18. Askegard-Giesmann JR, Caniano DA, Kenny BD. Rare but serious complications of central line insertion. *Semin Pediatr Surg* 2009;18(2):73-83.
19. Gordon CE, Feller-kopman D, Balk EM, Smetana GW. Pneumothorax following thoracocentesis : a systemic review and meta-analysis. *Arch Intern Med* 2010;170(4):332-9.
20. Kumar A, Chuan A. Ultrasound guided vascular access efficacy and safty. *Best Pract Res Clin Anaesthesiol* 2009;23(3):299-311.
21. Duncan DR, Morgenthaler TI, Ryu JH, Daniels CE. Reducing iatrogenic risk in thoracocentesis : establishing best practice via experiential training in a zero –risk environment. *Chest* 2009;135(5):1315-20.
22. Sigaut S, Skhiri A, Stany I, Golmar J, Nivoche Y, Constant I, et al. Ultrasound guided internal juglar vein access in children and infant: a meta-analysis published studies. *Pediatr Anaesth* 2009;19(12):1199-206.
23. Bagchi I, Nycyk JA. Familial spontaneous pneumothorax. *Arch Dis Child Fetal Neonatal Ed.* 2002 jul; 87(1): F70.
24. Wait MA, Estrera A. Changing clinical spectrum of spontaneous pneumothorax .*Am J Surg* 1992;164:528-31.
25. Parikh DH, Crabbe DC, Auldish AW, Rothenberg SS. Chapter 38:Pneumothorax.In: Bradnock TJ, Crabbe DC. *Pediatric Thoracic Surgery* .1st edition .London: Springer 2009:465-79.
26. Currie GP, Alluri R, Christie GL, Legge JS. Pneumothorax : an update. *Postgrad Med J* 2007;83:461-5.
27. Rankine JJ, Thomas AN, Fluechter D. Diagnosis of pneumothorax in critically ill adults .*Postgrad Med J* 2000;76:399-404.
28. Yarmus L, Feller-kopman D. Pneumothorax in critically ill patient. *Chest* 2012;141:1098-105.
29. Leigh-Smith S, Harris T. Tension pneumothorax –time for a re-think. *Emerg Med J* 2005;22:8-16.
30. Norris RM, Jones JG, Bishop JM. Respiratory gas exchange in patients with spontaneous pneumothorax. *Thorax* 1968;23:427-33.
31. Light RW, O'Hara VS, Moritz TE, McElhinney AJ, Butz R, Haakenson CM, et al. Intrapleural tetracycline for the prevention of recurrent spontaneous pneumothorax ;results of a Department of Veterans Affairs Co-Operative study. *JAMA* 1990;264:2224-30.

References

32. Lesur O, Delorme N, Frogamet JM, Bernadac P, Polu JM. Computed tomography in the aetiological assessment of idiopathic spontaneous pneumothorax. *Chest* 1990;98:341-7.
33. Tocino IM, Miller MH, Fairfax WR. Distribution of pneumothorax in the supine and semirecumbent critically ill adult. *AJR Am J Roentgenol* 1985;144:901-5.
34. Rhea JT, van Sonnenberg E, McLoud TC. Basilar pneumothorax in the supine adult. *Radiology* 1979;133:593-5.
35. Schramel FM, Golding RP, Haakman CD, Sutedja TG, de Jong KA, Postmus PE. Expiratory chest radiographs do not improve visibility of small apical pneumothoraces by enhanced contrast. *Eur Respir J* 1996;9:406-9.
36. Carr JJ, Reed JC, Choplin RH, Case LD. Pneumothorax detection: a problem in experimental design. *Radiology* 1993;186:23-6.
37. Ding W, Shen Y, Yang J, He X, Zhang M. Diagnosis of pneumothorax by radiography and ultrasonography: a meta-analysis. *Chest* 2011;140(4):859-66.
38. Brook OR, Beck-Razi N, Abadi S, Filatov J, Litmanovich D, Gaitini D, et al. Sonographic detection of pneumothorax by radiology residents as part of extended focused assessment with sonography for trauma. *J Ultrasound Med* 2009;28:749-55.
39. Paramasivam E, Bodenham A. Air leaks ,pneumothorax and chest drains . *Contin Educ Anaesth Crit Care Pain* 2008;8:204-9.
40. Soldati G, Testa A, Sher S, Pignataro G, La Sala M, Silveri NG. Occult traumatic pneumothorax: diagnostic accuracy of lung ultrasonography in the emergency department. *Chest* 2008;133:204-11.
41. Galbois A, Ait-Oufella H, Baudel JL, Kofman T, Bottero J, Viennot S, et al. Pleural ultrasound compared with chest radiographic detection of pneumothorax resolution after drainage. *Chest* 2010;138(3):648-55.
42. Reissig A, Kroegel C. Accuracy of transthoracic sonography in excluding post-interventional pneumothorax and hydropneumothorax. Comparison to chest radiography. *Eur J Radio*. 2005;53(3):463-70.
43. Vezzani A, Brusasco C, Palermo S, Launo C, Mergoni M, Corradi F. Ultrasound localization of central vein catheter and detection of postprocedural pneumothorax: an alternative to chest radiography. *Crit Care Med* 2010;38(2):533-8.
44. Mayo PH, Goltz HR, Tafreshi M, Doelken P. Safety of ultrasound-guided thoracentesis in patients receiving mechanical ventilation. *Chest* 2004;125(3):1059-62.
45. Kelly AM, Weldon D, Tsang AY, Graham CA. Comparison between two methods for estimating pneumothorax size from chest X-rays. *Respir Med* 2006;100(8):1356-9.

References

46. Phillips GD, Trotman-Dickenson B, Hodson ME, Geddes DM. Role of CT in the management of pneumothorax in patients with complex cystic lung disease. *Chest* 1997;112(1):275-8.
47. Kelly AM, Loy J, Tsang AY, Graham CA. Estimating the rate of re-expansion of spontaneous pneumothorax by a formula derived from computed tomography volumetry studies. *Emerg Med J* 2006;23(10):780-2.
48. Rahman NM, Davies RJO, Gleeson FV. Pleural interventions: management of acute and chronic pneumothorax. *Semin Respir Crit Care Med* 2008;29(4):427-40.
49. Sahn SA, Heffner JE. Spontaneous pneumothorax. *N Engl J Med* 2000;342(12):868-74.
50. Chen KY, Jerng JS, Liao WY, Ding LW, Kuo LC, Wang JY, et al. Pneumothorax in the ICU: patient outcomes and prognostic factors. *Chest* 2002;122(2):678-83.
51. Moore FO, Goslar PW, Coimbra R, Velmahos G, Brown CVR, Coopwood TB, et al. Blunt traumatic occult pneumothorax: is observation safe? Results of a prospective, AAST multicenter study. *J Trauma* 2011;70(5):1019-25.
52. Northfield TC. Oxygen therapy for spontaneous pneumothorax. *Br Med J* 1971;4:86-8.
53. Eber E, Midulla F. Pneumothorax and pneumomediastinum. In: Schwerk N, Brinkmann F, Grasemann H editors. *ERS Handbook of Pediatric Respiratory Medicine*. 1st edition. Charlesworth press 2013:485-90.
54. Massard G, Thomas P, Wihlm JM. Minimally invasive management for first and recurrent pneumothorax. *Ann Thorac Surg* 1998;66:592-9.
55. Rodriguez –Panadero F, Antony VB. Pleurodesis: state of art. *Euro Respir J* 1997;10:1648-54.
56. Henry M, Arnold T, Harvey J. Pleural Diseases Group. Standards of Care Committee, British Thoracic Society. *BTS guidelines for the management of spontaneous pneumothorax*. *Thorax* 2003; 58(suppl2): 39-52.
57. Carr JJ, Reed JC, Choplin RH, Pope TL Jr, Case LD. Plain and computed radiography for detecting experimentally induced pneumothorax in cadavers: implications for detection in patients. *Radiology* 1992; 183:193-99.
58. Brant WE, Helms CA. Pneumothorax. In: Klein JS and Ghostin JS editors *Fundamentals of diagnostic radiology*. 4th edition. Lippincott Williams & Wilkins, a Wolters Kluwers business, 2012:510-12.
59. Strange C. Pleural complications in the intensive care unit. *Clin Chest Med* 1999;20(2):317-27.
60. de Latorre FJ, Tomasa A, Klamburg J, Leon C, Soler M, Ruis J. Incidence of pneumothorax and pneumomediastinum in patients with aspiration pneumonia requiring ventilator support. *Chest* 1977; 72(2):141-44.

References

61. Rivera R, Tibballs J. Complications of endotracheal intubation and mechanical ventilation in infants and children. *Crit Care Med* 1992; 20(2):193-99.
62. de Lassence A, Timsit JF, Tafflet M, Azoulay E, Jamali S, Vincent F, et al. Pneumothorax in the intensive care unit: incidence, risk factors and outcome. *Anaesthesiology* 2006;104(1):5-13.
63. MacDuff A, Arnold A, Harvey J. Management of spontaneous pneumothorax: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010; suppl 2:ii 18-31.
64. Petersen GW, Baier H. Incidence of pulmonary barotrauma in a medical ICU. *Crit Care Med* 1983;11(2):67-9.
65. Mehta NM, Arnold JH. Mechanical ventilation in children with acute respiratory failure. *Curr Opin Crit Care* 2004;10(1):7-12.
66. Randolph AG, Meert KL, O'Neil ME, Hanson JH, LUCKETT PM, Arnold JH, et al. The feasibility of conducting clinical trials in infants and children with acute respiratory failure. *Am J Respir Crit Care Med* 2003; 167(10):1334-40.
67. Yapicioğlu H, Yildizdaş D, Bayram I, Sertdemir Y, Yilmaz HL. The use of surfactant in children with acute respiratory distress syndrome: efficacy in terms of oxygenation, ventilation and mortality. *Pulmonary Pharmacology & Therapeutics* 2003; 16(6):327-33.
68. Ramirez JB, Cid JL, Alapont VM. Prevalence of mechanical ventilation in pediatric intensive care units in Spain. *Anales de Pediatría* 2005;61(6):533-41.
69. Lefrant JY, Muller L, De LA Coussaye JE, Prudhomme M, Ripart J, Gouzes C, et al. Risk factors of failure and immediate complications of subclavian vein catheterization in critically ill patients. *Intensive Care Med* 2002; 28(8):1036-41.
70. Boussarsar M, Thierry G, Jaber S, Roudot-Thoraval F, Lemaire F, Brochard L. Relationship between ventilatory settings and barotrauma in acute respiratory distress syndrome. *Intensive Care Med* 2002; 28(4): 406-13.
71. Afessa B, Morales I, Cury JD. Clinical course and outcome of patients admitted to an ICU for status asthmaticus. *Chest* 2001; 120(5): 1616-21.
72. Kotanoglu MS, Karasu S, Tokat AO, Ozcan N, Ozcan A, Karakaya J, et al. Evaluation of iatrogenic pneumothorax cases in the intensive care unit: A retrospective investigation. *Sci. Res. Essays* 2011; 6(29):6243-45.
73. Zwillich CW, Pierson DJ, Creagh CE, Sutton FD, Schatz E, Petty LT, et al. Complications of assisted ventilation. A prospective study of 345 consecutive episodes. *Am J Med* 1974; 57(2):161-70.
74. Zimmerman JE, Dunbar BS, Klingenstein CH. Management of subcutaneous emphysema, pneumomediastinum and pneumothorax during respirator therapy. *Crit Care Med* 1975; 3(2):69-73.

References

75. ATLS. Advanced trauma life support. 6th ed. Chicago: American Collage of Surgeons, 1977.
76. Esteban A, Anzueto A, Frutos F, Alia I, Brochard L, Stewart TE, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation; a 28 day international study. *JAMA* 2002; 287(3):345-55.
77. Chien-Wei H, Shu-Fen S. Iatrogenic pneumothorax related to mechanical ventilation. *World J of Crit Care Med* 2014; 3(1):8-14.
78. Zhan C, Smith M, Stryer D. Accidental iatrogenic pneumothorax in hospitalized patients. *Med Care* 2006; 44(2):182-6.
79. Sassoon CS, Light RW, O'Hara VS, Moritz TE. Iatrogenic pneumothorax: etiology and mortality. Results of a Department of Veterans Affairs Cooperative Study. *Respiration* 1992; 59(4):215-20.
80. Despars JA, Sassoon CS, Light RW. Significance of iatrogenic pneumothoraces. *Chest* 1994; 105(4):1147-50.
81. Johnson NN, Toledo A, Endom EE. Pneumothorax, pneumomediastinum and pulmonary embolism. *Pediatr Clin North Am* 2010; 57(6):1357-83.
82. Kumar A, Pontoppidan H, Falke KJ, Wilson RS, Laver MB. Pulmonary barotrauma during mechanical ventilation. *Crit Care Med* 1973; 1(4): 181-6.
83. Gammon RB, Shin MS, Groves RH, Hardin M, Hsu C, Buchalter SE. Clinical risk factors for pulmonary barotrauma: a multivariate analysis. *Am J Respir Crit Care Med* 1995; 152(4):1235-40.
84. Lew TWK, Kwek T-K, Tai D, Earnest A, Loo S, Singh K, et al. Acute respiratory distress syndrome in critically ill patients with severe acute respiratory syndrome. *JAMA* 2003; 290(3):374-80.
85. Concepcion P, Carlos C, Teresa MM, Luis M, Rosa G, Isabel B, et al. Factors associated to the presence of pneumothorax in cystic fibrosis patients in the city of Madrid. *International J Clin Med* 2011; 2:212-17.
86. Cullen DJ, Caldera DL. The incidence of ventilator-induced pulmonary barotrauma in critically ill patients. *Anaesthesiology* 1979; 50(3):185-90.
87. Rotta AT, Steinhorn DM. Conventional mechanical ventilation in paediatrics. *J Pediatr* 2007; 83(suppl 2):s100-8.
88. Miller Mp, Sagy M. Pressure characteristics of mechanical ventilation and incidence of pneumothorax before and after the implementation of protective lung strategies in the management of pediatric patients with severe ARDS. *Chest* 2008; 134(5):696-73.
89. Anon (2000). Ventilation with low tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 2000;342:1301-8

References

90. Kendirli T, Kavaz A, Yalaki Z, Hişmi ÖB, Derelli E, Ince E. Mechanical ventilation in children. *The Turkish Journal of Pediatrics* 2006; 48:323-2
91. Pierson DJ. Complications associated with mechanical ventilation. *Crit Care Clin* 1990; 6(3):711-24.
92. Downs JB, Chapman RL. Treatment of bronchopleural fistula during continuous positive pressure ventilation. *Chest* 1976; 69(3):363-6.
93. Kirby RR, Downs JB, Civetta JM, Modell JH, Dannemiller FJ, Klein EF, et al. High level positive end expiratory pressure (PEEP) in acute respiratory insufficiency. *Chest* 1975; 67(2):156-63.
94. Steier M, Ching N, Roberts EB, Nealon TF. Pneumothorax complicating continuous ventilatory support. *Journal of Thoracic and Cardiovascular Surgery* 1974; 67(1):17-23.
95. Woodring JH. Pulmonary interstitial emphysema in the adult respiratory distress syndrome. *Crit Care Med* 1985; 13(10):786-91.
96. Haake R, Schlichtig R, Ulstad DR, Henschen RR . Barotrauma. pathophysiology, risk factors, and prevention. *Chest* 1987; 91(4): 608-13.
97. Weg JG, Anzueto A, Balk RA, Wiedemann HP, Pattishall EN, Schork MA, et al. The relation of pneumothorax and other air leaks to mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338(6):341-6.
98. Brochard L, Roudot-Thoraval F, Roupie E, Delclaux C, Chastre J, Fernandez-Mondejar E, et al. Tidal volume reduction for prevention of ventilator-induced lung injury in acute respiratory distress syndrome. The Multicenter Trial Group On Tidal Volume reduction in ARDS. *Am J Respir Crit Care Med* 1998; 158(6):1831-8.
99. Gattinoni L, Pesenti A. The concept of “baby lung”. *Intensive Care Med* 2005; 31:776-84.
100. Gattinoni L, Mascheroni D, Torresin A, Marcolin R, Fumagalli R, Vesconi S, et al. Morphological response to positive end expiratory pressure in acute respiratory failure. Computerized tomography study. *Intensive Care Med* 1986; 12(3):137-42.
101. Gattinoni L, Pesenti A, Bombino M, Baglioni S, Rivolta M, Rossi F, et al. Relationships between lung computed tomographic density, gas exchange, and PEEP in acute respiratory failure. *Anaesthesiology* 1988; 69(6):824-32.
102. Gattinoni L, Bombino M, Pelosi P, Lissoni A, Pesenti A, Fumagalli R, et al. Lung structure and function in different stages of severe adult respiratory distress syndrome. *JAMA* 1994; 271(22):1772-9.
103. Stewart TE, Meade MO, Cook DJ, Granton JT, Hodder RV, Lapinsky SE, et al. Evaluation of a ventilation strategy to prevent barotrauma in patients at high risk for acute respiratory distress syndrome. *N Engl J Med* 1998; 338(6):355-61.

References

104. Farias JA, Fernandez A, Monteverde E, Flores JC, Baltodano A, Menchacha A, et al. Mechanical ventilation in pediatric intensive care units during the season for acute lower respiratory infection: a multicentre study. *Pediatr Crit Care Med* 2012; 13(2):158-64.
105. Soares LC, Ribas D, Spring R, Silva JM, Miyague NI. Clinical profile of systemic inflammatory response after pediatric cardiac surgery with cardio-pulmonary bypass. *Arq BrasCardiol* 2010;94:127-33.
106. Wilkerson RG, Stone MB. Sensitivity of bedside ultrasonography and supine anteroposterior radiographs for identification of pneumothorax after a blunt trauma. *Acad Emerg Med* 2010; 17(1):11-7.
107. Rowan KR, Kirkpatrick AW, Liu D, Forkheim KE, Mayo JR, Nicolaou S. Traumatic pneumothorax detection with thoracic US: correlation with chest radiography and CT –initial experience. *Radiology* 2002; 225(1):210-4.

الدكتور محمد فوزي
مستشفى
صحة المستشفيات
د. اسامعيل عبدالمنعم
٢٠١٤/١٢/١٤
سواحي
٢٠١٤/١٢/١٤

STUDY OF PNEUMOTHORAX IN ALEXANDRIA
UNIVERSITY PEDIATRIC INTENSIVE CARE UNIT:
A 5 YEARS RETROSPECTIVE STUDY

دراسة لحالات الاسترواح الصدري للأطفال في وحدة العناية المركزة بمستشفى الأطفال
الجامعي بالأسكندرية : دراسة استيعابية على مدار خمس سنوات

Protocol of a thesis submitted
to the Faculty of Medicine
University of Alexandria
In partial fulfillment of the
requirements of the degree of
Master of Pediatrics

خطة بحث مقدمة
لكلية الطب
جامعة الإسكندرية
إيفاء جزئياً
لشروط الحصول على درجة
الماجستير في طب الأطفال

by

من

Reem Mohamad Gamal Newegy
MBBCh, Alex.
Resident
Alexandria University Children's Hospital
Department of Pediatrics
Faculty of Medicine
University of Alexandria
2012

ريم محمد جمال نويجي
بكالوريوس الطب والجراحة ، الإسكندرية
طبيب مقوم
مستشفى الأطفال الجامعي بالإسكندرية
قسم طب الأطفال
كلية الطب
جامعة الإسكندرية
٢٠١٢

سواحي
٢٠١٤/١٢/١٤

٢٠١٢

SUPERVISORS

المشرفون

Prof. Dr. Ahmed Ahmed El-Nawawy
Professor of Pediatrics,
Faculty of Medicine,
University of Alexandria.

أ/د/ أحمد أحمد النواوي
استاذ طب الاطفال
كلية الطب
جامعة الإسكندرية



Prof. Dr. Amina Sedky Al-Halawany
Professor of Pediatrics,
Faculty of Medicine,
University of Alexandria.

أ/د/ أمينة صدقي الحلواني
استاذ طب الاطفال
كلية الطب
جامعة الإسكندرية



CO-SUPERVISOR

المشرف المشارك

Dr. Manal Abdelmalik Antonios
Lecturer in Pediatrics,
Faculty of Medicine,
University of Alexandria.
For her experience in pediatric
pneumothorax

د/ منال عبد الملك الطونسيوس
مدرس طب الاطفال
كلية الطب
جامعة الإسكندرية
وذلك لخبرتها في مجال الاسترواح الصدري





ASSISTANT-RESEARCHER

3.

الباحث المساعد

Mohamed Abdelhamid Manaa

5th grade student

Faculty of Medicine,

University of Alexandria.

Mobile phone: 01224468643


E-mail: manaa22@yahoo.com


محمد عبد الحميد مناع

طالب بالفرقة الخامسة

كلية الطب

جامعة الإسكندرية





INTRODUCTION

Pneumothorax refers to the presence of air or gas in the pleural cavity between the visceral and parietal pleura. This condition is uncommon during childhood but can be life threatening.⁽¹⁾

Pneumothorax can be classified into: i) Primary spontaneous pneumothorax which occurs in children without known lung disease. ii) Secondary spontaneous pneumothorax occurs as a complication of chronic or acute lung disease such as asthma, cystic fibrosis and pneumonia. iii) Traumatic pneumothorax which is caused by blunt or penetrating trauma to the chest. iv) Iatrogenic pneumothorax which occurs as a complication of diagnostic or therapeutic procedures such as central line placement or as a consequence of mechanical ventilation.⁽¹⁾

In children, pneumothorax is reported as a complication in 1-4 % of central venous catheter insertion and 0.9-13% of mechanically ventilated cases in pediatric intensive care unit (PICU). The mortality rate of mechanically ventilated patients is 34.5-58.3 % versus the mortality rate of all patients in PICU of 12.2-22.6 %.⁽²⁻⁶⁾

An earlier study in the pediatric population showed that the prevalence of pneumothorax in ventilated patients was significantly higher in the era before protective lung strategies with low tidal volumes were the standard of care.⁽⁷⁾ Higher prevalences of pneumothoraces were seen in patients with acute respiratory distress syndrome (ARDS), but not in those treated with prone positioning or different ventilator strategies related to airway pressures alone.⁽⁸⁻¹⁰⁾



There are three mechanisms by which air can enter the pleural space: 1) A communication between the pleura and alveolar space via visceral pleural rupture. 2) A communication between the pleural space and the atmosphere most commonly due to penetrating chest trauma ⁽¹¹⁾ and 3) The presence of gas producing organisms within the pleural space. ⁽¹²⁾

The severity of the symptoms depends on the extent of lung collapse, rate of development and underlying clinical status of the patient. Pneumothorax may be either asymptomatic or presents with signs of respiratory distress including tachypnea, nasal flaring, accessory muscle use, anxiety or altered mental status. Breath sounds may be decreased or absent on the affected side. The percussion note may be tympanic. Pulsus paradoxus (>10 mmHg fall in systolic blood pressure on inspiration) may be noted. Signs of tension pneumothorax may include deviation of trachea away from affected side and if severe, may include cyanosis, jugular vein distension and deterioration of vital signs leading to pulseless electrical activity and death. ⁽¹³⁾

Pneumothorax should be considered a medical emergency and requires a high index of suspicion, prompt recognition and intervention. The diagnosis of pneumothorax in critically ill patients can be made by physical examination findings or radiographic studies including chest radiographs, ultrasonography or computerized tomography. ⁽¹¹⁾

The cause of pneumothorax, as well as the patient's underlying disease greatly influence the treatment course and overall prognosis in critically ill patients. ⁽¹⁴⁾ The primary goal during the management of pneumothorax is to evacuate air from pleural space and allow apposition of the lung and chest wall. Although many patients with pneumothorax can be managed conservatively, the majority of patients in the ICU require pleural intervention. ⁽¹⁵⁾



AIM OF THE WORK

The aim of this work is to conduct a five year (1st of January 2007-31st December 2011) retrospective study of the cases of pneumothorax among pediatric intensive care unit admissions in Alexandria University Children's Hospital.

MATERIAL

This study will be based on the data retrieved from the files of all children admitted at pediatric intensive care unit of Alexandria University Children's Hospital during five years (1st of January 2007-31st of December 2011) who were diagnosed as having pneumothorax either on admission or during hospital stay.


منال عبدالقادر
عبدالمعز
[Signature]

METHODS

Files of patients with pneumothorax at pediatric intensive care unit of Alexandria University Children's Hospital during the period between (1st January 2007-31st December 2011) will be reviewed regarding:

- History, physical examination, laboratory investigations especially cultures and radiological studies.
- Pediatric index of mortality 2 Score (PIM 2 Score) and Pediatric logistic organ dysfunction score (PELOD Score).
- Admission diagnosis.
- Ventilatory details (if any) including: timing, setting, setting changes and weaning.
- Pneumothorax details: as regards etiology, diagnosis, radiology, management and complications.
- Fate and complications including 28 day mortality.

The study will be done after the approval of Alexandria University Ethical Committee. Informed consents of the parents will be waived as the patients were discharged during the last five years and there will be no application of prospective procedures to the patients. All measures will be taken so as not to spread private data as names of these patients.

The block contains three handwritten signatures or initials in black ink. The first is a large, stylized signature on the left. The second is a smaller signature in the middle. The third is a large, circular signature on the right.

ETHICS OF RESEARCH

Research on human or human products:

- Prospective study: informed consent will be taken from patients. In case of incompetent patients the informed consent will be taken from the guardians.
- Retrospective study: confidentiality of records will be considered.
- DNA/genomic material: informed consent for DNA / genomic test and for research will be taken from patients. No further test will be carried out except with further approval of committee and patients. If the samples will travel outside Egypt the researcher will be responsible for transportation and security approval.
- All drugs used in the research are approved by the Egyptian Ministry of Health.


Research on animal:

- The animal species are appropriate for the test.
- After test, if animal will suffer, it will be euthanized and properly disposed.
- After operation, it will have a proper postoperative care.



RESULTS

The results of this study will be tabulated and analyzed using the appropriate statistical methods and SSPS version 20.0.0.

سازگار الله
مهاجر


DISCUSSION

The results will be discussed in view of achievement of the aim, and will be compared to those of similar published studies.

سید محمد علی
سید
P

REFERENCES

1. William G, Rene JF, Thomas S. Pediatric pneumothorax. Medscape 2011; 29-3.
2. Kaplan J, Brill RJ. Vascular access. In: Wheeler DS, Wong HR, Shanley TP, editors. Pediatric critical medicine: Basic science and clinical evidence. London: Springer 2007; 253-71.
3. Khilnani P, Sarma R, Singh R, Uttam R, Rajdev S, Makkar A, et al. Demographic profile and outcome analysis of tertiary level pediatric intensive care unit. Indian J Pediatr 2004; 71:587-91.
4. Kendirli T, Kavaz A, Yalaki Z, Ozturk HB, Derelli E, Ince E. Mechanical ventilation in children .The Turkish Journal Of Pediatrics 2006; 48:323-7.
5. Medjo B, Vunjak N, Rsovac S, Nikolic D, Kalanj J, Cuturilo G, et al. Indications and complications of mechanical ventilation in pediatric intensive care unit patients. Archives of Disease in Childhood 2008; 93:491.
6. Ozdemir H, Kantar A, Dinlen N, Coskun E, Ozyoruk D, Metin A. The first results of mechanical ventilation in a newly opened pediatric intensive care unit. Turk Pediatri Arsivi 2008; 43:99-101.
7. Miller MP, Sagy M. Pressure characteristics of mechanical ventilation and incidence of pneumothorax before and after the implementation of protective lung strategies in the management of pediatric severe ARDS Chest 2008; 1345:969-73.



8. Boussarsar M, Thierry G, Jaber S, Roudot-Thoraval F, Lemaire F, Brochard L. Relationship between ventilator settings and barotrauma in the acute respiratory distress syndrome. *Intensive Care Med* 2002; 284:406-13.
9. Kopterides P, Siempos II, Armaganidis A. Prone positioning in hypoxemic respiratory failure: meta-analysis of randomized controlled trials. *Crit Care* 2009; 241:89-100.
10. Briel M, Meade M, Mercat A, Brower RG, Talmor D, Walter SD, et al. Higher VS lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome : systematic review and meta-analysis. *JAMA* 2010; 3039:865-73.
11. Lonny Y, David FK. Pneumothorax in critically ill patient. *Chest* 2012; 141:1098-105.
12. Noppen M, De Keukeleire T. Pneumothorax. *Respiration* 2008; 762:121-7.
13. Ciorciari AJ. Trauma. In: Ellen FC, Jeffrey CG, editors. *Clinical manual of emergency pediatrics .5 th edition*. New York: Cambridge University Press, United States of America 2010; 702-23.
14. Rahman NM, Davies RVO, Gleeson FV. Pleural interventions: management of acute and chronic pneumothorax. *Semin Respir Crit Care Med* 2008; 294:427-40.
15. Sahn SA, Heflinger JE. Spontaneous pneumothorax. *N Engl J Med* 2000; 34212:868-74.



المخلص العربي

الاسترواح الصدرى في المرضى ذوي الحالات الحرجة لا يزال يمثل مشكلة شائعة في وحدات العناية المركزة، وهو ما يمثل حدث يحتمل أن يهدد الحياة، وخاصة إذا كان مرتبط بالتهوية الميكانيكية.

الاسترواح الصدرى يمكن أن يكون عفويا أو علاجي المنشأ، ويعتبر النوع الاخير هو الاكثر شيوعا في جميع أنحاء العالم.

وينبغي اعتبار حالات الاسترواح الصدرى من الحالات الطبية الطارئة والتي تتطلب مستوى عالى من الشك، والتعرف الفوري والتدخل السريع، ويمكن تشخيص حالات الاسترواح الصدرى في المرضى ذوي الحالات الحرجة من خلال نتائج الفحص البدنى او الدراسات الاشعاعية بما في ذلك الاشعة السينية للصدر والاشعة المقطعية.

والهدف من هذا العمل هو تقييم حالات الاسترواح الصدرى في اطفال العناية المركزة بمستشفى جامعة الاسكندرية خلال خمس سنوات.

وقد اجريت هذه الدراسة على جميع حالات الدخول في وحدة العناية المركزة للاطفال في الفترة بين ١ يناير ٢٠٠٨ الى ٣١ ديسمبر ٢٠١٢ وعددهم ١٢٩٨ حالة، وقد تم استعراض كافة الملفات للحالات و التي اظهرت ان ١٣٥ حالة عانت من الاسترواح الصدرى، وتم مقارنة هذه الحالات مع باقي الحالات التي لم تعاني من الاسترواح الصدرى وذلك من اجل دراسته معدل الحنوث والمخاطر والمضاعفات المرتبطة لاسترواح الصدرى.

و جرى استعراض ملفات جميع المرضى من حيث التاريخ المرضى، والفحص البدنى، والتشخيص، واحتمال الوفيات عند الدخول وتم حسابه باستخدام مؤشر وفيات الاطفال PIM2 score، ودرجه ضعف الاعضاء عند الدخول وعلى اساس يومية وقد تم حسابه باستخدام PELOD score، والحاجة الى التهوية الميكانيكية، وضبط جهاز التنفس الصناعى ومصير الحالات (تحسن او متوفى). كما تم استعراض ملفات حالات الاسترواح الصدرى لتوضيح مسببات المرض، شدته، مضاعفاته، الوسيله الاولى المستخدمه في التشخيص المبندى و طرق العلاج.

ولقد كشفت النتائج فروقات ذات دلالات احصائية من حيث ارتفاع معدل الوفيات، وطول مده الاقامه وصغر سن المريض، وانخفاض وزن الجسم وارتفاع احتمال الوفيات ودرجه ضعف الاعضاء عند الدخول في حالات الاسترواح الصدرى مقارنة للحالات التي لا تعاني من الاسترواح الصدرى.

كما اظهرت الدراسة ان الاسترواح الصدرى علاجي المنشأ اكثر شيوعا من الاسترواح الصدرى العفوى. وكان السبب الاكثر شيوعا للاسترواح الصدرى في هذه الدراسة هو الاذى الضغطى المتعلق بالتهوية الميكانيكية يليها ادخال القسطرة الوريدية المركزية، وكانت الوسيله الاكثر شيوعا لتشخيص الاسترواح الصدرى هي الاشعة السينية للصدر يليها الشك الاكلينيكي مع عدم وجود دور للموجات فوق الصوتية لتشخيص في هذه الدراسة.

كما اظهرت النتائج وجود فروقات ذات دلالة احصائية في حالات التهوية الميكانيكية حيث كانت هناك مدة اطول للتهوية الميكانيكية، وضبطيه تنفس صناعى عالية في كلا التنفس الصناعى التقليدى و التهوية التذبيبه عاليه التردد في حالات الاسترواح الصدرى مقارنة بالحالات التي لا تعاني من الاسترواح الصدرى.

وكذلك اظهرت النتائج ارتفاع ملحوظ بنسبة ثانى اكسيد الكربون وانخفاض فى نسبة الاوكسجين فى حالات الاسترواح الصدرى مقارنة بباقي الحالات، بينما لم تظهر نتائج معامل الالتهاب، وعدد كرات الدم البيضاء ودرجه تشبع الدم بالاوكسجين فروقات ملحوظه.

بمقارنه نتائج تحاليل غازات الدم فى حالات الاسترواح الصدرى فى وقت الدخول وعند حدوث الاسترواح الصدرى اظهرت النتائج ارتفاع ملحوظ فى نسبة حموضه الدم و نسبة ثانى اكسيد الكربون و انخفاض فى نسبة الاوكسجين و درجه تشبع الاوكسجين بالدم عند حدوث الاسترواح الصدرى.

وحيث ان السبب الاكثر شيوعا للاسترواح الصدرى فى دراستنا هو الاذى الضغطى المتعلق بالتهوية الميكانيكية لذلك من المهم جدا تطبيق الاستراتيجيات الوقائية للرنه بدقه متناهيه من اجل تقليل خطر حدوث الاسترواح الصدرى.

دراسة لحالات الاسترواح الصدرى للأطفال فى وحدة العناية المركزة
بمستشفى الأطفال الجامعى بالأسكندرية: دراسة استعادىة على مدار خمس
سنوات

رسالة مقدمة

لقسم طب الأطفال - كلية الطب - جامعة الإسكندرية
ضمن متطلبات درجة

الماجستير

فى

طب الأطفال

من

ريم محمد جمال نوىجى

بكالوريوس الطب والجراحة، ٢٠٠٦

كلية الطب، جامعة الإسكندرية

[٢٠١٥]

دراسة لحالات الاسترواح الصدري للأطفال في وحدة العناية المركزة
بمستشفى الأطفال الجامعي بالإسكندرية: دراسة استيعادية على مدار خمس
سنوات

رسالة مقدمة من

ريم محمد جمال نويجي

للحصول على درجة

الماجستير

في

طب الأطفال

التوقيع

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د/ أحمد أحمد النواوي
أستاذ طب الأطفال
قسم طب الأطفال
كلية الطب
جامعة الإسكندرية

أ.د/ محمد أحمد بدر
أستاذ طب الأطفال
قسم طب الأطفال
كلية الطب
جامعة الزقازيق

أ.د/ تيسير محمد زيتون
أستاذ ورئيس قسم الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الإسكندرية

التاريخ / /

موافقون

لجنة الإشراف

.....

أ.د/ أحمد أحمد النواوى

أستاذ طب الأطفال
قسم طب الأطفال
كلية الطب
جامعة الإسكندرية

.....

أ. د/ أمينة صدقى الحلوانى

أستاذ طب الأطفال
قسم طب الأطفال
كلية الطب
جامعة الإسكندرية

المشرف المشارك

.....

د/ منال عبد الملك انطونيوس

مدرس طب الأطفال
قسم طب الأطفال
كلية الطب
جامعة الإسكندرية