

ACKNOWLEDGEMENT

Thank to Allah

For

Accomplishment of this work

I wish to express my deepest gratitude to all those who assisted me to complete this work.

First and foremost, my thanks are directed to ***Prof. Dr. Nair Abou Rawash, for his Soul*** Professor of Biochemistry, Medical Research Institute, University of Alexandria, for his unlimited help.

Many thanks and appreciation to ***Prof. Dr. Madiha Hassan Helmy***, Professor of Biochemistry, Medical Research Institute, University of Alexandria, for her supervision and encouragement and for her kindness throughout the work and her continuous insistence on perfection, without her constant supervision, this thesis could not have achieved in its present form.

I am greatly indebted to ***Prof. Dr. Magda Ismail Youssif***, Professor of Histochemistry, Medical Research Institute, University of Alexandria, for fruitful suggestions and wise guidance created this thesis.

I am greatly indebted to ***Prof. Dr. Maher Abd el Naby Kamel Rashwan***, Professor of Biochemistry, Medical Research Institute, University of Alexandria, for his constant encouragement and continuous guidance as well as his perseverance

It is my pleasure to extend my thanks to all members of the staff of the Biochemistry Department.

LIST OF CONTENTS

Chapter	Page
ACKNOWLEDGMENTS.....	i
LIST OF CONTENTS.....	ii
LIST OF TABLES.....	iii
LIST OF FIGURES.....	v
LIST OF ABBREVIATIONS	ix
I. INTRODUCTION	1
II. AIM OF THE WORK.....	26
III. MATERIALS AND METHODS.....	27
IV. RESULTS.....	43
V. DISCUSSION.....	84
VI. SUMMARY AND CONCLUSIONS.....	90
VII. RECOMMENDATIONS	92
VIII. REFERENCES	93
APPENDIX	
PROTOCOL	
ARABIC SUMMARY	

LIST OF TABLES

Table	Page
(1) Criteria for the diagnosis of glucose intolerance and diabetes -----	14
(2) The serum concentration of fasting glucose (mg/dl) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	44
(3) The serum concentration of insulin ($\mu\text{g/ml}$) in the control rats and rats supplemented with sodium selenate and selenocysteine at different doses -----	45
(4) The HOMA-insulin resistance index calculations in the control rats and rats supplemented with sodium selenate and selenocysteine at different doses -----	46
(5) The hepatic content of phospho-insulin receptor (ng/mg protein) in the control rats and rats supplemented with sodium selenate and selenocysteine at different doses	48
(6) The muscle content of phospho-insulin receptor (ng/mg protein) in the control rats and rats supplemented with sodium selenate and selenocysteine at different doses	49
(7) White adipose tissue content of phospho insulin receptor (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses ----	50
(8) The muscle content of glucose transporter 4 (ng /mg protein) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	51
(9) White adipose tissue content of glucose transporter 4 (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	52
(10) The serum cholesterol (mg/dl) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	54
(11) Serum concentration of triglycerides (mg/dl) in the control rats and rats treated with sodium selenat an selenocysteine iat different doses -----	55
(12) Serum concentration of high density lipoprotein (mg/dl) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	56
(13) The serum concentration of low density lipoprotein (mg/dl) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	57
(14) Blood concentration of urea (mg/dl) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	59
(15) Serum concentration of creatinine (mg/dl) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	60
(16) Hepatic content of total glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	62

(17)	The muscle content of total glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	63
(18)	The white adipose tissue content of total glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenate and selenocysteine at different doses -----	64
(19)	The hepatic content of selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	65
(20)	The muscle content of selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	66
(21)	The white adipose tissue content of selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses-----	67
(22)	The hepatic content of non selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	68
(23)	Muscle content of non selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	69
(24)	The white adipose tissue content of non selenium dependent glutathione peroxidase (ng/mg protein) in control rats and rats treated with sodium selenate and selenocysteine at different doses-----	70
(25)	The hepatic content of selenoprotein P (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	71
(26)	The muscle content of selenoprotein P (ng/mg protein) in the control rats and rats treated with sodium selenat and selenocysteine at different doses -----	72
(27)	White adipose tissue content of selenoprotein P (ng/mg protein) in control rats and rats treated with sodium selenat and selenocysteine at different doses -----	73
(28)	Correlation between all parameters -----	74

LIST OF FIGURES

Figure	Page
(1) Insulin biosynthesis and processing -----	2
(2) Intracellular mechanisms through which glucose stimulates insulin secretion Intracellular mechanisms through which glucose stimulates insulin secretion ---	3
(3) A schematic illustration of general insulin signaling pathways involved in both metabolic and mitogenic effects -----	5
(4) Summary of insulin signaling pathways in the liver -----	7
(5) An overview of insulin signaling pathways in skeletal muscle.-----	9
(6) Summary of insulin function involved in lipogenesis and lipolysis in adipose tissue. -	11
(7) Human selenoprotein -----	18
(8) Scheme depicting a potential influence of selenium on components of the insulin signaling cascade.-----	24
(9) Standard curve of total protein. -----	29
(10) Standards Curve of Glut-4-----	32
(11) Standards Curve of Se-P-----	34
(12) Standard curve of insulin. -----	40
(13) The change in fasting glucose (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	44
(14) The serum concentration of insulin ($\mu\text{g/ml}$) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	45
(15) The change in HOMA-insulin resistance index in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	46
(16) The change in hepatic content of phospho-insulin receptor (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	48
(17) The change in muscle content of phospho insulin receptor (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	49
(18) The change in white adipose tissue content of phospho-insulin receptor (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	50

(19)	The change in muscle content of glucose transporter 4 (ng /mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	51
(20)	The change in white adipose tissue content of glucose transporter 4 (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	52
(21)	The change in serum cholesterol (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	54
(22)	The change in serum concentration of triglycerides (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	55
(23)	The change in serum concentration of HDL-Cholesterol (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	56
(24)	The change in serum concentration of LDL-Cholesterol (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	57
(25)	The change in serum urea (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	59
(26)	The change in serum concentration of creatinine (mg/dl) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	60
(27)	The change in hepatic activity of total glutathione peroxidase in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	62
(28)	The change in muscle activity in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	63
(29)	The change in white adipose tissue activity of total glutathione peroxidase in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	64
(30)	The change in hepatic activity of selenium dependent glutathione peroxidase (Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	65
(31)	The change in muscle activity of selenium dependent glutathione peroxidase (Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	66
(32)	The change in white adipose tissue activity of selenium dependent glutathione peroxidase (Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	67

(33)	The hepatic activity of non selenium dependent glutathione peroxidase (n-Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	68
(34)	The change in muscle activity of non selenium dependent glutathione peroxidase (n-Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	69
(35)	The change in white adipose tissue activity of non selenium dependent glutathione peroxidase (n-Se-GPx) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months -----	70
(36)	The change in hepatic content of selenoprotein P (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	71
(37)	The change in muscle content of selenoprotein P (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	72
(38)	The change in white adipose tissue content of selenoprotein P (ng/mg protein) in the rats supplemented with sodium selenate and selenocysteine at different doses for 3 months-----	73
(39)	Correlation curve between serum fasting glucose and selenoprotein p in liver in selenocysteine group. -----	74
(40)	Correlation curve between serum fasting glucose and selenoprotein p in muscle in selenocysteine group -----	75
(41)	Correlation curve between fasting serum glucose and selenoprotein p in adipose tissue in selenocysteine group.-----	75
(42)	Correlation curve between fasting serum glucose and activity of se-dependent GPX in liver in selenocysteine group-----	76
(43)	Correlation curve between fasting serum glucose and activity of Se-dependent GPX in muscle in both groups (sodium selenat and selenocysteine) -----	76
(44)	Correlation curve between fasting glucose and activity of Se-dependent GPX in adipose tissue in Sodium selenate group -----	77
(45)	Correlation curve between activity of Se-dependent glutathione peroxidase in liver and serum insulin in sodium selenat group -----	77
(46)	Correlation curve between Selenoprotein P in liver and glucose transporter 4 in muscle in both groups-----	78
(47)	Correlation curve between selenoprotein P and GLUT 4 in muscle in both groups---	78
(48)	Correlation curve between phspho insulin receptor and GLUT4 in muscle in selenocysteine treated group -----	79

(49)	Correlation between serum triglycerides and selenoprotein P in liver in sodium selenat group-----	79
(50)	Correlation between serum triglycerides and the activity of Se-dependent GPX in the liver in sodium selenat group -----	80
(51)	Correlation between serum triglycerides and the activity of Se-dependent GPX in the muscle in sodium selenat group -----	80
(52)	Correlation between the activity of Se-dependent GPX and selenoprotein P in liver in selenocysteine group -----	81
(53)	H&E stain of pancreas section from control rats -----	82
(54)	H&E stain of pancreas section from rats supplemented with 100µg/kg sodium selenate -----	82
(55)	H&E stain of pancreas section from rats supplemented with 100µg/kg selenocysteine -----	83