
SUMMARY

Care should be taken for cirrhotic patients undergoing liver resection under general anaesthesia concerning perioperative hepatocellular integrity and kidney functions.^(1,2)

Our aim is to compare Target controlled infusion (TCI) of propofol-fentanyl versus Desflurane (Des) based anesthesia in major liver resection for cirrhotic patients.

In a double blinded hospital based prospective randomized controlled study, 50 adult cirrhotic patients (Child A) were categorized into 2 groups (TCI group $n=25$ and Desflurane (D) group $n=25$) for anaesthesia maintenance. In TCI before induction all subjects received Fentanyl $3 \mu\text{g/kg}$ for 30 seconds and then an infusion of $2 \mu\text{g/kg/h}$ for 30 min, $1.5 \mu\text{g/kg/h}$ from 31–150 min, and $1 \mu\text{g/kg/h}$ until 30 min before skin closure and this is adjusted according to pharmacokinetics of Navigator. Propofol was administered with a syringe pump integrated with (controller and software) GE Healthcare Finland (Datex-Ohmeda, Helsinki, Finland). The Propofol venous blood target concentration (C_t) for induction of anesthesia was set at $4 \mu\text{g/ml}$. During maintenance anesthetist was able to increase or decrease Et Des by 1% increments, or the Propofol (C_t) by $0.5 \mu\text{g/ml}$ to keep Entropy between 40-60 and maintain patients' hemodynamics.

Liver and Kidney functions tests, Urinary Microalbuminurea, Glutathione-S-transferase (GST), were assayed preoperatively (T_1), immediately postoperative (T_2) and 48 hours postoperatively (T_3). Haemodynamic, transfusion requirements, recovery, complications, intensive care unit and hospital stay together with economic cost were recorded.

Significant reduction in systemic vascular resistance (SVR) occurred with TCI versus Des (779 ± 36 vs $836 \pm 8 \text{ dyn.sec.cm}^{-5}$, $P=0.000$), this was associated with a significant reduction in mean blood pressure and stroke volume respectively (85 ± 3 vs $91 \pm 4 \text{ mmHg}$, 76 ± 7 vs $86 \pm 4 \text{ ml}$, $P=0.000$). ALT and AST peaked in both groups at (T_2), (Des) group was 378 ± 80 and $407 \pm 35 \text{ U/L}$ while, for TCI group was 467 ± 38 and $413 \pm 39 \text{ U/L}$ respectively, this increase was significantly less in (Des) group $P=0.000$. At (T_3) both parameters dropped. There was no significant difference between groups regarding, GST, Microalbuminurea and ICU and hospital stay. Significant difference between both groups regarding Fentanyl intake, Des group was 270.00 ± 29.86 and TCI group was $584.82 \pm 126.32 \mu\text{gm}$. Extubation time was prolonged in TCI $15.20 \pm 2.629 \text{ min}$ versus $9.76 \pm 1.507 \text{ min}$ in Des, $P=0.000$. Des group was lower than TCI group in costs, 235 ± 26 versus 438 ± 57 Egyptian pounds or 33.70 ± 3.836 versus $62.65 \pm 8.233 \text{ U S Dollars}$, $P=0.000$. respectively during the same surgical duration and with the same surgical team.

Desflurane appears a more appropriate choice than Target-controlled infusion of Propofol-Fentanyl in cirrhotic patients undergoing major hepatic resection surgery from haemodynamic stability, recovery characteristics and costs, but neither is superior to the other with respect to their effect on both liver and kidney organs.

CONCLUSION

- Desflurane is a more appropriate choice than Target-controlled infusion of Propofol-Fentanyl in cirrhotic patients undergoing major hepatic resection surgery due to its better haemodynamic stability, recovery characteristics and costs, but neither is superior to the other in respect to their effect on both liver and kidney.
- Significant haemodynamic parameters changes were associated with liver resection which can be summarized in significant increase in cardiac output and significant reduction in systemic vascular resistance.
- Transoesophageal Doppler is a reliable minimally invasive monitor, but the liver mobilization during resection of hepatic tumors requires frequent repositioning of the Doppler probe, this repeatedly affect the TED probe position and hence readings.

RECOMMENDATIONS

1. Desflurane is a more appropriate choice than TCI Propofol-Fentanyl in cirrhotic patients undergoing major hepatic resections from haemodynamic prospective.
2. Neither Desflurane nor TCI Propofol-Fentanyl is superior to the other with respect to their effects on both the liver and kidney. Both can be used safely for liver surgery in cirrhotic patients.
3. Close cardiovascular monitoring is essential during major liver resection and is highly recommended.
4. Transoesophageal Doppler (TED) can be used reliably as a monitoring tool for haemodynamic parameters during and after major surgical procedures. TED as a minimally invasive monitor is recommended in hepatic patients particularly with coagulopathy.
5. Laboratory markers of hepatic (blood glutathione-S-transferase) and renal integrity (urinary microalbumin) can be used to monitor perioperative changes during and after major surgery.
6. Maintaining general anaesthesia with Desflurane is cost effective and promotes early recovery in this category of hepatic patients, this supports it's application during liver surgery.
7. Navigation system is a recommended tool when intravenous general anaesthesia is used.

REFERENCE

- (1) Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010 Dec 15;127(12):2893-917.
- (2) Di Bisceglie AM, Simpson LH, Lotze MT, Hoofnagle JH. Development of hepatocellular carcinoma among patients with chronic liver disease due to hepatitis C viral infection. *J Clin Gastroenterol* 1994 Oct;19(3):222-6.
- (3) Hassan MM, Zaghloul AS, El-Serag HB, Soliman O, Patt YZ, Chappell CL, et al. The role of hepatitis C in hepatocellular carcinoma: a case control study among Egyptian patients. *J Clin Gastroenterol* 2001 Aug;33(2):123-6.
- (4) El-Zayadi AR, Badran HM, Barakat EM, Attia M, Shawky S, Mohamed MK, et al. Hepatocellular carcinoma in Egypt: a single center study over a decade. *World J Gastroenterol* 2005 Sep 7;11(33):5193-8.
- (5) El Sharkawy OA, Refaat EK, Ibraheem AE, Mahdy WR, Fayed NA, Mourad WS, et al. Transoesophageal Doppler compared to central venous pressure for perioperative hemodynamic monitoring and fluid guidance in liver resection. *Saudi J Anaesth* 2013 Oct;7(4):378-86.
- (6) Niemann CU, Roberts JP, Ascher NL, Yost CS. Intraoperative hemodynamics and liver function in adult-to-adult living liver donors. *Liver Transpl* 2002 Dec;8(12):1126-32.
- (7) Bedawy AA, Fayed NA, Morad W, Madany S, Nematalah F, Yassen KA. Rotational Thrombo Elastometry and Standard Coagulation Tests for Hepatic Patients undergoing Major Liver Resection. *J Anaesth and Clin Research* 2013;3(9):1-5.
- (8) Ayanoglu HO, Ulukaya S, Yuzer Y, Tokat Y. Anesthetic management and complications in living donor hepatectomy. *Transplant Proc* 2003 Dec;35(8):2970-3.
- (9) Mc Kuskey R. Anatomy of the normal liver. In: Boyer TD, Manns MP, Sanyal AJ, editors. *Zakim and Boyer's Hepatology: A Textbook of Liver Disease*. 6 ed. Philadelphia: Elsevier Saunders; 2012. p. 3-19.
- (10) Mushlin PS, Gelman S. Anesthesia and the Liver . In: Barash PG, Cullen BF, Stoelting RK, editors. *Clinical Anesthesia*. 4 ed. Philadelphia: Lippincott Williams & Wilkins, 2001. p. 1067.
- (11) Madoff DC, Hicks ME, Vauthey JN, Charnsangavej C, Morello FA, Jr., Ahrar K, et al. Transhepatic portal vein embolization: anatomy, indications, and technical considerations. *Radiographics* 2002 Sep;22(5):1063-76.
- (12) Benedikt HJ, Pannen A. Hepatic blood flow during anesthesia and surgery. European Society of Anaesthesiologists refresher courses. 2000; Vienna 2000.
- (13) Berendes E, Lippert G, Loick HM, Brussel T. Effects of enflurane and isoflurane on splanchnic oxygenation in humans. *J Clin Anesth* 1996 Sep;8(6):456-68.

- (14) Frink EJ, Jr., Morgan SE, Coetzee A, Conzen PF, Brown BR, Jr. The effects of sevoflurane, halothane, enflurane, and isoflurane on hepatic blood flow and oxygenation in chronically instrumented greyhound dogs. *Anesthesiology* 1992 Jan;76(1):85-90.
- (15) Gelman S, Frenette L. Effects of anaesthetics on liver blood flow. *Baillière's Clinical Anaesthesiology* 1992 Dec;6(4):729-50.
- (16) Fujita Y. Effects of PEEP on splanchnic hemodynamics and blood volume. *Acta Anaesthesiol Scand* 1993 May;37(4):427-31.
- (17) Pannen BH, Bauer M, Noldge-Schomburg GF, Zhang JX, Robotham JL, Clemens MG, et al. Regulation of hepatic blood flow during resuscitation from hemorrhagic shock: role of NO and endothelins. *Am J Physiol* 1997 Jun;272(6 Pt 2):H2736-H2745.
- (18) Pannen BH, Kohler N, Hole B, Bauer M, Clemens MG, Geiger KK. Protective role of endogenous carbon monoxide in hepatic microcirculatory dysfunction after hemorrhagic shock in rats. *J Clin Invest* 1998 Sep 15;102(6):1220-8.
- (19) Lautt WW, Legare DJ, Ezzat WR. Quantitation of the hepatic arterial buffer response to graded changes in portal blood flow. *Gastroenterology* 1990 Apr;98(4):1024-8.
- (20) Lebrec D, Hillon P, Munoz C, Goldfarb G, Nouel O, Benhamou JP. The effect of propranolol on portal hypertension in patients with cirrhosis: a hemodynamic study. *Hepatology* 1982 Sep;2(5):523-7.
- (21) Hedner U, Ezban M. Recent advances in clot biology and assessment of clotting. *Trauma* 2008 Oct 1;10(4):261-70.
- (22) Laffan MA, Manning RA. Investigation of haemostasis. In: Lewis SA, Bain B, Bates I, editors. *Practical haematology*. 9 ed. London: Churchill Livingstone; 2001. p. 339-91.
- (23) Cesarman-Maus G, Hajjar KA. Molecular mechanisms of fibrinolysis. *Br J Haematol* 2005 May;129(3):307-21.
- (24) Greenberg DL, Davie LW. Blood coagulation factors: their complementary DNA, genes and expression. In: Coluan RW, Hiroh J, Marder VJ, Clower AW, George JN, editors. *Hemostasis and Thrombosis: Basic Principles and Clinical Practice*. Philadelphia: PA: Lippincott Williams and Wilkins; 2001. p. 21-57.
- (25) Gelman S. General anesthesia and hepatic circulation. *Can J Physiol Pharmacol* 1987 Aug;65(8):1762-79.
- (26) Lai HC, Lai HC, Wang KY, Lee WL, Ting CT, Liu TJ. Severe pulmonary hypertension complicates postoperative outcome of non-cardiac surgery. *Br J Anaesth* 2007 Aug;99(2):184-90.
- (27) Mullin EJ, Metcalfe MS, Maddern GJ. How much liver resection is too much? *Am J Surg* 2005 Jul;190(1):87-97.

- (28) Teh SH, Christein J, Donohue J, Que F, Kendrick M, Farnell M, et al. Hepatic resection of hepatocellular carcinoma in patients with cirrhosis: Model of End-Stage Liver Disease (MELD) score predicts perioperative mortality. *J Gastrointest Surg* 2005 Dec;9(9):1207-15.
- (29) Schroeder RA, Marroquin CE, Bute BP, Khuri S, Henderson WG, Kuo PC. Predictive indices of morbidity and mortality after liver resection. *Ann Surg* 2006 Mar;243(3):373-9.
- (30) Cucchetti A, Ercolani G, Vivarelli M, Cescon M, Ravaioli M, La BG, et al. Impact of model for end-stage liver disease (MELD) score on prognosis after hepatectomy for hepatocellular carcinoma on cirrhosis. *Liver Transpl* 2006 Jun;12(6):966-71.
- (31) Cescon M, Cucchetti A, Grazi GL, Ferrero A, Vigano L, Ercolani G, et al. Indication of the extent of hepatectomy for hepatocellular carcinoma on cirrhosis by a simple algorithm based on preoperative variables. *Arch Surg* 2009 Jan;144(1):57-63.
- (32) Van den Broek MA, Olde Damink SW, Dejong CH, Lang H, Malago M, Jalan R, et al. Liver failure after partial hepatic resection: definition, pathophysiology, risk factors and treatment. *Liver Int* 2008 Jul;28(6):767-80.
- (33) Moemen ME, Gaafar TY, Farag WA, El-Gammal SM, Faramawy M, El-Khouly A. Prognostic Categorization in Cirrhotic Patients Undergoing Abdominal Surgery: A Randomized Trial. *Eg J Anaesth* 200;20:7-27.
- (34) Webster NR. Scoring systems and surgery in cirrhotic patients. Editorial. *Eg J Anaesth* 2004;20:1-3.
- (35) Bates SM, Weitz JI. Coagulation assays. *Circulation* 2005 Jul 26;112(4):e53-e60.
- (36) Kottke-Marchant K, Corcoran G. The laboratory diagnosis of platelet disorders. *Arch Pathol Lab Med* 2002 Feb;126(2):133-46.
- (37) Hayward CP, Harrison P, Cattaneo M, Ortel TL, Rao AK. Platelet function analyzer (PFA)-100 closure time in the evaluation of platelet disorders and platelet function. *J Thromb Haemost* 2006 Feb;4(2):312-9.
- (38) Johansson PI. The blood bank: from provider to partner in treatment of massively bleeding patients. *Transfusion* 2007 Aug;47(2 Suppl):176S-81S.
- (39) Bennett JJ, Blumgart LH. Assessment of hepatic reserve prior to hepatic resection. *J Hepatobiliary Pancreat Surg* 2005;12(1):10-5.
- (40) Buell JF, Rosen S, Yoshida A, Labow D, Limsrichamrern S, Cronin DC, et al. Hepatic resection: effective treatment for primary and secondary tumors. *Surgery* 2000 Oct;128(4):686-93.
- (41) Chiappa A, Zbar AP, Audisio RA, Leone BE, Biella F, Staudacher C. Factors affecting survival and long-term outcome in the cirrhotic patient undergoing hepatic resection for hepatocellular carcinoma. *Eur J Surg Oncol* 2000 Jun;26(4):387-92.
- (42) Poon RT, Fan ST, Lo CM, Liu CL, Wong J. Intrahepatic recurrence after curative resection of hepatocellular carcinoma: long-term results of treatment and prognostic factors. *Ann Surg* 1999 Feb;229(2):216-22.

- (43) Patel T. Surgery in the patient with liver disease. *Mayo Clin Proc* 1999 Jun;74(6):593-9.
- (44) Lai EC, Chu KM, Lo CY, Mok FP, Fan ST, Lo CM, et al. Surgery for malignant obstructive jaundice: analysis of mortality. *Surgery* 1992 Nov;112(5):891-6.
- (45) Behrns KE, Tsiotos GG, DeSouza NF, Krishna MK, Ludwig J, Nagorney DM. Hepatic steatosis as a potential risk factor for major hepatic resection. *J Gastrointest Surg* 1998 May;2(3):292-8.
- (46) Mansour A, Watson W, Shayani V, Pickleman J. Abdominal operations in patients with cirrhosis: still a major surgical challenge. *Surgery* 1997 Oct;122(4):730-5.
- (47) Olthoff KM, Brown RS, Jr., Delmonico FL, Freeman RB, McDiarmid SV, Merion RM, et al. Summary report of a national conference: Evolving concepts in liver allocation in the MELD and PELD era. December 8, 2003, Washington, DC, USA. *Liver Transpl* 2004 Oct;10(10 Suppl 2):A6-22.
- (48) Dunn W, Jamil LH, Brown LS, Wiesner RH, Kim WR, Menon KV, et al. MELD accurately predicts mortality in patients with alcoholic hepatitis. *Hepatology* 2005 Feb;41(2):353-8.
- (49) Kooby DA, Stockman J, Ben-Porat L, Gonen M, Jarnagin WR, Dematteo RP, et al. Influence of transfusions on perioperative and long-term outcome in patients following hepatic resection for colorectal metastases. *Ann Surg* 2003 Jun;237(6):860-9.
- (50) Poon RT, Fan ST, Lo CM, Liu CL, Lam CM, Yuen WK, et al. Improving perioperative outcome expands the role of hepatectomy in management of benign and malignant hepatobiliary diseases: analysis of 1222 consecutive patients from a prospective database. *Ann Surg* 2004 Oct;240(4):698-708.
- (51) De Boer MT, Molenaar IQ, Porte RJ. Impact of blood loss on outcome after liver resection. *Dig Surg* 2007;24(4):259-64.
- (52) Fattovich G, Giustina G, Degos F, Tremolada F, Diodati G, Almasio P, et al. Morbidity and mortality in compensated cirrhosis type C: a retrospective follow-up study of 384 patients. *Gastroenterology* 1997 Feb;112(2):463-72.
- (53) Boldt J. Clinical review: hemodynamic monitoring in the intensive care unit. *Crit Care* 2002 Feb;6(1):52-9.
- (54) Lam CM, Lo CM, Liu CL, Fan ST. Biliary complications during liver resection. *World J Surg* 2001 Oct;25(10):1273-6.
- (55) Sun HC, Qin LX, Wang L, Ye QH, Wu ZQ, Fan J, et al. Risk factors for postoperative complications after liver resection. *Hepatobiliary Pancreat Dis Int* 2005 Aug;4(3):370-4.
- (56) Walia A. Anesthetic management for liver resection. *J Gastrointest Surg* 2006 Feb;10(2):168-9.
- (57) Fallon MB, Abrams GA. Pulmonary dysfunction in chronic liver disease. *Hepatology* 2000 Oct;32(4 Pt 1):859-65.
- (58) Redai I, Emond J, Brentjens T. Anesthetic considerations during liver surgery. *Surg Clin North Am* 2004 Apr;84(2):401-11.

- (59) Jones RM, Moulton CE, Hardy KJ. Central venous pressure and its effect on blood loss during liver resection. *Br J Surg* 1998 Aug;85(8):1058-60.
- (60) Rees M, Plant G, Wells J, Bygrave S. One hundred and fifty hepatic resections: evolution of technique towards bloodless surgery. *Br J Surg* 1996 Nov;83(11):1526-9.
- (61) Melendez JA, Arslan V, Fischer ME, Wuest D, Jarnagin WR, Fong Y, et al. Perioperative outcomes of major hepatic resections under low central venous pressure anesthesia: blood loss, blood transfusion, and the risk of postoperative renal dysfunction. *J Am Coll Surg* 1998 Dec;187(6):620-5.
- (62) Jamieson GG, Corbel L, Champion JP, Launois B. Major liver resection without a blood transfusion: is it a realistic objective? *Surgery* 1992 Jul;112(1):32-6.
- (63) Wu CC, Ho WM, Cheng SB, Yeh DC, Wen MC, Liu TJ, et al. Perioperative parenteral tranexamic acid in liver tumor resection: a prospective randomized trial toward a "blood transfusion"-free hepatectomy. *Ann Surg* 2006 Feb;243(2):173-80.
- (64) Lutz JT, Valentin-Gamazo C, Gorlinger K, Malago M, Peters J. Blood-transfusion requirements and blood salvage in donors undergoing right hepatectomy for living related liver transplantation. *Anesth Analg* 2003 Feb;96(2):351-5, table.
- (65) Mariette D, Smadja C, Naveau S, Borgonovo G, Vons C, Franco D. Preoperative predictors of blood transfusion in liver resection for tumor. *Am J Surg* 1997 Apr;173(4):275-9.
- (66) Lentschener C, Roche K, Ozier Y. A review of aprotinin in orthotopic liver transplantation: can its harmful effects offset its beneficial effects? *Anesth Analg* 2005 May;100(5):1248-55.
- (67) Henry DA, Carless PA, Moxey AJ, O'Connell D, Stokes BJ, McClelland B, et al. Anti-fibrinolytic use for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2007;(4):CD001886.
- (68) Shirabe K, Shimada M, Gion T, Hasegawa H, Takenaka K, Utsunomiya T, et al. Postoperative liver failure after major hepatic resection for hepatocellular carcinoma in the modern era with special reference to remnant liver volume. *J Am Coll Surg* 1999 Mar;188(3):304-9.
- (69) Nelson DR, Lim HL, Oliver D, Qian KP, Davis GL, Lau JY. alpha-Glutathione S-transferase as a marker of hepatocellular damage in chronic hepatitis C virus infection. *Am J Clin Pathol* 1995 Aug;104(2):193-8.
- (70) Coppola N, De SG, Marrocco C, Scarano F, Scolastico C, Tarantino L, et al. *Helicobacter* spp. and liver diseases. *Infez Med* 2003 Dec;11(4):201-7.
- (71) Mannervik B. The isoenzymes of glutathione transferase. *Adv Enzymol Relat Areas Mol Biol* 1985;57:357-417.
- (72) Hayes JD, Pulford DJ. The glutathione S-transferase supergene family: regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit Rev Biochem Mol Biol* 1995;30(6):445-600.

- (73) Mannervik B, Awasthi YC, Board PG, Hayes JD, Di IC, Ketterer B, et al. Nomenclature for human glutathione transferases. *Biochem J* 1992 Feb 15;282 (Pt 1):305-6.
- (74) Beckett GJ, Hayes JD. Glutathione S-transferases: biomedical applications. *Adv Clin Chem* 1993;30:281-380.
- (75) Beckett GJ, Dyson EH, Chapman BJ, Templeton AJ, Hayes JD. Plasma glutathione S-transferase measurements by radioimmunoassay: a sensitive index of hepatocellular damage in man. *Clin Chim Acta* 1985 Feb 28;146(1):11-9.
- (76) Rees GW, Trull AK, Doyle S. Evaluation of an enzyme-immunometric assay for serum alpha-glutathione S-transferase. *Ann Clin Biochem* 1995 Nov;32 (Pt 6):575-83.
- (77) Corrigall AV, Kirsch RE. Glutathione S-transferase distribution and concentration in human organs. *Biochem Int* 1988 Mar;16(3):443-8.
- (78) Strange RC, Howie AF, Hume R, Matharoo B, Bell J, Hiley C, et al. The development expression of alpha-, mu- and pi-class glutathione S-transferases in human liver. *Biochim Biophys Acta* 1989 Dec 8;993(2-3):186-90.
- (79) Giannini E, Mastracci L, Botta F, Romagnoli P, Fasoli A, Risso D, et al. Liver iron accumulation in chronic hepatitis C patients without HFE mutations: relationships with histological damage, viral load and genotype and alpha-glutathione S-transferase levels. *Eur J Gastroenterol Hepatol* 2001 Nov;13(11):1355-61.
- (80) Dickson RC, Lauwers GY, Rosen CB, Cantwell R, Nelson DR, Lau JY. The utility of noninvasive serologic markers in the management of early allograft rejection in liver transplantation recipients. *Transplantation* 1999 Jul 27;68(2):247-53.
- (81) Prabhu KS, Reddy PV, Jones EC, Liken AD, Reddy CC. Characterization of a class alpha glutathione-S-transferase with glutathione peroxidase activity in human liver microsomes. *Arch Biochem Biophys* 2004 Apr 1;424(1):72-80.
- (82) Richard MJ, Portal B, Meo J, Coudray C, Hadjian A, Favier A. Malondialdehyde kit evaluated for determining plasma and lipoprotein fractions that react with thiobarbituric acid. *Clin Chem* 1992 May;38(5):704-9.
- (83) Schmidt CC, Suttner SW, Piper SN, Nagel D, Boldt J. Comparison of the effects of desflurane and isoflurane anaesthesia on hepatocellular function assessed by alpha glutathione S-transferase. *Anaesthesia* 1999 Dec;54(12):1207-11.
- (84) Chhibber A, Dziak J, Kolano J, Norton JR, Lustik S. Anesthesia care for adult live donor hepatectomy: our experiences with 100 cases. *Liver Transpl* 2007 Apr;13(4):537-42.
- (85) Martin JL, Plevak DJ, Flannery KD, Charlton M, Poterucha JJ, Humphreys CE, et al. Hepatotoxicity after desflurane anesthesia. *Anesthesiology* 1995 Nov;83(5):1125-9.
- (86) Berghaus TM, Baron A, Geier A, Lamerz R, Paumgartner G. Hepatotoxicity following desflurane anesthesia. *Hepatology* 1999 Feb;29(2):613-4.
- (87) Eger EI. New inhaled anesthetics. *Anesthesiology* 1994 Apr;80(4):906-22.

- (88) Durak I, Guven T, Birey M, Ozturk HS, Kurtipek O, Yel M, et al. Halothane hepatotoxicity and hepatic free radical metabolism in guinea pigs; the effects of vitamin E. *Can J Anaesth* 1996 Jul;43(7):741-8.
- (89) Carbonell LF, Nadal JA, Llanos MC, Hernandez I, Nava E, Diaz J. Depletion of liver glutathione potentiates the oxidative stress and decreases nitric oxide synthesis in a rat endotoxin shock model. *Crit Care Med* 2000 Jun;28(6):2002-6.
- (90) Rohm KD, Suttner SW, Boldt J, Schollhorn TA, Piper SN. Insignificant effect of desflurane-fentanyl-thiopental on hepatocellular integrity--a comparison with total intravenous anaesthesia using propofol-remifentanyl. *Eur J Anaesthesiol* 2005 Mar;22(3):209-14.
- (91) Ray DC, Robbins AG, Howie AF, Beckett GJ, Drummond GB. Effect of spinal anaesthesia on plasma concentrations of glutathione S-transferase. *Br J Anaesth* 2002 Feb;88(2):285-7.
- (92) Hartman JC, Pagel PS, Proctor LT, Kampine JP, Schmeling WT, Warltier DC. Influence of desflurane, isoflurane and halothane on regional tissue perfusion in dogs. *Can J Anaesth* 1992 Oct;39(8):877-87.
- (93) Mazze RI. Metabolism of the inhaled anaesthetics: implications of enzyme induction. *Br J Anaesth* 1984;56 Suppl 1:27S-41S.
- (94) Merin RG, Bernard JM, Doursout MF, Cohen M, Chelly JE. Comparison of the effects of isoflurane and desflurane on cardiovascular dynamics and regional blood flow in the chronically instrumented dog. *Anesthesiology* 1991 Mar;74(3):568-74.
- (95) Koblin DD. Characteristics and implications of desflurane metabolism and toxicity. *Anesth Analg* 1992 Oct;75(4 Suppl):S10-S16.
- (96) Ray DC, Drummond GB. Halothane hepatitis. *Br J Anaesth* 1991 Jul;67(1):84-99.
- (97) Tiainen P, Lindgren L, Rosenberg PH. Changes in hepatocellular integrity during and after desflurane or isoflurane anaesthesia in patients undergoing breast surgery. *Br J Anaesth* 1998 Jan;80(1):87-9.
- (98) Kenna JG, Jones RM. The organ toxicity of inhaled anesthetics. *Anesth Analg* 1995 Dec;81(6 Suppl):S51-S66.
- (99) Kenna JG, Van Pelt FNAM. The metabolism and toxicity of inhaled anaesthetic agents. *Anaesth Pharm Rev* 1994;2:29-42.
- (100) Murray JM, Phillips AS, Fee JP. Comparison of the effects of isoflurane and propofol on hepatic glutathione-S-transferase concentrations during and after prolonged anaesthesia. *Br J Anaesth* 1994 May;72(5):599-601.
- (101) Beckett GJ, Chapman BJ, Dyson EH, Hayes JD. Plasma glutathione S-transferase measurements after paracetamol overdose: evidence for early hepatocellular damage. *Gut* 1985 Jan;26(1):26-31.
- (102) Beckett GJ, Hayes PC, Hussey AJ, Bouchier IA, Hayes JD. Plasma glutathione S-transferase measurements in patients with alcoholic cirrhosis. *Clin Chim Acta* 1987 Oct 30;169(1):85-9.

- (103) Hussey AJ, Aldridge LM, Paul D, Ray DC, Beckett GJ, Allan LG. Plasma glutathione S-transferase concentration as a measure of hepatocellular integrity following a single general anaesthetic with halothane, enflurane or isoflurane. *Br J Anaesth* 1988 Feb;60(2):130-5.
- (104) Murray JM, Rowlands BJ, Trinick TR. Indocyanine green clearance and hepatic function during and after prolonged anaesthesia: comparison of halothane with isoflurane. *Br J Anaesth* 1992 Feb;68(2):168-71.
- (105) Piper SN, Haisch G, Kumle B, Walz GA, Breining T, Mattinger P, et al. [Effects of esmolol- and sodium nitroprusside-induced controlled hypotension on hepatocellular integrity in patients undergoing endonasal sinus surgery]. *Anesthesiol Intensivmed Notfallmed Schmerzther* 2003 Dec;38(12):781-6.
- (106) Miller R. *Anesthesia*. New York, NY: Churchill Livingstone; 2005.
- (107) Sneyd JR. Recent advances in intravenous anaesthesia. *Br J Anaesth* 2004 Nov;93(5):725-36.
- (108) Farid H, O'Connell T. Hepatic resections: changing mortality and morbidity. *Am Surg* 1994 Oct;60(10):748-52.
- (109) Wang WD, Liang LJ, Huang XQ, Yin XY. Low central venous pressure reduces blood loss in hepatectomy. *World J Gastroenterol* 2006 Feb 14;12(6):935-9.
- (110) Marsh B, White M, Morton N, Kenny GN. Pharmacokinetic model driven infusion of propofol in children. *Br J Anaesth* 1991 Jul;67(1):41-8.
- (111) Schnider TW, Minto CF, Shafer SL, Gambus PL, Andresen C, Goodale DB, et al. The influence of age on propofol pharmacodynamics. *Anesthesiology* 1999 Jun;90(6):1502-16.
- (112) Milne SE, Troy A, Irwin MG, Kenny GN. Relationship between bispectral index, auditory evoked potential index and effect-site EC50 for propofol at two clinical end-points. *Br J Anaesth* 2003 Feb;90(2):127-31.
- (113) Wakeling HG, Zimmerman JB, Howell S, Glass PS. Targeting effect compartment or central compartment concentration of propofol: what predicts loss of consciousness? *Anesthesiology* 1999 Jan;90(1):92-7.
- (114) Absalom AR, Kenny GN. Closed-loop control of propofol anaesthesia using bispectral index: performance assessment in patients receiving computer-controlled propofol and manually controlled remifentanyl infusions for minor surgery. *Br J Anaesth* 2003 Jun;90(6):737-41.
- (115) Schuttler J, Ihmsen H. Population pharmacokinetics of propofol: a multicenter study. *Anesthesiology* 2000 Mar;92(3):727-38.
- (116) Gepts E, Camu F, Cockshott ID, Douglas EJ. Disposition of propofol administered as constant rate intravenous infusions in humans. *Anesth Analg* 1987 Dec;66(12):1256-63.
- (117) Ecoffey C, Viviani X, Billard V, Cazalaa JB, Molliex S, Servin F, et al. [Target controlled infusion (TCI) anesthesia using propofol. Assessment of training and practice in the operating room]. *Ann Fr Anesth Reanim* 2001 Mar;20(3):228-45.

- (118) Servin FS. TCI compared with manually controlled infusion of propofol: a multicentre study. *Anaesthesia* 1998 Apr;53 Suppl 1:82-6.
- (119) Egan TD. Target-controlled drug delivery: progress toward an intravenous "vaporizer" and automated anesthetic administration. *Anesthesiology* 2003 Nov;99(5):1214-9.
- (120) Sevransky J. Clinical assessment of hemodynamically unstable patients. *Curr Opin Crit Care* 2009 Jun;15(3):234-8.
- (121) Polanco PM, Pinsky MR. Practical issues of hemodynamic monitoring at the bedside. *Surg Clin North Am* 2006 Dec;86(6):1431-56.
- (122) Block FE, Jr. What is heart rate, anyway? *J Clin Monit* 1994 Nov;10(6):366-70.
- (123) Mathews L. Paradigm shift in hemodynamic monitoring. *The Internet Journal of Anesthesiology* 2007;11:2.
- (124) Gan TJ, Soppitt A, Maroof M, el-Moalem H, Robertson KM, Moretti E, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 2002 Oct;97(4):820-6.
- (125) Laupland KB, Bands CJ. Utility of esophageal Doppler as a minimally invasive hemodynamic monitor: a review. *Can J Anaesth* 2002 Apr;49(4):393-401.
- (126) Dark PM, Singer M. The validity of trans-esophageal Doppler ultrasonography as a measure of cardiac output in critically ill adults. *Intensive Care Med* 2004 Nov;30(11):2060-6.
- (127) Cariou A, Monchi M, Joly LM, Bellenfant F, Claessens YE, Thebert D, et al. Noninvasive cardiac output monitoring by aortic blood flow determination: evaluation of the Sometec Dynemo-3000 system. *Crit Care Med* 1998 Dec;26(12):2066-72.
- (128) King SL, Lim MS. The use of the oesophageal Doppler monitor in the intensive care unit. *Crit Care Resusc* 2004 Jun;6(2):113-22.
- (129) Turner MA. Doppler-based hemodynamic monitoring: a minimally invasive alternative. *AACN Clin Issues* 2003 May;14(2):220-31.
- (130) Knirsch W, Kretschmar O, Tomaske M, Stutz K, Nagdyman N, Balmer C, et al. Comparison of cardiac output measurement using the CardioQP oesophageal Doppler with cardiac output measurement using thermodilution technique in children during heart catheterisation. *Anaesthesia* 2008 Aug;63(8):851-5.
- (131) Berton C, Cholley B. Equipment review: new techniques for cardiac output measurement--oesophageal Doppler, Fick principle using carbon dioxide, and pulse contour analysis. *Crit Care* 2002 Jun;6(3):216-21.
- (132) English JD, Moppett IK. Evaluation of a trans-oesophageal Doppler probe in awake subjects. *Anaesthesia* 2005 Jul;60(7):720-1.
- (133) Martorano PP, Barboni E, Buscema G, Di RA. A novel anaesthetical approach to patients with brugada syndrome in neurosurgery. *Case Rep Anesthesiol* 2013;2013:280826.

- (134) Sinclair S, James S FAU - Singer, Singer M. Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial.(0959-8138 (Print)).
- (135) Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem* 1974 Nov 25;249(22):7130-9.
- (136) Arslan M, Kurtipek O, Dogan AT, Unal Y, Kizil Y, Nurlu N, et al. Comparison of effects of anaesthesia with desflurane and enflurane on liver function. *Singapore Med J* 2009 Jan;50(1):73-7.
- (137) Singaravelu S, Barclay P. Automated control of end-tidal inhalation anaesthetic concentration using the GE Aisys Carestation. *Br J Anaesth* 2013 Apr;110(4):561-6.
- (138) Francese E. Usage of Reference Management Software at the University of Torino. *JLIS it*; Vol 4, No 2 (2013) 2013 Feb 1.
- (139) Boutron I, Guittet L, Estellat C, Moher D, Hróbjartsson Ar, Ravaud P. Reporting Methods of Blinding in Randomized Trials Assessing Nonpharmacological Treatments. *PLoS Med* 2007 Feb 20;4(2):e61.
- (140) Field A. Sample Size Calculation. In: Field A, editor. *Discovering Statistics Using SPSS*. 2 ed. London, California, New Delhi: SAGE Publications Ltd; 2006. p. 143-217.
- (141) Bland JM, Altman DG. Multiple significance tests: the Bonferroni method. *BMJ* 1995 Jan 21;310(6973):170.
- (142) Henriksen JH, Moller S. Cardiac and systemic haemodynamic complications of liver cirrhosis. *Scand Cardiovasc J* 2009 Aug;43(4):218-25.
- (143) Kawasaki T, Moriyasu F, Kimura T, Someda H, Fukuda Y, Ozawa K. Changes in portal blood flow consequent to partial hepatectomy: Doppler estimation. *Radiology* 1991 Aug;180(2):373-7.
- (144) Boermeester MA, Straatsburg IH, Houdijk AP, Meyer C, Frederiks WM, Wesdorp RI, et al. Endotoxin and interleukin-1 related hepatic inflammatory response promotes liver failure after partial hepatectomy. *Hepatology* 1995 Nov;22(5):1499-506.
- (145) Palmer RM. The discovery of nitric oxide in the vessel wall. A unifying concept in the pathogenesis of sepsis. *Arch Surg* 1993 Apr;128(4):396-401.
- (146) Boermeester MA, Houdijk AP, Straatsburg IH, van Noorden CJ, van Leeuwen PA. Organ blood flow after partial hepatectomy in rats: modification by endotoxin-neutralizing bactericidal/permeability-increasing protein (rBPI23). *J Hepatol* 1999 Nov;31(5):905-12.
- (147) Koblin DD, Eger EI, Johnson BH, Konopka K, Waskell L. I-653 resists degradation in rats. *Anesth Analg* 1988 Jun;67(6):534-8.
- (148) Takizawa D, Hiraoka H, Goto F, Yamamoto K, Horiuchi R. Human kidneys play an important role in the elimination of propofol. *Anesthesiology* 2005 Feb;102(2):327-30.

- (149) Chen CL, Chen YS, de Villa VH, Wang CC, Lin CL, Goto S, et al. Minimal blood loss living donor hepatectomy. *Transplantation* 2000 Jun 27;69(12):2580-6.
- (150) Smyrniotis V, Farantos C, Kostopanagiotou G, Arkadopoulos N. Vascular control during hepatectomy: review of methods and results. *World J Surg* 2005 Nov;29(11):1384-96.
- (151) Ray DC, Bomont R, Mizushima A, Kugimiya T, Forbes HA, Beckett GJ. Effect of sevoflurane anaesthesia on plasma concentrations of glutathione S-transferase. *Br J Anaesth* 1996 Sep;77(3):404-7.
- (152) Suttner SW, Schmidt CC, Boldt J, Huttner I, Kumle B, Piper SN. Low-flow desflurane and sevoflurane anesthesia minimally affect hepatic integrity and function in elderly patients. *Anesth Analg* 2000 Jul;91(1):206-12.
- (153) Ko JS, Gwak MS, Choi SJ, Kim GS, Kim JA, Yang M, et al. The effects of desflurane and propofol-remifentanyl on postoperative hepatic and renal functions after right hepatectomy in liver donors. *Liver Transpl* 2008 Aug;14(8):1150-8.
- (154) Tao KM, Yang LQ, Liu YT, Tao Y, Song JC, Wu FX, et al. Volatile anesthetics might be more beneficial than propofol for postoperative liver function in cirrhotic patients receiving hepatectomy. *Med Hypotheses* 2010 Dec;75(6):555-7.
- (155) Lendvay V, Draegni T, Rostrup M, Kirkeboen KA. Propofol/Remifentanyl Vs Desflurane/Fentanyl in Open Hemicolectomy Surgery. *Anesthe Clinic Res* 2010;1:102-7.

**Effect Of Target-Controlled Infusion Of Propofol-Fentanyl
Versus Desflurane On Hepatocellular Integrity In Cirrhotic
Patients Undergoing Major Hepatic Resection.**

تأثير التسريب الهداف لعقاري البروبوفول و الفنتانيل مقارنة
باستنشاق الديسفلورين على الكفاءة الحيوية لخلايا الكبد في
مرضى التليف الكبدي أثناء جراحات الاستئصال الجزئي للكبد.

Protocol of a thesis submitted
to the Faculty of Medicine
University of Alexandria
In partial fulfillment of the
requirements of the degree of
**Doctor in Anesthesia and
Surgical Intensive Care**

خطة بحث مقدمة
لكلية الطب
جامعة الإسكندرية
إيفاء جزئياً
لشروط الحصول على درجة
دكتور في التخدير
والعناية المركزة الجراحية

By


من

Amr Farouk Ismail El Safty

عمرو فاروق إسماعيل الصفتي

MBBCh, Alex.
Master of Anesthesia, Alex.
Department of Anesthesia
and Surgical Intensive Care
Faculty of Medicine
University of Alexandria
2012

بكالوريوس الطب والجراحة ، الإسكندرية
ماجستير التخدير، الاسكندرية
قسم التخدير
والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية
٢٠١٢



SUPERVISORS**المشرفون****Prof. Dr. Ahmed Mohamed El Attar****أ.د/ أحمد محمد العطار**

Professor of Anaesthesia and Surgical
Intensive Care,
Faculty of Medicine,
University of Alexandria.

أستاذ التخدير و العناية المركزة
الجراحية
كلية الطب
جامعة الإسكندرية

Prof. Dr. Khaled Ahmed Yassen**أ.د/ خالد أحمد يس**

Professor and head of Anaesthesia and
Surgical Intensive Care department,

أستاذ و رئيس قسم التخدير و العناية
المركزة الجراحية

National Liver Institute,
University of Menoufiya.

معهد الكبد القومي
جامعة المنوفية

Co- SUPERVISOR**المشرف المشارك****Dr. Ragab Saad Beltagy****د/ رجب سعد بلتاغي**

Lecturer of Anaesthesia and Surgical
Intensive Care

مدرس التخدير و العناية المركزة
الجراحية


Faculty of Medicine,
University of Alexandria.

كلية الطب
جامعة الإسكندرية

For his experience in cardiac
measurements.

وذلك لخبرته في قياسات القلب.





ASSISTANT RESEARCHER

Mennah Atef Abou Saeid

5th grade student,

Faculty of Medicine,

University of Alexandria.

E-mail:mena_atef@yahoo.com

Mobile:0115170601

الباحث المساعد

منة عاطف ابو سعيد

طالبة بالفرقة الخامسة

كلية الطب

جامعة الإسكندرية



INTRODUCTION

Hepatocellular carcinoma (HCC) ranks fifth among the most prevalent cancers in the world and is the third most common cause of cancer related mortality.⁽¹⁾ It is frequently found in patients with chronic liver disease resulting from infection with hepatitis C virus (HCV).⁽²⁾


Resection is still the first choice for treatment of HCC⁽³⁾ even at the intermediate or advanced stages.⁽⁴⁾ Radical resection for intermediate-advanced HCC is indicated as follows: (1) single HCC with large or huge tumor nodule, swelling outward, clear border and a pseudo capsule. (2) multiple HCC with 3 or fewer nodules localized in one lobe or segment of the liver.⁽⁵⁾

The body can cope with removal of up to two-thirds of the liver and up to 70%–80% of the normal liver can be resected with functional compensation occurring within weeks of an operation.⁽⁶⁾

Major liver resection may result in complications due to metabolic insufficiency, hemodynamic changes and impairment of pulmonary gas exchange, which could lead to hepatic and multiorgan dysfunction.⁽⁷⁾

The site and volume of planned resection must be carefully assessed in the preoperative period, this, in addition to the patient's co-morbidities.⁽⁸⁾

During or after the operation which may require blood transfusions or a further operation, blood loss remains one of the main predictors of both perioperative morbidity and mortality after


A handwritten signature in black ink, appearing to read 'R. F. El-Ni', followed by a large, stylized flourish or mark.

liver resection.⁽⁹⁾ More recent studies with larger numbers of both transfused and non-transfused patients have been able to confirm the detrimental effects of transfusion on the development of postoperative complications and perioperative death after liver resections.⁽⁹⁾

Target controlled infusion (TCI) devices allow the anesthetist to provide anesthesia by controlling the theoretical (predicted) concentration of the drug in the central compartment. Therefore, rapid changes in the depth of anesthesia are possible with similar ease to that achieved with inhalational anesthesia. But unlike inhalational anesthesia, where the end-tidal concentration of vapors can be monitored, on-line blood concentration monitoring is not practical at the present time.⁽¹⁰⁾ Also, (TCI) of opioids and propofol were advocated to provide anaesthesia for major liver resection aiming to fasten patient recovery with minimal stay in the postoperative intensive care.⁽¹¹⁾

On the other hand, Desflurane has been reported to be more innocent than other volatile agents. It is almost resistant to biotransformation and only 0.01% can be metabolized.⁽¹²⁾

After major liver resections in cirrhotic patients, also, postoperative hepatocellular integrity needed to be assessed through determination of glutathione S-transferase (GST) levels which seems to be a more specific indicator of liver injury.⁽¹³⁾ An increase in GST concentrations may be detected before changes occur in routine liver function tests (aminotransferase activity, bilirubin concentrations) due to their lack of specificity and sensitivity.⁽¹³⁾




Combination of advanced, minimally invasive, hemodynamic assessment and monitoring such as trans-esophageal doppler(TED) improves the perioperative management and outcome of those high-risk surgical patients.⁽¹⁴⁾

TED is easy to use and avoids the complications associated with other monitoring devices. It takes minutes to be inserted, correct signal is easy to achieve, decisions on fluid therapy, inotropes, pressors and dilators usage is facilitated.⁽¹⁵⁾

Several studies have demonstrated that intraoperative fluid optimization guided by TED significantly improves outcomes, as evidenced by a decreased length of stay ranging from 30% to 40%.More over, reductions in the incidence of post operative nausea and vomiting, a shorter recovery time of gut function, and resumption of enteral nutrition have also been reported in the volume optimized groups.⁽¹⁶⁾

Until now, there has been no available literatures or research-based evidences regarding the effect of both anesthetic techniques on the perioperative events occurring among cirrhotic patients undergoing major hepatic resection. Also, the impact of preoperative cirrhosis and the decision whether to use (TCI) using propofol opioids or desflurane needs to be more investigated.



AIM OF THE WORK

- To compare between Target-controlled infusion of propofol-fentanyl versus desflurane based anesthesia in major liver resection in cirrhotic patients as regards subclinical hepatic injury, renal affection, hemodynamic parameters and postoperative course.



Patients And Methods:


The study will involve fifty adult cirrhotic (child A) patients going to be admitted to the National Liver Institute, Menoufiya University, undergoing major liver resection and they will be categorized into two groups, target controlled infusion group (TCI) and desflurane group (Des).

The sample size was calculated by using Epi Info software with the power of 80%, $\alpha=0.05$. Randomization will be done by using the closed envelope technique.

A written informed consent will be taken from each patient. The study will be carried out after Research and Ethics Committee approval from both: Faculty of Medicine, Alexandria University and National Liver Institute, Menoufiya University, Egypt.

Exclusion criteria:

- Patients with a history of esophageal disease.
- Patients undergoing re-operation.
- Patients with a history of allergic reactions to drugs.
- Those who had undergone a recent anaesthesia (within the previous 7 days).
- Patients who bled profusely during their operation, who are haemodynamically unstable, or who need inotropic support.
- Patients with preoperative renal dysfunction.



Anaesthesia monitoring includes:

- Five leads Electrocardiography.
- Continuous invasive arterial blood pressure via radial artery (mmHg).
- Continuous central venous pressure via right internal jugular vein catheter (cmH₂O).
- Peripheral pulse oxymetry (%).
- Capnography to monitor end tidal (ET) CO₂(mmHg).
- Fraction inspired oxygen concentration (%).
- Urine output (ml/hour).
- Neuromuscular blockade (Train of four).
- Oesophageal temperature(degree centigrade).
- Entropy to monitor depth of anaesthesia.
(Anaesthesia Workstation, General Electric, Helsinki, Finland)
- Trans-esophageal Doppler (cardioQ) haemodynamic monitoring. (Deltex Medical, Chichester, UK).

Anaesthetic technique:

A 20 gauge intravenous cannula will be inserted in the non dominant hand and used for induction of anaesthesia. Afterwards, at least two large peripheral venous lines (16 gauge or larger), multi-lumen central venous catheter in the right internal jugular vein and an arterial catheter also in the non dominant hand (radial artery will be used after doing Allen test) will be inserted.



In group Des general anesthesia will be induced with fentanyl in a dose of (1microgram/kg), propofol (2 mg/kg) and rocuronium (1 mg/kg). End-tidal desflurane concentrations during the induction will be limited to 1 MAC. Endotracheal intubation will be done after complete muscle relaxation is confirmed by nerve stimulator and anesthesia will be maintained with a mixture of air, oxygen and Desflurane at a fresh gas flow of 1 litre/min. Ventilation will be controlled to maintain ETCO_2 between 32 and 36 mmHg (Anaesthesia Workstation, General Electric, Helsinki, Finland).

In group TCI immediately before induction of anesthesia with propofol, all subjects will receive fentanyl 3 $\mu\text{g}/\text{kg}$ for 30 seconds and will be followed by a continuous infusion of fentanyl 2 $\mu\text{g}/\text{kg}/\text{h}$ for 30 min, 1.5 $\mu\text{g}/\text{kg}/\text{h}$ from 31–150 min, and 1 $\mu\text{g}/\text{kg}/\text{h}$ until 30 min before skin closure⁽¹⁰⁾. Propofol will be administered with a syringe pump integrated with (controller and software) GE Healthcare Finland (Datex-Ohmeda, Helsinki, Finland). The propofol venous blood target concentration (C_t). for induction of anesthesia was set at 4 $\mu\text{g}/\text{ml}$ for the younger patients(less than fifty years) and 3 $\mu\text{g}/\text{ml}$ for the elderly (more than fifty years).⁽¹¹⁾ If anesthesia was not induced within 5 min, the C_t will be increased sufficiently to complete the induction of anesthesia. When consciousness will be lost, rocuronium 1mg/kg, will be given and trachea will be intubated.

During maintenance of anesthesia, the anesthetist will be able to increase or decrease end tidal desflurane concentration by 1% increments in (group Des), or the propofol (C_t). by 0.5 $\mu\text{g}/\text{ml}$




in (group TCI) at any time according to entropy and patients hemodynamics.

In both groups, Entropy will be kept between 40-60. If MAP or HR remained increased after 5 min supplemental dose of fentanyl (0.5 µg/ kg) will be given. Atropine 0.5 mg will be given intravenously if heart rate drops below 45 beats/min. Anesthetics will be decreased also in response to hypotension not responsive to replacement of intraoperative fluid loss or treatment of bradycardia. Additional rocuronium will be administered as appropriate in bolus doses and guided by nerve stimulator.

Intravenous crystalloids in the form of Ringer's acetate will be infused at a rate of 6 ml/kg/hr to replace fluid deficit and basal fluid requirements. Packed RBCs are administered to keep haematocrit more than 25%. Blood products are given, if at all, guided by the coagulation laboratory intraoperative findings. Urine output will be maintained at least 1 mL/kg/hour. Rapid infusion device is available for emergency use. Intra-operative normothermia will be maintained using a forced air-warming blanket (Bair Hugger; Arizant, United Kingdom), in addition to a warm intravenous fluid device.


Prophylaxis against deep venous thrombosis includes the use of elastic stockings and a sequential compression device for both lower limbs until early ambulation. Daily administration of subcutaneous low molecular weight heparin from the first postoperative day is practiced.

All patients are planned to be extubated on table at the end of surgery unless decided otherwise according to the



operative circumstances. At the end of surgery (whether extubated or not), all patients of both groups will be transferred to the intensive care unit (ICU).

Postoperative complications including bleeding, sepsis, nausea, vomiting, chest infection, and renal dysfunction will be monitored closely and will be dealt with appropriately.

A handwritten signature in black ink, consisting of several stylized, overlapping loops and strokes, positioned below the text.

MEASUREMENTS

1) Haemodynamic parameters will be monitored continuously and recorded before induction (t_0), immediately after induction before intubation (t_1), 15 min after the intubation (t_2), during dissection (t_3), during hepatic resection (t_4), and near the end of surgery (t_5).

- Heart rate (beat/min).
- Mean arterial blood pressure (mmHg).
- Central venous pressure (cmH₂O).
- (CO) cardiac output (L/min).
- (SV) stroke volume (ml)
- (SVR) systemic vascular resistance (dyn.sec.cm⁻⁵).
- (FTc) corrected flow time (msec).

2) Laboratory investigations :

a) Liver function tests will include:

-AST& ALT(U/L), total bilirubin(mg/dl), albumin(gm/dl),
Prothrombin(%).

-Glutathion S transferase (GST), (IU/ml).


b) Kidney functions tests will include serum urea & creatinine(mg/dl) and microalbumin in urine(μ gm/ml).

C) Metabolic parameters & electrolytes will include:

-Arterial blood gases.

-Serum electrolytes as sodium& potassium(meq/l).

Samples of GST will be kept at room temperature for almost one hour, then serum will be separated by centrifugation



for five minutes at 3000 r/min and 1ml aliquots will be frozen at -80 degree centigrade until analysis.

All laboratory samples will be collected preoperatively, immediately postoperative and 48 hours postoperatively.

3) Fluid balance (intake and output), intraoperatively and 24 hours postoperatively.

4) Time from discontinuation of inhalational agent until the trachea is extubated is recorded.

5) Modified Aldrete Scoring system for recovery score.⁽¹⁷⁾

6) Postoperative course will be assessed as follow:

a) The amount of transfused blood and blood products (fresh frozen plasma and/or platelets) in ml needed intraoperatively and during first 48 hours postoperatively.

b) Postoperative complications will be recorded .


c) Duration of ICU and hospital stay will be recorded.

7) The total amount of inhalational agent in (ml) used intraoperatively will be calculated automatically by using the Aisys® GE Healthcare Finland (Datex-Ohmeda, Helsinki, Finland) anaesthesia machine and then will be recorded. Also, the amount of propofol(ml) used in TCI group and the anesthetic cost(LE) in both groups according to the current international prices will be recorded.




Analysis Of The Results

The data will be recorded by both manual and automatic charting and most operative data will be recorded in computerized files. Data will be subjected for statistical analysis by appropriate statistical tests using the SPSS (software package for windows), and P value of <0.05 will be considered to be statistically significant.




Discussion

The results are going to be discussed, with reference to the previous literatures, in the aim of identifying the effects of target-controlled infusion of propofol-fentanyl versus desflurane on hepatocellular integrity in cirrhotic patients undergoing major hepatic resection.



References


- 1) Livraghi T, Kissaló HM, Line PD. Treatment options in hepatocellular carcinoma today. *Scandinavian Journal of Surgery* 2011;100: 22–9
- 2) Di Bisceglie A, Simpson L, Lotze M. Development of hepatocellular carcinoma among patients with chronic liver disease due to hepatitis C viral infection. *J Clin Gastroenterol* 1994;19:222–6.
- 3) Wu MC. Status and prospect of comprehensive treatment of liver cancer. *Zhonghua Gandan Waike Zazhi* 2006.12:1–4
- 4) Karasawa Y, Masaki T, Watanabe S, Kuriyama S, Maeta H. Significance of resection surgery in multidisciplinary treatment of advanced hepatocellular carcinoma with multiple intrahepatic lesions. *J Surg Oncol* 2003;82:98–103
- 5) Ruzzenente A, Capra F, Pachera S, Iacono C, Piccirillo G, Lunardi M, Pistoso S, Valdegamberi A, D'Onofrio M, Guglielmi. Is liver resection justified in advanced hepatocellular carcinoma? Results of an observational study in 464 patients. *J Gastrointest Surg* 2009;13:1313–20
- 6) Bennett, Blumgart L H. Assessment of hepatic reserve prior to hepatic resection, *J Hepatobiliary Pancreat Surg* 2005; 12:10–15.
- 7) Clus U, Niemann, John P, Roberts, Nuncy L, Ascher, und C, Spencer Y. Intraoperative hemodynamic and liver function in adult-to-adult living liver donors. *Liver Transplantation* 2002.8:1126–33.



- 8) Ann Walia. Anesthetic Management for Liver Resection. *The Society for Surgery of the Alimentary Tract* 2006; 10: 2-5.
- 9) Marieke T. de Boer I, Quintus Molenaar , Robert J. Porte Impact of Blood Loss on Outcome after Liver Resection. *Dig Surg* 2007; 24: 259-64
- 10) Yu-hong L , Jian-hong XU, Jian-jun Y, Jie T, Jian-guo XU. Predictive performance of 'Diprifusor' TCI system in patients during upper abdominal surgery under propofol/fentanyl anesthesia. *J Zhejiang Univ SCI* 2005;1:43-8.
- 11) Milne SE, Horgan PG, Kenny GN. Target-controlled infusions of propofol and remifentanyl with closed-loop anaesthesia for hepatic resection. *Anaesthesia*. 2002;57(1):93-9.
- 12) Tung D, Yoshida EM, Wang CS, Steinbrecher UP. Severe desflurane hepatotoxicity after colon surgery in an elderly patient . *Can J Anaesthesia* 2005;52:133-6.
- 13) Arslan M, Kurtipek O, Dogan A T, Inal Y, Kizil Y, Nurlu N, Kamaci S, Kavutcu M. Comparison of effects of anaesthesia with desflurane and enflurane on liver Function. *Singapore Med J* 2009;50(1):73-81.
- 14) Goddard NG, Menadue LT, Wakeling HG. A case for routine oesophageal Doppler fluid monitoring during major surgery becoming a standard of care. *Br J Anaesth* 2007; 99: 559-65.
- 15) Singer M. Oesophageal Doppler. *Current opinion in critical care* 2009; 15; 244-8
- 16) Wakeling HG, McFall MR, Jenkins CS, Woods WG, Miles WF, Barclay GR, Fleming SC. Intraoperative oesophageal Doppler

guided fluid management shortens postoperative hospital stay after major bowel surgery. *Br J Anaesth* 2005; 95: 634–42.

17) Aldrete JA. The post-anesthesia recovery score revisited. *J Clin Anesth* 1995;7:89–91

Three handwritten signatures in black ink, likely representing the authors or reviewers of the document.

الملخص العربي

المرجعية الدراسية:

انه من الواجب توخي الحذر عند اعطاء المرضى المصابون بتليف الكبد تخدير كلي عند استئصال جزء من الكبد للمحافظة علي وظائف الكبد و الكلي.

الهدف من البحث:

المقارنه بين التسريب بالتحكم المستهدف لعقاري البروبوفول و الفنتانيل (TCI) مقابل عقار الدسفلورين (Des) الاستشافي في الاستئصال الجزئي للكبد في مرضي التليف الكبدي.

طريقه البحث:

بعد موافقه لجنه الاخلاقيات تم تقسيم خمسون مريضاً مصابون بتليف الكبد تصنيف تشيك الف الي مجموعتين: مجموعه TCI ومجموعه Des كل مجموعه خمس و عشرون مريضاً.

في مجموعه TCI قبل البدء في عملية التخدير جميع المرضى تم اعطائهم عقار الفنتانيل ثلاثه ميكروجرام /كجم لمدته ثلاثون ثانيه و اعقبه تسريب مستمر للفنتانيل اثنان ميكروجرام /كجم/ساعه لمدته ثلاثون دقيقه و واحد و نصف ميكروجرام /كجم/ساعه من بعد ثلاثون دقيقه حتي مائه و خمسون دقيقه ثم واحد ميكروجرام /كجم/ساعه حتي ثلاثون دقيقه قبل اغلاق الجلد. وتم حقن عقار البروبوفول عن طريق مضخة الحقن المتكمله مع وحده تحكم البرمجيات و قد كان التركيز

المستهدف للبروبوفول في الدم اربعه ميكروجرام /لتر بالنسبه للمرضي الاصغر سنا و ثلاثه ميكروجرام /لتر للمسنين. اثناء استمرار التخدير طبيب التخدير كان قادراً علي زياده او نقصان عقار البروبوفول نصف ميكروجرام /لتر في مجموعه TCI و تركيز الدسفلوران بنسبه واحد % في اي وقت وفقاً لانتروبي المرضى بين اربعون الي ستون.

تم فحص وظائف الكبد و الكلي في الدم و نسبه الميكروالبومين في البول و نسبه انزيم الجلوتاثاين اس ترانسفيراز قبل العمليه و مباشره بعد العمليه و ثمانيه و اربعون ساعه بعد العمليه.

تم قياس العلامات القياسيه وكميه الدم و البلازم المنقول للمريض كذلك دلائل الافاقه و الاعراض الجانييه و مكوث المريض في العنايه المركزه او المستشفى كذلك الوقت حتي اخراج انبويه الحجزه وكذلك التكلفة الاقتصاديه.

النتائج:

حدث انخفاض ملحوظ في مقاومه الاجهزه الوعائيه في مجموعه TCI مقارنة بمجموعه Des (779±36 vs 836±8 dyn.sec.cm⁻⁵, P=0.000), من القلب (85±3 مقابل 91±4 mmhg و 76±7 مقابل 86±4 ملل. حيث P<0.01 علي التوالي).

بمقارنه وظائف الكبد وجد ارتفاع ملحوظ في قيمه كل من ALT&AST في المجموعتان مباشره بعد العمليه حيث كان في مجموعه Des (378±80 مقابل 407±35 U/L) وفي مجموعه TCI (413±39 مقابل 467±38 U/L) علي التوالي. و قد حدث هبوط في المجموعتان ثمانيه و اربعون ساعه بعد العمليه. ولا يوجد اي اختلاف ملحوظ في المجموعتان من حيث نسبه انزيم الجلوتاثاين اس ترانسفيراز و نسبه الميكروالبومين في البول وكذلك مكوث المريض فب العنايه المركزه او المستشفى.

الوقت حتي اخراج انبويه الحجزه كان اطول في مجموعه TCI (15.2±2.629) دقيقه مقابل (9.76±1.507) دقيقه في مجموعه Des حيث P=0.000. ومن حيث التكلفة الاقتصاديه كانت تكلفه مجموعه Des اقل من مجموعه TCI حيث كان (235±26) مقابل (438±57) جنيهاً مصرياً و كان (33.7±3.8) مقابل (62.65±8.233) دولار امريكي حيث P=0.000.

الاستنتاج:

عقار الدسفلوران الاستثنائي اختيار اصلح عن عقاري البروبوفول و الفنتانيل بالتسريب بالتحكم المستهدف في المرضى المصابون بتليف الكبد عند الاستئصال الجزئي للكبد من حيث استقرار العلامات الديناميكية و دلائل الفوقان و كذلك من حيث التكلفة الاقتصادية ولكنهما متماثلين من حيث التأثير علي وظائف الكبد و الكلي.

تأثير التسريب الهادف لعقاري البروبوفول و الفنتانيل مقارنة باستنشاق
الديسفلورين على الكفاءة الحيوية لخلايا الكبد في مرضى التليف الكبدي أثناء
جراحات الاستئصال الجزئى للكبد

رسالة علمية

مقدمة لكلية الطب – جامعة الإسكندرية
إيفاءاً جزئياً لشروط للحصول على درجة

الدكتوراه فى التخدير و العناية المركزة الجراحية

مقدمة من

عمرو فاروق إسماعيل الصفتي

بكالوريوس الطب والجراحة – الإسكندرية
ماجستير التخدير و العناية المركزة الجراحية

كلية الطب
جامعة الإسكندرية
٢٠١٥

تأثير التسريب الهادف لعقاري البروبوفول و الفنتانيل مقارنة باستنشاق الديسفلورين على الكفاءة الحيوية لخلايا الكبد في مرضى التليف الكبدي أثناء جراحات الاستئصال الجزئي للكبد

مقدمة من

عمرو فاروق إسماعيل الصفتي

بكالوريوس الطب والجراحة- الإسكندرية

للحصول على درجة

الدكتوراه في التخدير و العناية المركزة الجراحية

موافقون

لجنة المناقشة والحكم على الرسالة

.....

أ.د/ فوزية محمد أبو الفتوح
أستاذ التخدير و العناية المركزة الجراحية
كلية الطب
جامعة القاهرة

.....

أ.د/ أحمد محمد العطار
أستاذ التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

.....

أ.د/ نجوى محمود القبية
أستاذ التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

التاريخ:

السادة المشرفون

.....

أ.د/ احمد محمد العطار

أستاذ التخدير والعناية المركزة الجراحية

كلية الطب

جامعة الإسكندرية

.....

أ.د/ خالد احمد يس

أستاذ التخدير والعناية المركزة الجراحية

معهد الكبد القومى

جامعة المنوفية

.....

د/ رجب سعد بلتاجي

مدرس التخدير والعناية المركزة الجراحية

كلية الطب

جامعة الإسكندرية