

Advisors' Committee:

Prof. Dr. Ahmed Magdy Beda

Professor of Opthamology

Faculty of Medicine

University of Alexandria

.....

Dr. Amir Ramadan Ahmed

Lecturer of Opthamology

Faculty of Medicine

University of Alexandria

.....

ACKNOWLEDGMENT

First of all I thank **Allah** the most gracious for helping me to accomplish this research and for providing me with such very encouraging and supportive supervisors.

I would like to express my deepest gratitude to **Prof. Dr. Ahmed Magdy Beda**, who suggested this topic. I appreciate his unforgettable support as well as his generous efforts in the evaluation of this work. It is honorable to be supervised by him.

I am greatly indebted and grateful to **Dr. Amir Ramadan Ahmed**, for his fruitful suggestions, insistence on perfection together with the invaluable assistance he devoted in the supervision of this study.

LIST OF CONTENTS

Chapter	Page
ACKNOWLEDGMENT.....	i
LIST OF CONTENTS	ii
LIST OF TABLES	iii
LIST OF FIGURES	iv
LIST OF ABBREVIATIONS.....	vi
I. INTRODUCTION	1
II. AIM OF THE WORK	15
III. SUBJECTS AND METHODS	16
IV. RESULTS	17
V. DISCUSSION.....	32
VI. SUMMARY	34
VII. CONCLUSIONS	35
VIII. REFERENCES.....	36
PROTOCOL	
ARABIC SUMMARY	

LIST OF TABLES

Table		Page
(1)	Chorioretinal Diseases With Changes in the Choroidal Thickness	6
(2)	Comparison between the studied groups according to demographic data.....	17
(3)	Comparison between the studied groups according to side.....	19
(4)	Distribution of studied sample according to type in the diabetics group (n=20).....	20
(5)	Comparison between the studied groups according to horizontal choroidal thickness	21
(6)	Comparison between the studied groups according to vertical choroidal thickness.....	23
(7)	Comparison between Type of diabetic retinopathy according to Average Horizontal (H)(μm) and Average vertical (V)(μm) choroidal thickness	25

LIST OF FIGURES

Figure		Page
(1)	Optical coherence tomography image demonstrating image averaging using Cirrus HD-OCT (Carl Zeiss Meditec, Inc).....	3
(2)	Optical coherence tomography image demonstrating enhanced depth imaging on Spectralis (Heidelberg Engineering)	4
(3)	Optical coherence tomography image of the normal choroid taken on Spectralis with EDI and over sampling.	5
(4)	Optical coherence tomography image of the normal choroid taken on Cirrus HD-OCT with 20 images averaged	5
(5)	Optical coherence tomography image of the choroid of a patient with central serous chorioretinopathy taken on Cirrus HD-OCT with 20 images averaged.....	7
(6)	A. Optical coherence tomography image of the choroid of a patient with exudative AMD taken on Cirrus HD-OCT with 20 images averaged.....	8
(7)	Optical coherence tomography image of the choroid of a patient with high myopia taken on Cirrus HD-OCT (Carl Zeiss Meditec, Inc) with 20 images averaged.....	10
(8)	Optical coherence tomography image of the choroid of 2 patients with retinitis pigmentosa taken on Cirrus HD-OCT (Carl Zeiss Meditec, Inc) with 20 images averaged	12
(9)	Optical coherence tomography image of the choroid of a patient with late phase of retinitis pigmentosa taken on Cirrus HD-OCT (Carl Zeiss Meditec, Inc) with 20 images averaged.....	12
(10)	Comparison between the studied groups according to sex.....	18
(11)	Comparison between the studied groups according to age	18
(12)	Comparison between the studied groups according to side.....	19
(13)	Distribution of studied sample according to type in the diabetics group	20
(14)	Comparison between the studied groups according to Horizontal temporal (HT)(μm)and Horizontal foveal (HF)(μm) and Horizontal Nasal (HN)(μm) and average Horizontal (H)(μm) choroidal thickness	22
(15)	Comparison between the studied groups according to vertical temporal (VT)(μm) and vertical foveal (VF)(μm) and vertical nasal (VN) (μm) and average vertical (V)(μm) choroidal thickness	24
(16)	Comparison between Types of diabetic retinopathy according to Average Horizontal (H) ^(μm) choroidal thickness	26
(17)	Comparison between Type of diabetic retinopathy according to Average vertical (V)(μm) choroidal thickness.....	26
(18)	Comparison between Macular edema (NPDR/PDR) according to Average H	28

Figure	Page
(19) Comparison between Macular edema (NPDR/PDR) according to Average V	28
(20) (Optical coherence tomography image of the normal choroid taken on Spectralis with EDI and over sampling. The central subfoveal choroidal thickness was 491 μ in case A and 413 μ in cases B	29
(21) Optical coherence tomography image of the choroid of eyes with non proliferative diabetic retinopathy taken on Spectralis with EDI and over sampling. The central subfovea choroidal thickness was 222 μ in case A and 325 μ in case B	30
(22) Optical coherence tomography image of the choroid of an eye with non proliferative diabetic retinopathy with cystoid macular edema and vitromacular traction taken on Spectralis with EDI and over sampling. The central subfovea choroidal thickness was 287 μ	31
(23) Optical coherence tomography image of the choroid of an eye with severe proliferative diabetic retinopathy taken on Spectralis with EDI and over sampling. The central subfovea choroidal thickness was 212 μ	31

LIST OF ABBREVIATIONS

NPDR	Non Proliferative Diabetic Retinopathy
PDR	Proliferative Diabetic Retinopathy
CME	Cystoid Macular Edema
HT	Horizontal Temporal Choroidal Thickness
HF	Horizontal Foveal Choroidal Thickness
HN	Horizontal Nasal Choroidal Thickness
H	Horizontal Choroidal Thickness
VT	Vertical Temporal Choroidal Thickness
VF	Vertical Foveal Choroidal Thickness
VN	Vertical Nasal Choroidal Thickness
V	Vertical Choroidal Thickness
CSFT	Central sub field macular Thickness
CSFV	Central Sub Field Macular Volume
TV	Total Macular Volume