

# **RECOMMENDATIONS**

## RECOMMENDATIONS

- 1- As there is a strong relationship between NES and BMI, psychiatrists must not neglect the major role of treatment of obesity hand in hand with the treatment of depression and NES.
- 2- There is an emphasis for the need to identify individuals with the atypical depression as it was found that the atypical subtype of MDD is a strong predictor of obesity and thereafter, therapeutic measures to diminish the consequences of increased appetite during depressive episodes with atypical features are advocated.
- 3- Future studies are needed to explore weight changes and the relationship between NES and BMI as possible risk factors for each other, especially among depressed patients of all weight ranges.
- 4- Further researches are essential to augment our understanding of NES as it occurs in conjunction with depression in order to discover the effects and causes of NES among depression patients.
- 5- Further researches are needed to study the relationship between NES and antidepressant drugs for the assessment of their effects, and how NES might be better managed and/or prevented with their use.
- 6- In patients with combined NES and depression it is preferred to consider the effect of certain types of SSRIs in the treatment.
- 7- Avoid as much as possible certain antidepressants as mirtazapine, clomipramine and trazodone in treatment of NES among patient with depression as they may have some risk in inducing NES which will make the treatment difficult.
- 8- Educate patients and family members on the side effects of antidepressants (change in appetite, weight gain, risk of glucose intolerance, and lipid abnormalities).
- 9- Adjust, if possible, the patient's antidepressant treatment in accordance with his or her metabolic profile.

# REFERENCES

## REFERENCES

1. Colles SL, Dixon JB, O'Brien PE. Night eating syndrome and nocturnal snacking: association with obesity, binge eating and psychological distress. *Int J Obes (Lond)* 2007; 31: 1722-30.
2. Tanofsky-Kraff M, Yanovski SZ. Eating disorder or disordered eating? Non-normative eating patterns in obese individuals. *Obes Res* 2004; 12: 1361-6.
3. Stunkard A, Berkowitz R, Wadden T, Tanrikut C, Reiss E, Young L. Binge eating disorder and the night eating syndrome. *Int J Obes Relat Metab Disord* 1996; 20: 1-6.
4. Stunkard AJ, Grace WJ, Wolff HG. The night-eating syndrome; a pattern of food intake among certain obese patients. *Am J Med* 1955; 19: 78-86.
5. De Zwaan M, Burgard M, Schenck CH, Mitchell JE. Night time eating: a review of the literature. *Eur Eat Disord Rev* 2003; 11: 7-24.
6. Striegel-Moore RH, Dohm FA, Hook JM, Schreiber GB, Crawford PB, Daniels SR. Night eating syndrome in young adult women: prevalence and correlates. *Int J Eat Disord* 2005; 37: 200-6.
7. Kuldau J, Rand CS. The night eating syndrome in the morbidly obese. *Int J Eat Disord* 1986; 5: 143-8.
8. Rand CS, Kuldau JM. Morbid obesity: a comparison between a general population and obesity surgery patients. *Int J Obes Relat Metab Disord* 1993; 17: 657-61.
9. Rand CS, Macgregor AM, Stunkard AJ. The night eating syndrome in the general population and among postoperative obesity surgery patients. *Int J Eat Disord* 1997; 22: 65-9.
10. Birketvedt GS, Florholmen J, Sundsfjord J, Osterud B, Dinges D, Bilker W, et al. Behavioral and neuroendocrine characteristics of the night eating syndrome. *JAMA* 1999; 282: 657-63.
11. Rand CS, Kuldau JM. Eating patterns in normal weight individuals: bulimia, restrained eating and the night eating syndrome. *Int J Eat Disord*. 1986; 5: 75-84.
12. Adami GF, Meneghelli A, Scopinaro N. Night eating and binge eating disorder in obese patients. *Int J Eat Disord* 1999; 25: 335-8.
13. Gluck ME, Geliebter A, Satov T. Night eating syndrome is associated with depression, low self-esteem, reduced daytime hunger, and less weight loss in obese outpatients. *Obes Res* 2001; 9: 264-7.
14. Berry EM, Growdon JH, Wurtman JJ, Caballero B, Wurtman RJ. A balanced carbohydrate: Protein diet in the management of Parkinson's disease. *Neurology* 1991; 41: 1295-7.
15. Yokogoshi H, Wurtman RJ. Meal composition and plasma amino acid ratios: Effects of various proteins or carbohydrates, and of various protein concentrations. *Metabolism* 1986; 35:637-642.
16. Birketvedt GS, Sundsfjord J, Florholmen JR. The HPA-axis in the Night Eating Syndrome. *Am J Physiol Endocrinol Metab* 2002; 282: 366-9.

17. Kennedy SH, Garfinkel PE, Panti V, Costa D, Brown GM. Changes in melatonin levels but not cortisol levels are associated with depression in patients with depression and eating disorders. *Arch Gen Psychiatry* 1989; 46: 73-8.
18. Schenk CH, Mahowald MW. Review of nocturnal sleep-related eating disorders. *Int J Eat Disord* 1994; 16: 343-56.
19. Aronoff NJ, Geliebter A, Zammit G. Gender and body mass index are related to the night-eating syndrome in obese outpatients. *J Am Diet Assoc* 2001; 101: 102-4.
20. O'Reardon TP, Ringel BL, Dinges DF, Allison KC, Rogers NL, Martino NS, et al. Circadian eating and sleeping patterns in the night eating syndrome. *Obes Res* 2004; 12: 1789-96.
21. Allison KC, Ahima RS, O'Reardon JP, Dinges DF, Sharma V, Cummings DE, et al. Neuroendocrine profiles associated with energy intake, sleep, and stress in the night eating syndrome. *J Clin Endocrinol Metab* 2005; 90: 6214-7.
22. Thorpy MJ. Classification of sleep disorders. *J Clin Neurophysiol* 1990; 7: 67-81.
23. Spaggiari MC, Granella F, Parrino L, Marchesi C, Melli I, Terzano MG. Nocturnal eating syndrome in adults. *Sleep* 1994; 17: 339-44.
24. Schenk CH, Hurwitz TD, Bundlie SR, Mahowald MW. Sleep-related eating disorders: polysomnographic correlates of a heterogeneous syndrome distinct from daytime eating disorders. *Sleep* 1991; 14: 419-31.
25. Pawlow LA, O'Neil PM, Malcolm RJ. Night eating syndrome: effects of brief relaxation training on stress, mood, hunger, and eating patterns. *Int J Obes Relat Metab Disord* 2003; 27: 970-8.
26. Adami GF, Campostano A, Marinari GM, Ravera G, Scopinaro N. Night eating in obesity: a descriptive study. *Nutrition* 2002; 18: 587-9.
27. Stunkard AJ, Allison KC. Two forms of disordered eating in obesity: binge eating and night eating. *Int J Obes Relat Metab Disord* 2003; 27: 1-12.
28. Allison KC, Engel SG, Crosby RD, de ZM, O'Reardon JP, Wonderlich SA, et al. Evaluation of diagnostic criteria for night eating syndrome using item response theory analysis. *Eat Behav* 2008; 9: 398-407.
29. Powers PS, Perez A, Boyd F, Rosemurgy A. Eating pathology before and after bariatric surgery: a prospective study. *Int J Eat Disord* 1999; 25: 293-300.
30. Ceru-Bjork C, Andersson I, Rossner S. Night eating and nocturnal eating-two different or similar syndromes among obese patients. *Int J Obes Relat Metab Disord* 2001; 25: 365-72.
31. Napolitano MA, Head S, Babyak MA, Blumenthal JA. Binge eating disorder and night eating syndrome: psychological and behavioral characteristics. *Int J Eat Disord* 2001; 30: 193-203.
32. Goel N, Stunkard AJ, Rogers NL, Van Dongen HP, Allison KC, O'Reardon JP, et al. Circadian rhythm profiles in women with night eating syndrome. *J Biol Rhythms* 2009; 24: 85-94.
33. Rogers NL, Dinges DF, Allison KC, Maislin G, Martino N, O'Reardon JP, et al. Assessment of sleep in women with night eating syndrome. *Sleep* 2006; 29: 814-819.

34. de Zwaan M, Roerig DB, Crosby RD, Karaz S, Mitchell JE. Night time eating: a descriptive study. *Int J Eat Disord* 2006; 39: 224-32.
35. Geliebter A. New developments in binge eating disorder and the night eating syndrome. *Appetite* 2002; 39: 175-7.
36. Marshall HM, Allison KC, O'Reardon JP, Birketvedt G, Stunkard AJ. Night eating syndrome among non obese persons. *Int J Eat Disord* 2004; 35: 217-22.
37. Allison KC, Crow SJ, Reeves RR, West DS, Foreyt JP, Dilillo VG, et al. Binge eating disorder and night eating syndrome in adults with type 2 diabetes. *Obesity* 2007; 15: 1287-93.
38. Morse SA1, Ciechanowski PS, Katon WJ, Hirsch IB. Isn't this just bedtime snacking? The potential adverse effects of night-eating symptoms on treatment adherence and outcomes in patients with diabetes. *Diabetes Care* 2006; 29: 1800-4.
39. Grilo CM, Masheb RM. Night-time eating in men and women with binge eating disorder. *Behav Res Ther* 2004; 42: 397-407.
40. Lundgren JD, Allison KC, Stunkard AJ. Familial aggregation in the night eating syndrome. *Int J Eat Disord* 2006; 39: 516-8.
41. Lundgren JD, Rempfer MV, Brown CE, Goetz J, Hamera E. The prevalence of night eating syndrome and binge eating disorder among overweight and obese individuals with serious mental illness. *Psychiatry Res* 2010; 175: 233-6.
42. Lundgren JD, Allison KC, Crow S, O'Reardon JP, Berg KC, Galbraith J, et al. Prevalence of the night eating syndrome in a psychiatric population. *Am J Psychiatry* 2006; 163: 156-8.
43. Wadden TA, Phelan S. Assessment of quality of life in obese individuals. *Obes Res* 2002; 10(Suppl 1): 50S-7S.
44. Aschoff J. Circadian timing. *Ann NY Acad Sci* 1984; 423: 442-68.
45. Sinha MK, Ohannesian JP, Heiman ML, Kriauciunas A, Stephens TW, Magosin S, et al. Nocturnal rise of leptin in lean, obese, and non-insulin-dependent diabetes mellitus subjects. *J Clin Invest* 1996; 97: 1344-7.
46. Birketvedt GS, Sundsfjord J, Florholmen JR. Hypothalamic-pituitary-adrenal axis in the night eating syndrome. *Am J Physiol Endocrinol Metab* 2002; 282: E366-9.
47. Tholin S, Lindroos A, Tynelius P, Akerstedt T, Stunkard AJ, Bulik CM, et al. Prevalence of night eating in obese and nonobese twins. *Obesity (Silver Spring)* 2009; 17: 1050-5.
48. Root TL, Thornton LM, Lindroos AK, Stunkard AJ, Lichtenstein P, Pedersen NL, et al. Shared and unique genetic and environmental influences on binge eating and night eating: a Swedish twin study. *Eat Behav* 2010; 11: 92-8.
49. Vander Wal JS. Night eating syndrome: a critical review of the literature. *Clin Psychol Rev* 2012; 32: 49-59.
50. Vetrugno R, Manconi M, Ferini-Strambi L, Provini F, Plazzi G, Montagna P. Nocturnal eating sleep-related eating disorder or night eating syndrome? A videopolysomnographic study. *Sleep* 2006; 29: 949-54.
51. Fassino S. Sleep and eating disorders: a review. *G Ital Psicopatologia* 2005; 11: 354-62.

52. Allison KC, Lundgren JD, O'Reardon JP, Geliebter A, Gluck ME, Vinai P, et al. Proposed diagnostic criteria for night eating syndrome. *Int J Eat Disord* 2010; 43: 241-7.
53. Stunkard AJ, Allison KC, Geliebter A, Lundgren JD, Gluck ME, O'Reardon JP. Development of criteria for a diagnosis: lesson from the night eating syndrome. *Compr Psychiatry* 2009; 50: 391-9.
54. O'Reardon JP, Peshek A, Allison KC. Night eating syndrome: diagnosis, epidemiology and management. *CNS Drugs* 2005; 19: 997-1008.
55. Vinai P, Allison KC, Cardetti S, Carpegna G, Ferrato N, Masante D, et al. Psychopathology and treatment of night eating syndrome: a review. *Eat Weight Disord* 2008; 13: 54-63.
56. Allison KC, Lundgren JD, O'Reardon JP, Martino NS, Sarwer DB, Wadden TA, et al. The night eating questionnaire (NEQ): psychometric properties of a measure of severity of the night eating syndrome. *Eating Behav* 2008; 9: 62-72.
57. Stunkard AJ, Messick S. The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J Psychosom Res* 1985; 29: 71-83.
58. Wadden TA, Sarwer DB, Womble LG, Foster GD, McGuckin BG, Schimmel A. Psychosocial aspects of obesity and obesity surgery. *Surg Clin North Am* 2001; 81: 1001-24.
59. Vander Wal JS, Waller SM, Klurfeld DM, McBurney MI, Dhurandhar NY. Night eating syndrome: evaluation of two screening instruments. *Eat Behav* 2005; 6: 63-73.
60. O'Reardon JP, Stunkard AJ, Allison KC. Clinical trial of sertraline in the treatment of night eating syndrome. *Int J Eat Disord* 2004; 35: 16-26.
61. Allison KC, Tarves EP. Treatment of night eating syndrome. *Psychiatr Clin North Am* 2011; 34: 785-96.
62. American Academy of Sleep Medicine. *International Classification of Sleep Disorders: Diagnostic and Coding Manual*. 2<sup>nd</sup> ed. Westchester, IL: American Academy of Sleep Medicine, 2005.
63. Winkelman JW, Herzog DB, Fava M. The prevalence of sleep-related eating disorder in psychiatric and non-psychiatric population. *Psychol Med* 1999; 29: 1461-6.
64. Howell MJ, Schenck CH. Treatment of nocturnal eating disorders. *Curr Treat Options Neurol* 2009; 11: 333-9.
65. Howell MJ, Schenck CH, Crow SJ. A review of nighttime eating disorder. *Sleep Med Rev* 2009; 13: 23-34.
66. Eiber R, Friedman S. Correlation between eating disorders and sleep disturbances. *Encephale* 2001; 27: 429-4.
67. Miyaoka T, Yasukawa R, Tsubouchi K, Miura S, Shimizu Y, Sukegawa T, et al. Successful treatment of nocturnal eating/drinking syndrome with selective serotonin reuptake inhibitors. *Int Clin Psychopharmacol* 2003; 18: 175-7.
68. Stunkard AJ, Allison KC, Lundgren JD, Martino NS, Heo M, Etemad B, et al. A paradigm for facilitating pharmacotherapy at a distance: sertraline treatment of the night eating syndrome. *J Clin Psychiatry* 2006; 67: 1568-72.

69. O'Reardon JP, Allison KC, Martino NS, Lundgren JD, Heo M, Stunkard AJ. A randomized, placebo-controlled trial of sertraline in the treatment of night eating syndrome. *Am J Psychiatry* 2006; 163: 893-8.
70. Winkelman JW. Treatment of nocturnal eating syndrome and sleep-related eating disorder with topiramate. *Sleep Med* 2003; 4: 243-6.
71. Morgenthaler TI, Silber MH. Amnestic sleep-related eating disorder associated with Zolpidem. *Sleep Med* 2002; 3: 323-7.
72. Friedman S, Even C, Dardennes R, Guelfi JD. Light therapy, obesity, and night-eating syndrome. *Am J Psychiatry* 2002; 159: 875-6.
73. Friedman S, Even C, Dardennes R, Guelfi JD. Light therapy, nonseasonal depression, and night eating syndrome. *Can J Psychiatry* 2004; 49: 790.
74. Allison KC, Lundgren JD, Moore RH, O'Reardon JP, Stunkard AJ. Cognitive behavior therapy for night eating syndrome: a pilot study. *Am J Psychother* 2010; 64: 91-106.
75. Kubitz KA, Landers DM, Petruzzello SJ, Han M. The effects of acute and chronic exercise on sleep. A meta-analytic review. *Sports Med* 1996; 21: 277-91.
76. Youngstedt SD, O'Connor PJ, Dishman RK. The effects of acute exercise on sleep: a quantitative synthesis. *Sleep* 1997; 20: 203-14.
77. Carek PJ, Laibstain SE, Carek SM. Exercise for the treatment of depression and anxiety. *Int J Psychiatry Med* 2011; 41: 15-28.
78. Sandra S. *Depression: Questions You Have – Answers You Need*. People's Medical Society. ISBN 1997; 978-1-882606-14-6.
79. Katon W, Ciechanowski P. Impact of major depression on chronic medical illness. *J Psychosom Res* 2002; 53: 859–63.
80. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorder, Text Revision*. 4<sup>th</sup> ed. Washington, DC: American Psychiatric Association, 2000.
81. A Message From APA President Dilip Jeste, M.D., on DSM-5. Available at: <http://www.psychnews.org/files/DSM-message.pdf>.
82. O'Hara MW, Neunaber DJ, Zekoski EM. Prospective study of postpartum depression: prevalence, course, and predictive factors. *J Abnorm Psychol* 1984; 93: 158-71.
83. Kendell RE, Chalmers JC, Platz C. Epidemiology of puerperal psychoses. *Br J Psychiatry* 1987; 150: 662-73.
84. SIGN. *Postnatal Depression and Puerperal Psychosis: A National Clinical Guideline*. Scottish Intercollegiate Guidelines Network. 2002; 32.
85. USPSTF. Screening for depression in adults: U.S. preventive services task force recommendation statement. *Ann Intern Med* 2009; 151: 784-92.
86. ACOG Committee Opinion No. 343: psychosocial risk factors: perinatal screening and intervention. *Obstet Gynecol* 2006; 108: 469-77.
87. Peindl KS, Wisner KL, Hanusa BH. Identifying depression in the first postpartum year: guidelines for office-based screening and referral. *J Affect Disord* 2004; 80: 37-44.

88. Wisner KL, Parry BL, Piontek CM. Clinical practice. Postpartum depression. *N Engl J Med* 2002; 347: 194-9.
89. Beck CT. Postpartum depression: it isn't just the blues. *Am J Nurs* 2006; 106: 40-50.
90. Imsiragic AS, Begic D, Martic-Biocina S. Acute stress and depression 3 days after vaginal delivery--observational, comparative study. *Coll Antropol* 2009; 33: 521-7.
91. Scrandis DA, Sheikh TM, Niazi R, Tonelli LH, Postolache TT. Depression after delivery: risk factors, diagnostic and therapeutic considerations. *ScientificWorldJournal* 2007; 7: 1670-82.
92. Spinelli MG. Antepartum and postpartum depression. *J Gend Specif Med* 1998; 1: 33-6.
93. ACOG Committee on Practice Bulletins--Obstetrics. ACOG Practice Bulletin: Clinical management guidelines for obstetrician-gynecologists number 92, April 2008 (replaces practice bulletin number 87, November 2007). Use of psychiatric medications during pregnancy and lactation. *Obstet Gynecol* 2008; 111: 1001-20.
94. Marian E. Incorporating recognition and management of perinatal and postpartum depression into pediatric practice. *Pediatrics* 2010; 126: 1032-9.
95. Pignone MP, Gaynes BN, Rushton JL, Burchell CM, Orleans CT, Mulrow CD, et al. Screening for depression in adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2002; 136: 765-76.
96. Peeke PM, Chrousos GP. Hypercortisolism and obesity. *Ann N Y Acad Sci.* 1995;771:665-676.
97. Kyrou I, Chrousos GP, Tsigos C. Stress, visceral obesity, and metabolic complications. *Ann N Y Acad Sci.* 2006;1083:77-110.
98. Bjorntorp P. Visceral obesity: a 'civilization syndrome' *Obes Res.* 1993;1:206-222.
99. Nieuwenhuizen AG, Rutters F. The hypothalamic-pituitary-adrenal-axis in the regulation of energy balance. *Physiol Behav.* 2008;94:169-177.
100. Hotamisligil GS. Inflammation and metabolic disorders. *Nature.* 2006;444:860-867.
101. Lee SW, Tsou AP, Chan H, et al. Glucocorticoids selectively inhibit the transcription of the interleukin 1 beta gene and decrease the stability of interleukin 1 beta mRNA. *Proc Natl Acad Sci U S A.* 1988;85:1204-1208.
102. Cronstein BN, Kimmel SC, Levin RI, et al. A mechanism for the antiinflammatory effects of corticosteroids: the glucocorticoid receptor regulates leukocyte adhesion to endothelial cells and expression of endothelial-leukocyte adhesion molecule 1 and intercellular adhesion molecule 1. *Proc Natl Acad Sci U S A.* 1992;89:9991-9995.
103. Raadsheer FC, Hoogendijk WJ, Stam FC, et al. Increased numbers of CRH expressing neurons in the hypothalamic paraventricular nucleus of depressed patients. *Neuroendocrinology.* 1994;60:436-444.
104. O'Toole JM, Sekula LK, Rubin RT. Pituitary-adrenal cortical axis measures as predictors of sustained remission in major depression. *Biol Psychiatry.* 1997;42:85-89.
105. Hansen-Grant SM, Pariante CM, Kalin NH, et al. Neuroendocrine and immune system pathology in psychiatric disease. In: Schatzberg AF, Nemeroff CB, eds. *Textbook of Psychopharmacology*, 2nd ed. Washington, DC: American Psychiatric Press; 1998:171-194.

106. Holsboer F, Barden N. Antidepressants and hypothalamic-pituitary-adrenocortical regulation. *Endocr Rev.* 1996;17:187-205.
107. Sapolsky RM. Stress, glucocorticoids, and damage to the nervous system: the current state of confusion. *Stress.* 1996;1:1-19.
108. Kling MA, Roy A, Doran AR, et al. Cerebrospinal fluid immunoreactive corticotropin-releasing hormone and adrenocorticotropin secretion in Cushing's disease and major depression: potential clinical implications. *J Clin Endocrinol Metab.* 1991;72:260-271.
109. Nemeroff CB. Reduced corticotropin releasing factor binding sites in the frontal cortex of suicide victims. *Arch Gen Psychiatry.* 1988;45:577-579.
110. Glowa JR, Gold PW. Corticotropin releasing hormone produces profound anorexigenic effects in the rhesus monkey. *Neuropeptides.* 1991;18:55-61.
111. E Atlantis, M Baker. Obesity effects on depression: systematic review of epidemiological studies. *Int J Obes (London)* 2008; 32: 881–91.
112. Stunkard AJ, Faith MS, Allison KC. Depression and obesity. *Biol Psychiatry* 2003; 54: 330-7.
113. Bornstein SR, Schuppenies A, Wong ML, Licinio J. Approaching the shared biology of obesity and depression: the stress axis as the locus of gene-environment interactions. *Mol Psychiatry* 2006; 11: 892-902.
114. Bean MK, Stewart K, Olbrisch ME. Obesity in America: implications for clinical and health psychologists. *J Clin Psychol Med Settings* 2008; 15: 214-24.
115. Emery CF, Fondow MD, Schneider CM, Christofi FL, Hunt C, Busby AK, Gastric bypass surgery is associated with reduced inflammation and less depression: a preliminary investigation. *Obes Surg* 2007; 17: 759-63.
116. Shoelson SE, Herrero L, Naaz A. Obesity, inflammation, and insulin resistance. *Gastroenterology* 2007; 132: 2169-80.
117. Vaccarino V, Johnson BD, Sheps DS, Reis SE, Kelsey SF, Bittner V. et al. National Heart, Lung, and Blood Institute Depression, inflammation, and incident cardiovascular disease in women with suspected coronary ischemia: the National Heart, Lung, and Blood Institute-sponsored WISE study. *J Am Coll Cardiol* 2007; 50: 2044-50.
118. Bremner MA, Beekman AT, Deeg DJ, Penninx BW, Dik MG, Hack CE, et al. Inflammatory markers in late-life depression: results from a population-based study. *J Affect Disord* 2008; 106: 249-55.
119. Milaneschi Y, Corsi AM, Penninx BW, Bandinelli S, Guralnik JM, Ferrucci L. Interleukin-1 receptor antagonist and incident depressive symptoms over 6 years in older persons: the InCHIANTI Study. *Biol Psychiatry* 2009; 65: 973-8.
120. Pasquali R, Vicennati V. Activity of the hypothalamic-pituitary-adrenal axis in different obesity phenotypes. *Int J Obes Relat Metab Disord* 2000; 24(Suppl 2): S47-S9
121. Walker BR. Activation of the hypothalamic-pituitary-adrenal axis in obesity: cause or consequence? *Growth Horm IGF Res* 2001; 11(suppl A): S91-S5.
122. Belanoff JK, Kalehzan M, Sund B, Fleming Ficek SK, Schatzberg AF. Cortisol activity and cognitive changes in psychotic major depression. *Am J Psychiatry* 2001; 158: 1612-6.

123. Holsboer F, The corticosteroid receptor hypothesis of depression. *Neuropsychopharmacology* 2000; 23: 477-501.
124. Lee WJ, Lee YC, Ser KH, Chen JC, Chen SC. Improvement of insulin resistance after obesity surgery: a comparison of gastric banding and bypass procedures. *Obes Surg* 2008; 18: 1119-25.
125. Huber JD. Diabetes, cognitive function, and the blood-brain barrier. *Curr Pharm Des* 2008; 14: 1594-600.
126. Ajilore O, Haroon E, Kumaran S, Darwin C, Binesh N, Mintz J, et al. Measurement of brain metabolites in patients with type 2 diabetes and major depression using proton magnetic resonance spectroscopy. *Neuropsychopharmacology* 2007; 32: 1224-31.
127. Atlantis E, Ball K. Association between weight perception and psychological distress. *Int J Obes (Lond)* 2008; 32: 715-21.
128. Derenne JL, Beresin EV. Body image, media, and eating disorders. *Acad Psychiatry* 2006; 30: 257-61.
129. Hoek HW, van Harten PN, Hermans KM, Katzman MA, Matroos GE, Susser ES. The incidence of anorexia nervosa on Curacao. *Am J Psychiatry* 2005; 162: 748-52.
130. Beesdo K, Jacobi F, Hoyer J, Low NC, Höfler M, Wittchen HU. Pain associated with specific anxiety and depressive disorders in a nationally representative population sample. *Soc Psychiatry Psychiatr Epidemiol* 2010; 45: 89-104.
131. Gadalla T, Piran N. Psychiatric comorbidity in women with disordered eating behavior: a national study. *Women Health* 2008; 48: 467-84.
132. Binder, E.B., Salyakina, D., Lichtner, P., Wochnik, G.M., Ising, M., Pütz, B. et al. **Polymorphisms in FKBP5 are associated with increased recurrence of depressive episodes and rapid response to antidepressant treatment.** *Nat Genet.* 2004; 36: 1319–1325
133. Faith M.S., Matz P.E., and Jorge, M.A. **Obesity-depression associations in the population.** *J Psychosom Res.* 2002; 53: 935–942
134. Björntorp P. Do stress reactions cause abdominal obesity and comorbidities? *Obes Rev* 2001; 2: 73-86.
135. Björntorp P. The regulation of adipose tissue distribution in humans. *Int J Obes Relat Metab Disord* 1996; 20: 291-302.
136. Ishak WW, Ha K, Kapitanski N, Bagot K, Fathy H, Swanson B, et al. The impact of psychotherapy, pharmacotherapy, and their combination on quality of life in depression. *Harv Rev Psychiatry* 2011; 19: 277-89.
137. Qaseem A, Snow V, Denberg TD, Forcica MA, Owens DK. Using second-generation antidepressants to treat depressive disorders: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2008; 149: 725-33.
138. APA. Practice Guideline for the Treatment of Patients with Major Depressive Disorder. 3<sup>rd</sup> ed. American Psychiatric Association. 2010.
139. Henigsberg N, Mahableshwarkar AR, Jacobsen P, Chen Y, Thase ME. A randomized, double-blind, placebo-controlled 8-week trial of the efficacy and tolerability of multiple doses of Lu AA21004 in adults with major depressive disorder. *J Clin Psychiatry* 2012; 73: 953-9.

140. Moll E, Neumann N, Schmid-Burgk W, Stabl M, Amrein R. Safety and efficacy during long-term treatment with moclobemide. *Clin Neuropharmacol.* 1994;17 Suppl 1:S74-87.
141. Garland EJ, Remick RA, Zis AP. Weight gain with antidepressants and lithium. *J Clin Psychopharmacol* 1988; 8: 323-30.
142. Berken GH, Weinstein DO, Stern WC. Weight gain. A side-effect of tricyclic antidepressants. *J Affect Disord* 1984; 7: 133-8.
143. Zimmerman U, Kraus T, Himmerick H, Schuld A, Pollmacher T. Epidemiology, implications and mechanisms underlying drug-induced weight gain in psychiatric patients. *J Psych Research* 2003; 37: 193-220.
144. Frank E, Kupfer DJ, Bulik CM, Levenson JA. Imipramine and weight gain during the treatment of recurrent depression. *J Affect Disord* 1990; 20: 165-72.
145. Fogelson DL. Weight gain during Fluoxetine treatment. *J Clin Psychopharmacol.* 1991 11:220-1.
146. Benazzi F. Weight gain in depression remitted with antidepressants: pharmacological or recovery effect? *Psychother Psychosom.* 1998 Jul-Oct;67(4-5):271-4.
147. Fava M. Weight gain and antidepressants. *J Clin Psychiatry.* 2000;61 Suppl 11:37-41.
148. Harvey BH, Bouwer CD. Neuropharmacology of paradoxical weight gain with selective serotonin reuptake inhibitors. *Clin Neuropharmacol* 2000; 23: 90-7.
149. Fava M, Judge R, Hoog SL, Nilsson ME, Koke SC. Fluoxetine versus sertraline and paroxetine in major depressive disorder: changes in weight with long-term treatment. *J Clin Psychiatry* 2000; 61: 863-7.
150. Bouwer CD, Harvey BH. Phasic craving for carbohydrate observed with citalopram. *Int Clin Psychopharmacol* 1996; 11: 273-8.
151. Himmerich H, Fulda S, Schaaf L, Beitinger PA, Schuld A, Pollmächer T. Changes in weight and glucose tolerance during treatment with mirtazapine. *Diabetes Care* 2006; 29: 170.
152. Montgomery SA, Reimtz PE, Zivkov M. Mirtazapine versus amitriptyline in the long-term treatment of depression: a double-blind placebo-controlled study. *Int ClinPsychopharmacol* 1998; 13: 63-73.
153. Silverstone PH, Ravindran A. Once-daily venlafaxine extended release (XR) compared with fluoxetine in outpatients with depression and anxiety. Venlafaxine XR 360 Study Group. *J Clin Psychiatry.* 1999 Jan;60(1):22-8.
154. Jain AK, Kaplan RA, Gaade KM, Wadden TA, Allison DB, Brewer ER, Leadbetter RA, Richard N, Haight B, Jamerson BD, et al. Bupropion SR Vs. Placebo for weight loss in obese patients with depressive symptoms. *Obes Res.* 2002 61:1049-56.
155. Weisler RH, Johnston JA, Lineberry CG, Samara B, Branconnier RJ, Billow AA. Comparison of bupropion and trazodone for the treatment of major depression. *J Clin Psychopharmacol.* 1994 Jun;14(3):170-9.
156. Chouinard G. Bupropion and amitriptyline in the treatment of depressed patients. *J Clin Psychiatry.* 1983 May;44(5 Pt 2):121-9.

157. Feighner J, Hendrickson G, Miller L, Stern W. Double-blind comparison of doxepin versus bupropion in outpatients with a major depressive disorder. *J Clin Psychopharmacol*. 1986 Feb;6(1):27-32.
158. Feiger AD, Bielski RJ, Bremner J, Heiser JF, Trivedi M, Wilcox CS, et al. Double-blind, placebo-substitution study of nefazodone in the prevention of relapse during continuation treatment of outpatients with major depression. *Int Clin Psychopharmacol* 1999; 14: 19-28.
159. Sussman N. The implications of weight changes with antipsychotic treatment. *J Clin Psychopharmacol*. 2003 Jun;23(3 Suppl 1):S21-6.
160. Jelliffe DP, Jelliffe EFP, Zerfas A, Neumann GG. *Community nutritional assessment*. Oxford: Oxford University Press, 1989.
161. Obesity. preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000; 894: i-xii, 1-253.
162. Kotz S, Balakrishnan N, Read CB, Vidakovic B. *Encyclopedia of statistical sciences*. 2<sup>nd</sup> ed. Hoboken, N.J.: Wiley-Interscience, 2006.
163. Kirkpatrick LA, Feeney BC. *A simple guide to IBM SPSS statistics for version 20.0*. Student ed. Belmont, Calif.: Wadsworth, Cengage Learning, 2013.
164. Pattern SB, Williams JVA, Lavorato DH, Brown L, McLaren L, Eliasziw M. et al. Major depression, antidepressant medication and the risk of obesity. *Psychother Psychosom* 2009; 78: 182–6.
165. Simon GE, Von Korff M, Saunders K, Miglioretti DL, Crane PK, van Belle G, et al. Association between obesity and psychiatric disorders in the US adult population. *Arch Gen Psychiatry* 2006; 63: 824–30.
166. Rosmond R, Bjorntorp P. Psychiatric ill-health of women and its relationship to obesity and body fat distribution. *Obes Res* 1998; 6: 338–45.
167. Ma J, Xiao L. Obesity and depression in US women: results from the 2005–2006 National Health and Nutritional Exam Survey. *Obesity* 2010; 18: 347–53.
168. Heo M, Pietrobelli A, Fontaine KR, Sirey JA, Faith MS. Depressive mood and obesity in US adults: comparison and moderation by sex, age, and race. *Int J Obes* 2006; 30: 513–9.
169. Strine TW, Mokdad AH, Dube SR, Balluz LS, Gonzalez O, Berry JT, et al. The association of depression and anxiety with obesity and unhealthy behaviors among community-dwelling US adults. *Gen. Hosp. Psychiatry* 2008; 30: 127–37.
170. Penninx BW, Beekman AT, Honig A, Deeg DJ, Schoevers RA, van Eijk JT, et al. Depression and cardiac mortality: results from a community-based longitudinal study. *Arch Gen Psychiatry* 2001; 58: 221–7.
171. Ng TP, Niti M, Tan WC, Cao Z, Ong KC, Eng P. Depressive symptoms and chronic pulmonary disease. *Arch Intern Med* 2007; 167: 60–7.
172. Wells KB, Rogers W, Burman MA, Camp P. Course of depression in patients with hypertension, myocardial infarction or insulin-dependent diabetes. *Am J Psychiatry* 1993; 150: 632–8.

173. Rosmond R. Obesity and depression: same disease, different names. *Med Hypotheses* 2004; 62: 976–9.
174. Blaine B. Does depression cause obesity? A meta-analysis of longitudinal studies of depression and weight control. *J Health Psychol* 2008; 13: 1190–7.
175. Orhan FO, Ozer UG, Ozer A, Altunoren O, Celik M, Kareaslan MF. Night eating syndrome among patients with depression. *Isr J Psychiatry Relat Sci* 2011; 48: 212-7.
176. Kucukgoncu S, Tek C, Bestepe E, Musket C, Guloksuz S. Clinical features of night eating syndrome among depressed patients. *Eur Eat Disord Rev* 2014; 22: 102-8.
177. J. W. Newcomer, “Second-generation (atypical) antipsychotics and metabolic effects: a comprehensive literature review,” *CNS Drugs*, vol. 19, supplement 1, pp. 1–93, 2005.
178. Allison KC, Grilo CM, Masheb RM, Stunkard AJ. Binge eating disorder and night eating syndrome a comparative study of disordered and eating. *J Consult Clin Psychol* 2005; 73: 1107-15.
179. Calugi S, Dalle Grave R, Marchesini G. Night eating syndrome in class II-III obesity: metabolic and psychopathological features. *Int J Obes (Lond)* 2009; 33: 899-904.
180. Rand CS, Kuldau JM. Morbid obesity: a comparison between a general population and obesity surgery patients. *Int J Obes Relat Metab Disord* 1993; 17: 657-61.
181. Lundgren JD, Allison KC, O'Reardon JP, Stunkard AJ. A descriptive study of non-obese persons with night eating syndrome and a weight-matched comparison group. *Eat Behav* 2008; 9: 343-51.
182. Dalle Grave R, Calugi S, Ruocco A, Marchesini G. Night eating syndrome and Weight loss outcome in obese patient. *Int J Eat Disord* 2011; 44: 150-6.
183. Striegel-Moore RI~I, Franko DL, Thompson D, Affenito S, Kraemer HC. Night eating: prevalence and demographic correlates. *Obesity (Silver Spring)* 2006; 14: 139-47.
184. Greeno CG, Wing RR, Marcus MD. Nocturnal eating in binge eating disorder and matched-weight controls. *Int J Eat Disord* 1995; 18: 343-9.
185. Striegel-Moore RH, Dohm FA, Hook JM, Schreiber GB, Crawford PB, Daniels SR. Night eating syndrome in young adult women: prevalence and correlates. *Int J Eat Disord* 2005; 37: 200-6.
186. Lundgren JD, Williams KB, Heitmann BL. Nocturnal eating predicts tooth loss among adults: results from the Danish MONIVA study. *Eating Behaviors* 2010; 11: 170-4.
187. Striegel-Morre RH, Franko DL, Thompson D, Affenito S, May A, Kraemer HC. Exploring the typology of night eating syndrome. *Int J Eat Disord* 2008; 41: 411-8.
188. Gluck ME, Vent CA, Saibe AD, Votruba SB, Krakoff J. Higher 24.h respiratory eaters. *Obesity (Silver spring)* 2011; 19: 319-32.
189. Lasserre AM, Glaus J, Vandeleur CL, Marques-Vidal P, Vaucher J, Bastardot F, et al. Depression with atypical features and increase in obesity, body mass index, waist circumference, and fat mass A prospective, population-based study. *JAMA Psychiatry* 2014; 71: 880-8.

# **APPENDIX**

# NEQ

**1. How hungry are you usually in the morning?**

- Not at all       A little       Somewhat       Moderately       Very

**2. When do you usually eat for the first time?**

- Before 9 a.m.       9:01 a.m. to 12:00 p.m.       12:01 p.m. to 3:00 p.m.  
 3:01 p.m. to 6:00 p.m.       After 6:01 p.m.

**3. Do you have cravings or urges to eat snacks after supper but before bedtime?**

- Not at all       A little       Somewhat       Very much so       Extremely so

**4. How much control do you have over your eating between supper and bedtime?**

- Not at all       A little       Somewhat       Very much so       Complete

**5. How much of your daily food intake do you consume after suppertime?**

- 0%       1-25%       26-50%       51-75%       76-100%  
(none)      (Up to a quarter)      (About half)      (More than half)      (All)

**6. Are you currently feeling blue or down in the dumps?**

- Not at all       A little       Somewhat       Very much so       Extremely so

**7. When you are feeling blue, is your mood lower in the:**

- Early morning       Late morning       Afternoon       Early evening  
 Late evening / nighttime       Check if your mood doesn't change during the day

**8. How often do you have trouble getting to sleep?**

- Never       Sometimes       About half the time       Usually       Always

**9. Other than only to use the bathroom, how often do you get up at least once in the middle of the night?**

- None at all       Less than once a week       About once a week  
 More than once a week       Every night

**If 0 on # 9, Please stop here**

**10. Do you have cravings or urges to eat snacks when you wake up at night?**

- Not at all       A little       Somewhat       Very much so       Extremely so

**11. Do you need to eat in order to get back to sleep when you awake at night?**

- Not at all       A little       Somewhat       Very much so       Extremely so

**12. When you get up in the middle of the night, how often do you snack?**

- Never       Sometimes       About half the time       Usually       Always

**If 0 on # 12, Please stop here**

**13. When you snack in the middle of the night, how aware are you of your eating?**

- Not at all       A little       Somewhat       Very much so       Completely

**14. How much control do you have over your night-time eating?**

- Not at all       A little       Somewhat       Very much       Complete

# **PROTOCOL**

**STUDY OF THE PREVELANCE OF NIGHT EATING  
SYNDROME IN DEPRESSED PATIENTS AND ITS  
RELATION TO ANTIDEPRESSANT DRUGS**

دراسة معدل انتشار متلازمة تناول الطعام ليلا في مرضى الاكتئاب و علاقته  
بالعقاقير المضادة للاكتئاب

Protocol of a thesis submitted  
to the Faculty of Medicine  
University of Alexandria  
In partial fulfillment of the  
requirements of the degree of  
**Master of neuropsychiatry**

خطة بحث مقدمة  
لكلية الطب  
جامعة الإسكندرية  
إيفاء جزئياً  
لشروط الحصول على درجة  
الماجستير في أمراض المخ و الأعصاب و  
الطب النفسي

By

من

**Mai essam hamed rizk**  
MBBCh, Aîex.  
Visiting Resident  
El-Hadara University Hospital  
Départment of Neurology and Psychiatry  
Faculty of Medicine  
University of Alexandria  
2013

مي عصام حامد رزق  
بكالوريوس الطب والجراحة ، الإسكندرية  
طبيب مقيم زائر  
مستشفى الحضرة الجامعي  
قسم الأمراض العصبية و النفسية  
كلية الطب  
جامعة الإسكندرية

٢٠١٣



**SUPERVISORS****المشرفون****Prof. Dr. Tarek Kamal Molokhia**

Professor of Psychiatry,  
Faculty of Medicine,  
University of Alexandria.



أ.د/ طارق كمال ملوخية

أستاذ الطب النفسي  
كلية الطب  
جامعة الإسكندرية


**Dr. Ossama Abo-Elmagd Elkhoully**

Assistant Professor of Psychiatry,  
Faculty of Medicine,  
University of Alexandria.



د/ أسامة أبو المجد الخولي

أستاذ مساعد الطب النفسي  
كلية الطب  
جامعة الإسكندرية



3.

**ASSISTANT RESEARCHER**

**الباحث المساعد**

**Mohamed Ali Mohamed Mahmoud**

**محمد علي محمد محمود**

6<sup>th</sup> grade student

طالب بالفرقة السادسة

Faculty of Medicine,

كلية الطب

University of Alexandria.

جامعة الإسكندرية

**Mobile:** 01113302888

**E-mail:** mohammed.shendidy@facebook.com


## INTRODUCTION

The reciprocal links associating obesity (whether general or central obesity) and depression (whether major or atypical) has been reported in many clinical and epidemiological studies, and seem to correlate more in adulthood, especially among women.<sup>(1-6)</sup>

Generally, both are major public health problems of high prevalence, that share leading threats being both associated with a number of chronic conditions as cardiovascular disease, metabolic disorders and cancers.<sup>(7-9)</sup>

Among their interlinking relations, some literature view obesity to be the clinical manifestation of a subtype of depression similar to that of atypical depression, while others view that they are separate constructs, having an influencing effect on each other; even being a risk factor to the development of one another.<sup>(10-12)</sup>

For instance, as far as obesity is concerned, in a prospective cohort of 74,332 men and women, it was reported that those with raised body mass index (BMI) were associated with an increased risk of depression,<sup>(13)</sup> while in another study elderly men with BMI over 30 had a 31% increased risk of developing depression.<sup>(14)</sup> In obese adolescences followed over a 4-year period, symptoms of depression were significantly related to increases in BMI for girls but not boys.<sup>(15)</sup> However, still, concrete data cannot as yet distinguish whether obesity is a consequence of, a contributor to, or a correlate of psychological disorders when linked to depression.<sup>(13-15)</sup>





Vice versa, from depression perspectives, combining data from 16 studies, confirmed that, depressed compared to non depressed people were at significantly higher risk for developing obesity<sup>(16)</sup> and that the risk among depressed people for later obesity was particularly high for adolescent females. This later was reported also in another study,<sup>(17)</sup> declaring that major depression in late adolescent girls was associated with a 2.3-fold increased risk of obesity in adulthood.<sup>(16,17)</sup>

Explanations have suggested some genetic and/or environmental cues that permit hypothalamic pituitary dysregulation, decrease physical activity, change the eating habits (especially become night eaters) use of antidepressants (more with tricyclic, monoaminoxidase inhibitors and some newer antidepressants), can all increase the risk of metabolic dysregulation in depressed individuals, ending in obesity.<sup>(18-20)</sup>

In this context, circadian desynchrony i.e disruption of circadian rhythmicity of the biological clock that fine tunes the sleep/wake cycle, has recently been nailed to the etiopathogenic links of depression, behavioral eating disorders specially night eating syndrome (NES) and obesity. Indeed, the association between NES and depressive symptoms in particular was noted in literature and has ranked up to 70% of NES subjects reporting a depressed mood in one of the studies.<sup>(21-24)</sup>

It's worth noting that depression in NES follows an atypical pattern with distinctive circadian features where by mood is usually better in the morning and worse during the evening and night, unlike depression which is worse in the morning and improves throughout the day.<sup>(10,21,24)</sup>




The denominator bottom line theme found in common in such association was the existence of hypothalamic-pituitary neurochemical dysregulation involving transmitters (monaminergic, gabergeric,...etc), hormones (cortisol, insulin, leptin...etc), and cytokines and neuropeptides (interleukines, adiponectin,...etc).<sup>(25-27)</sup>

From all aforementioned etiopathogenic raised links it was of interest to investigate in this study the relation between depression and obesity especially in subsets of patient presenting with NES and to correlate these findings to their anti-depressant drug treatment.

## AIM OF THE WORK

*This work aims to:*

- 1- Evaluate the prevalence of night eating syndrome in patients with depression.
- 2- Study the relationship between night eating syndrome and antidepressant drugs.



## SUBJECTS


After approval of Ethical Committee of Faculty of Medicine and written informed consent from patients, this study will be carried out among psychiatric outpatients in Alexandria Main University Hospitals over the period from February 2013 to July 2013 and patients will be divided into 2 groups (naïve and others receiving antidepressant therapy for at least 2 months)

### Inclusion criteria

1. Patients diagnosed as having depression according to DSM "IV" TR criteria.
2. Patients aged from 18 – 60 years.
3. Both sexes are included.
4. Patients will be grouped into naïve and others receiving antidepressant therapy for at least 2 months.
5. Patients must be accepting by informed consent.

### Exclusion criteria

1. Presence of organic illness contributes to either obesity or depression eg. Thyroid dysfunction.
2. Refusal of participation (none consenting).
3. Depression with psychotic features.
4. Patients receiving more than one antidepressant.
5. Other axis I or II disorders according to DSM "IV" TR.



## METHODS

All studied sample will be subjected to the following tools and techniques after the intake of consent:

1. Structured interview to collect:
  - A. Sociodemographic data as age, residence, educational level, marital status, occupation and socioeconomic status.
  - B. Medical, psychiatric and drug history.
  - C. Family history of psychiatric illness.
  - D. Onset of increase in weight and its relation to the onset of antidepressant intake.
  - E. The type of antidepressant in use.
  - F. Patterns of eating adopted.
  - G. Presence or absence of sleep disorders.
2. Clinical psychiatric assessment and diagnosis based on DSM-I VTR.
3. Weight and height measurements to define BMI.


Accordingly patients will be categorized into:

- Very obese: from 35 and above.
  - Obese: from 30 to 34.9.
  - Overweight: from 25 to 29.9.
4. Analysis by Arabic version of night eating questionnaire to define whether or not he is NES.<sup>(28)</sup>




## RESULTS

The results obtained from this study will be tabulated and statistically analyzed using the standard statistical methods.

A handwritten signature in black ink, appearing to be 'S. M. A.', written over a horizontal line.A handwritten signature in black ink, consisting of a stylized, cursive-like mark.

## DISCUSSION

The results obtained from this study will be discussed in view of achievement of the aim and compared with any available published data in the same field of the research.


A handwritten signature in black ink, appearing to be 'S. M. A.', written in a cursive style.A handwritten signature in black ink, appearing to be 'P.', written in a cursive style.

## REFERENCES


1. Pattern SB, Williams JVA, Lavorato DH. Major depression, antidepressant medication and the risk of obesity. *Psychother Psychosom* 2009; 78: 182–6.
2. Simon GE, Von Korff M, Saunders K. Association between obesity and psychiatric disorders in the US adult population. *Arch Gen Psychiatry* 2006; 63: 824–30.
3. Rosmond R, Bjorntorp P. Psychiatric ill-health of women and its relationship to obesity and body fat distribution. *Obes Res* 1998; 6: 338–45.
4. Ma J, Xiao L. Obesity and depression in US women: results from the 2005–2006 National Health and Nutritional Exam Survey. *Obesity*. 2010; 18: 347–53.
5. Heo M, Pietrobelli A, Fontaine KR, Sirey JA, Faith MS. Depressive mood and obesity in US adults: comparison and moderation by sex, age, and race. *Int J Obest* 2006; 30: 513–19.
6. Strine TW, Mokdad AH, Dube SR, Balluz LS, Gonzalez O, Berry JT, et al. The association of depression and anxiety with obesity and unhealthy behaviors among community-dwelling US adults. *Gen. Hosp. Psychiatry* 2008; 30: 127–37.
7. Penninx BW, Beekman AT, Honig A. Depression and cardiac mortality: results from a community-based longitudinal study. *Arch Gen Psychiatry* 2001; 58: 221–7.

8. Ng TP, Niti M, Tan WC. Depressive symptoms and chronic pulmonary disease. *Arch Intern Med* 2007; 167: 60–7.
9. Wells KB, Rogers W, Burman MA. Course of depression in patients with hypertension, myocardial infarction or insulin-dependent diabetes. *Am J Psychiatry* 1993; 150: 632–8.
10. Rosmond R. Obesity and depression: same disease, different names. *Med Hypotheses* 2004; 62: 976–9.
11. Blaine B. Does depression cause obesity?: A meta-analysis of longitudinal studies of depression and weight control. *J Health Psychol* 2008; 13: 1190–7.
12. Atlantis E, Baker M. Obesity effects on depression: systematic review of epidemiological studies. *Int J Obes (London)* 2008; 32: 881–91.
13. Bjerkese et O, Romundstad P, Evans J, Gunnell D. Association of adult body mass index & height with anxiety, depression and suicide in the general population; the HUNT study. *Am J Epidemiol* 2008; 167: 193-202.
14. Almeida SP, Calver J, Jamrozik K. Obesity and metabolic syndrome increase the risk of incident depression in older men: the Health in Men Study. *Am J Geriatr Psychiatry* 2009; 17: 889–98.
15. Boutelle KN, Hannan P, Fulkerson JA, Crow SJ, Stice E. Obesity as a prospective predictor of depression in adolescent females. *Health Psychol* 2010; 29: 293-8.



- 
16. Blaine B. Does depression cause obesity? : A meta analysis of longitudinal studies of depression & weight control. *J Health Psychol* 2008; 13: 1190-7.
  17. Richardson LP, Davis R, Poulton R. A longitudinal evaluation of adolescent depression and adult obesity. *Archives of Pediatrics and Adolescent Medicine* 2003; 157: 739-45.
  18. Pattern SB, Williams JVA, Lavorato DH. Major depression, antidepressant medication and the risk of obesity. *Psychother Psychosom* 2009; 78: 182-6.
  19. Vicennati V, Pasquali R. Abnormalities of the hypothalamic-pituitary-adrenal axis in nondepressed women with abdominal obesity and relations with insulin resistance: evidence for a central and a peripheral alteration. *Obes Res* 1996; 4: 245-52.
  20. Bjorntorp P. Do stress reactions cause abdominal obesity and comorbidities? *Obes Res* 2001; 2: 73-86.
  21. Gluck ME, Geliebter A, Satov T. Night eating syndrome is associated with depression, low self-esteem, reduced daytime hunger, and less weight loss in obese outpatients. *Obes Res* 2001; 9: 264-7.
  22. O'Reardon JP, Ringel BL, Dinges DF, Allison KC, Rogers NL, Martino NS, et al. Circadian eating and sleeping patterns in the night eating syndrome. *Obes Res* 2004; 12: 1789-96.
  23. Allison KC, Ahima RS, O'Reardon JP, Dinges DF, Sharma V, Cummings DE, et al. Neuroendocrine profiles associated with energy intake, sleep, and stress in the night eating syndrome. *J Clin Endocrinol Metab* 2005; 90: 6214-7.
- 
- 

24. Birketvedt GS, Florholmen J, Sundsfjord J, Osterud B, Dinges D, Bilker W, et al. Behavioral and neuroendocrine characteristics of the night-eating syndrome. *JAMA* 1999; 282: 657-63.
25. Marques-Deak AH, Neto FL, Dominguez WV. Cytokine profiles in women with different subtypes of major depressive disorder. *J Psychiatr Res* 2007; 41: 152-9.
26. Rosmond R, Chagnon YC, Holm G, Chagnon M, Perusse L, Lindell K, et al. A glucocorticoid receptor gene marker is associated with abdominal obesity, leptin, and dysregulation of the hypothalamic-pituitary-adrenal axis. *Obes Res* 2000; 8: 211-8.
27. Ahima RS, Qi Y, Singhal NS, Jackson MB, Scherer PE. Brain adipocytokine action and metabolic regulation. *Diabetes*. 2006; 55: S145-S54.
28. Sylvia, Escott-Stump. Night eating syndrome indicators and questionnaire. In: David BT, Matthew JH (eds). *Nutrition and diagnosis-related care* 6<sup>th</sup> ed. USA, Maryland: Lippincott William & Wilkins; 2008. pp 557.



# **ARABIC SUMMARY**

## الملخص العربي

إن العلاقة المتبادلة التي تربط السمنة (سواء العامة أو المركزية) بالاكتئاب قد وردت في العديد من الدراسات و يبدو انها ارتبطت أكثر بسن البلوغ و خاصة بين النساء.

عموما، كلاهما من الأسباب الضخمة لمشاكل صحية عالية الانتشار، و التي تشترك في تهديدات صحية، كونهما يرتبطان بالعديد من الأمراض المزمنة مثل أمراض القلب و الأوعية الدموية، اضطرابات التمثيل الغذائي و السرطان.

في اطار هذه الروابط المتبادلة، البعض يرى السمنة أنها مظهر اكلينيكي لنوع فرعي من أنواع الاكتئاب مثل الاكتئاب الغير نمطي، في حين يرى البعض الآخر أنهما منفصلان بينما لهما تأثير على بعضهما البعض أو كون أحدهما عامل مسبب لتطور الآخر.

مؤخرا، دخلت متلازمة الأكل الليلي هذه الحلقة المفرغة كونها مرتبطة بقوة مع كل من السمنة و الاكتئاب في علاقة ثنائية الاتجاه مع كل واحد منهما.

لوحظت متلازمة تناول الطعام ليلا لأول مرة بين المرضى الذين يعانون من السمنة، كما تم دعم العلاقة بينهما مع تقديرات تشير الى أنها أكثر شيوعا بين الأشخاص البدناء (٦-١٦%) مقارنة مع عموم الناس (١.٥%).

و فيما يتعلق بعلاقة متلازمة تناول الطعام ليلا مع الاكتئاب، فانه ليس من الواضح ما اذا كان الاكتئاب هو سبب، هو نتيجة أو سمة من سمات متلازمة تناول الطعام ليلا.

على الرغم من أنه تم اجراء العديد من الدراسات على العلاقة بين متلازمة تناول الطعام ليلا و الاكتئاب، الا أن معظمهم كان من منظور متلازمة تناول الطعام ليلا. و في الوقت الحاضر هناك قليل من الدراسات بشأن معدل متلازمة تناول الطعام ليلا في مرضى الاكتئاب و علاقتها بمضادات الاكتئاب.

لذلك أجري هذا البحث لتقدير معدل انتشار متلازمة تناول الطعام ليلا بين مرضى الاكتئاب و علاقته بالعقاقير المضادة للاكتئاب. و قد أجريت هذه الدراسة على ٤٠٠ مريض نفسي تتراوح أعمارهم بين ١٨-٦٠ عاما، من الجنسين في مستشفيات جامعة الاسكندرية خلال الفترة من فبراير ٢٠١٣ حتى يوليو ٢٠١٣، و قد تم تقسيم المرضى الى مجموعتين متساويتين (٢٠٠ مريض في كل مجموعة). واحدة على مضى الاكتئاب دون علاج و الأخرى على مرضى الاكتئاب و على العلاج لمدة شهرين على الأقل.

### وقد خضع المشاركون لما يلي:

- ١- استبيان معد ذاتيا لجمع كافة المعلومات المتعلقة بالامور الاجتماعية و الديموغرافية و الصحية و النفسية و التاريخ الدوائي بما في ذلك نوع مضادات الاكتئاب المستخدمة، بداية زيادة الوزن و علاقته ببداية العلاج.
- ٢- التقييم النفسي على أساس (DSM-IV) لتشخيص اضطراب الاكتئاب و استبعاد اضطرابات محور ١ (AXIS I) الأخرى.
- ٣- تم حساب القياسات الأنتروبومترية (قياسات الجسم البشري) و مؤشر كتلة الجسم.
- ٤- مقابلة نفسية لجميع الأشخاص الذين تم تحليلهم باستخدام معايير التشخيص المقترحة لمتلازمة تناول الطعام ليلا و هو ما كان مؤكدا باستبيان تناول الطعام ليلا الذي تم تعريبه و من ثم تم جمع درجات جميع العناصر ما عدا سؤال ١٣ (عن الوعي اثناء تناول الطعام) للوصول الى الدرجة الكلية حيث تتراوح بين ٠-٥٢. و كانت الدرجة القطعية المستخدمة في هذه الدراسة من ٢٥.

### ولقد اظهرت الدراسة النتائج التالية:

- ٢٦.٥% من الحالات في المجموعة أ تم تشخيصهم بمتلازمة تناول الطعام ليلا في حين تم تشخيص ٩% فقط من حالات المجموعة ب.
- في كل مجموعة، كانت حالات زيادة الوزن و السمنة أعلى احصائيا من الحالات ذات الوزن الطبيعي.
- فيما يتعلق بمؤشر كتلة الجسم، كان هناك فروق ذات دلالة احصائية بين المجموعتين ( $p < 0.001$ ) كما كانت الفروق بين حالات زيادة الوزن و السمنة بين المجموعتين ذات دلالة احصائية (٨٣% و ٥٧%) على التوالي.
- حالات متلازمة تناول الطعام ليلا الايجابية في كلتا المجموعتين كانت ذات سن أصغر من الحالات السلبية.

• الفروق بين الحالات السلبية و الايجابية لمتلازمة الأكل الليلي فيما يتعلق بمؤشر كتلة الجسم، كانت ذات دلالة احصائية ( $p < 0.001$ ). بمعنى آخر، مؤشر كتلة الجسم أعلى في الحالات الايجابية لمتلازمة الأكل الليلي عن الحالات السلبية. وقد تم تأكيد ذلك بالعلاقة الطردية بين درجات استبيان الأكل الليلي و مؤشر كتلة الجسم حيث ازدادت الدرجات مع تزايد مؤشر كتلة الجسم.

• ارتبطت متلازمة تناول الطعام ليلا بشكل كبير مع الاكتئاب الغير نمطي في المجموعة أ، و هذا ما أكدته درجات استبيان الأكل الليلي التي كانت ذات دلالة احصائية في المقارنة بين الاكتئاب الغير نمطي مع الأنواع الأخرى. في حين المجموعة ب ، ارتبطت حالات متلازمة الأكل الليلي بالاكتئاب الغير نمطي و الاكتئاب الأعظم.

• كان هناك فروق احصائية بين الحالات السلبية و الايجابية لمتلازمة تناول الطعام ليلا في المجموعة ب فيما يتعلق بنوع مضادات الاكتئاب المستخدمة. و قد وجد هذا مع ميرتازابين، ترازودون و كلوميبرامين.

**وبناء على نتائج الدراسة الحالية توصلت الدراسة إلى التوصيات التالية:**

• بما أن متلازمة الأكل الليلي شائعة بين مرضى الاكتئاب ، لذلك يجب استبعادها من قبل الأطباء في مرضى الاكتئاب خاصة بين البالغين و الذين يعانون من الاكتئاب الغير نمطي.

• بما أن هناك علاقة قوية بين متلازمة تناول الطعام ليلا و مؤشر كتلة الجسم، لذا يجب ان لا يهمل الأطباء النفسيين الدور الرئيسي لعلاج السمنة جنباً الى جنب مع علاج الاكتئاب و متلازمة تناول الطعام ليلا.

• هناك تأكيد على الحاجة لتحديد الأفراد ذات الاكتئاب الغير نمطي، حيث وجد أن هذا النوع من الاكتئاب كان مؤشراً قوياً للسمنة و بالتالي أخذ التدابير العلاجية للتقليل من عواقب زيادة الشهية خلال نوبات الاكتئاب.

• هناك الحاجة لدراسات مستقبلية لدراسة تغيرات الوزن و العلاقة بين متلازمة الأكل الليلي و مؤشر كتلة الجسم كعوامل خطر محتملة لبعضهم البعض.

• المزيد من الأبحاث ضرورية لتأكيد فهمنا لمتلازمة تناول الطعام ليلا التي تحدث بالتزامن مع الاكتئاب من أجل اكتشاف أسباب و آثار متلازمة تناول الطعام ليلا بين مرضى الاكتئاب.

• هناك الحاجة لمزيد من الأبحاث لدراسة العلاقة بين متلازمة تناول الطعام ليلا و العقاقير المضادة للاكتئاب لتقييم آثارها و كيفية العلاج الأفضل لمتلازمة تناول الطعام ليلا أو منعه.

• في المرضى الذين يعانون من متلازمة تناول الطعام ليلا و الاكتئاب معاً، يفضل أن ننظر الى تأثير بعض أنواع مثبتات امتصاص السيروتونين الانتقائية في العلاج.

• تجنب قدر الامكان مضادات اكتئاب معينة مثل ميرتازابين، كلوميبرامين و ترازودون في علاج متلازمة تناول الطعام ليلا بين مرضى الاكتئاب حيث قد يكون لديهم بعض المخاطر في احداث متلازمة تناول الطعام ليلا الأمر الذي سيجعل العلاج صعباً.

• تثقيف المرضى وأفراد الأسرة على الآثار الجانبية لمضادات الاكتئاب ( تغيرات الشهية ، زيادة الوزن، خطر الحساسية المفرطة تجاه الجلوكوز و تشوهات الدهون).

• ضبط مضادات الاكتئاب للمرضى إن أمكن وفقاً للبيانات الأيضية للمرضى.

# الملخص العربي

لجنة الإشراف

.....

أ.د./ طارق كمال ملوخية  
أستاذ الطب النفسي  
كلية الطب  
جامعة الإسكندرية

.....

د./ أسامة أبو المجد الخولي  
أستاذ مساعد الطب النفسي  
كلية الطب  
جامعة الإسكندرية

دراسة معدل انتشار متلازمة تناول الطعام ليلا في مرضى الاكتئاب و علاقته بالعقاقير  
المضادة للاكتئاب

مقدمة من

مي عصام حامد رزق  
بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٦

للحصول على درجة

الماجستير

في

أمراض المخ والأعصاب والطب النفسي

موافقون

لجنة المناقشة والحكم على الرسالة

.....

أ.د/ نهلة السيد ناجي  
أستاذ الطب النفسي  
كلية الطب  
جامعة عين شمس

.....

أ.د/ طارق كمال ملوخية  
أستاذ الطب النفسي  
كلية الطب  
جامعة الإسكندرية

.....

د/ سهبا عبد اللطيف الغباشي  
أستاذ مساعد الطب النفسي  
كلية الطب  
جامعة الاسكندرية



# دراسة معدل انتشار متلازمة تناول الطعام ليلا في مرضى الاكتئاب و علاقته بالعقاقير المضادة للاكتئاب

رسالة

مقدمة الي كلية الطب - جامعة الإسكندرية  
إستيفاء للدراسات المقررة للحصول على درجة

الماجستير

في

أمراض المخ والأعصاب والطب النفسي

مقدمة من

مي عصام حامد رزق

بكالوريوس في الطب والجراحة- كلية الطب- جامعة الإسكندرية (٢٠٠٦)

كلية الطب

جامعة الإسكندرية

٢٠١٥