



Conclusion & Recommendations



6. CONCLUSION AND RECOMINDATIONS

From the present study, it was concluded that:

- The technique used in this study succeeded in conserved water consumption, wastewater discharged and reduced the production costs.
- The technique used in this study succeeded in separating of the nutritional material from the wastewater.
- The addition of new dried separated material from the discharge to rats diet did not cause or simple adverse changes in physiological or biochemical parameters tested and showed improvement in growth rates.

Recommendations:

It was recommended that:

- Decreasing the amount of the raw water which is used in rice starch plant as followed in the study would help to preserve the water source, decrease the wastewater as well and decrease the production cost.
- Using the organic extracts which containing a high proportion of protein as new by-product in rice starch plant as additives for animal feed will raise the company's revenue.
- Further studies have to be taken on the techniques used for precipitation to reduce the time and improve the quality of water for reuse more safety and regaining the discharged food stuff to be reused in animal feeding in a way to minimize its desirable defect caused by the precipitating agents.



References



REFERENCES

1. FAO WATER. (2008). Water at a Glance: The relationship between water, agriculture, food security and poverty, Water Development and Management Unit, Food and Agriculture Organization of the United Nations, <http://www.fao.org/nr/water/docs/waterataglance.pdf> .
2. Abdel-Shafy, H. I. and Aly, R. O. (2002). Water issue in Egypt: resources, pollution and protection endeavors. Review article. Central European Journal of Occupational and Environmental Medicine, 8(1): 3-21.
3. Abdin, A. E. and Gaafar, I. (2009). Rational water use in Egypt, Technological perspectives for rational use of water resources in the Mediterranean region. Bari: CIHEAM: 88:11-27.
4. Abu-Zeid, M. (1991). Water resources assessment for Egypt. Ministry of water resources and irrigation. Egypt.
5. Dakwala, M.; Mohanty, B. and Bhargava, R. (2011). Waste water minimization of starch industry using water pinch technology. International workshop advances in cleaner production. San Paulo – Brazil.1-12.
6. UNESCO. (2006). Water a shared responsibility. The United Nations World Water Development Report, 2(8): 277-303.
7. Mori, H. (2003). Technology transfer manual of industrial wastewater treatment. Overseas environmental cooperation center, Japan, 3(1): 107-120.
8. El-Kady, A. A. (2010). Biotechnological Studies on Utilization of Olive Oil Wastes. M.Sc. Thesis, Department of Biotechnology, Institute of Graduate studies and Research, University of Alexandria, Egypt.
9. Kalvani, N.; Sharma, V. and Singh, L. (2012). Various techniques for the modification of starch and the applications of its derivatives. International Research Journal of Pharmacy, 3(5):25-29.
10. Seoul, S. L.; Kyonggi-do, J. L.; Seoul, D. S. and Seoul, H. L. (1999). Comparison of protein extraction solutions for rice starch isolation and effects of residual protein content on starch pasting properties. Starch/Stärke. 51: 120-125.
11. El-Alem, A. H. (1997). An Economic study on the use of agricultural and food processing residues as animal feed. M.Sc. Thesis, Department of Environmental Studies, Institute of Graduate Studies and Research, Alexandria University, Egypt.
12. Oladunjoye, I. O.; Ojebiyi, O, O. and Amao, O. A. (2010). Effect of feeding processed cassava (*Manihot esculenta* Crantz) peel meal based diet on the performance characteristics, egg quality and blood profile of laying chicken. Agricultura Tropica et Subtropica (Czech Republic). 43 (2): 119-126.
13. Moussa, H. M.; Barghash, R. M. and Hassan, A. A. (2011). The Current Situation of the Animal Feed Gap in Egypt. Journal of Basic and Applied Scientific Research. Agricultural Economics Department of National Research Centre, Cairo, Egypt
14. Egyptian Ministry of Agriculture and Land Reclamation. (2010). Statistics of livestock. Animal wealth development sector, Economic Affairs Sector. 2-49.

15. Winpenny, J.; Heinz, I.; Koo-Oshima, S.; Salgot, M.; Collado, J.; Hernandez, F. and Torricelli, R. (2010). The wealth of waste: The economics of wastewater use in agriculture. FAO Water Reports (FAO), / FAO, Rome (Italy). Land and Water Div. 35:1-129.
16. Ernest, H. and Ronald, A. S. (2006). Water rates: An industrial user's view. Journal of Environmental Quality, (78): 33-42.
17. United Nations. (2003). Water for people, water for life - UN World Water Development Report, 1(1):1-22.
18. Gad, A. and Ramadan, A. R. (2009). Water rationalization in Egypt the perspective of the virtual water concept. In: Technological perspectives for rational use of water resources in the Mediterranean region. 88: 301-310.
19. Abou-Rayan, M. A. and Djebedjian, B. (2000). Egypt's water demand, supply and management policies, presented at the workshop and training course, Mediterranean Cooperation for Water Desalination Policies in Perspective of Sustainable Development, Paris.
20. Agana, B. A. (2013). Factors required to enhance water recycling at two major Australian industrial manufacturing sites. Ph.D. thesis, Victoria University of Australia.
21. Meybeck, M.; Chapman, D. and Helmer, R. (1989). Global assessment of fresh water quality: a first assessment, Basil black well, Oxford, 307.
22. Madhukar, R. and Srikantaswamy, S. (2013). Impact of industrial effluents on the water quantity of varishabavathi river and byramangala lake in bidadi industrial area, Karnataka, India. International Journal of Geology, Earth & Environmental Sciences. 3 (2): 132-141
23. UNEP. (1993). Re-use of water. An overview. WHO/ FAO/ UNCHS/ UNEP. Workshop on health, agriculture and environmental aspects of the use of wastewater, Mexico City., Mexico.
24. Anonymous. (2005). Technical Manual: Reclaimed water for beneficial re-use. New Jersey Division of water quality, USA.
25. Kaushik, G., Satya, S. and Naik, S.N. (2009). Food processing a tool to pesticide residue dissipation-A review. Food Research International, 42 (1): 26-40.
26. European Commission (EC). (2006).Integrated pollution prevention and control: Reference Document on Best Available Techniques in the Food, Drink and Milk Industries.
27. Winger, R. and Wall, G. (2006). Food product innovation. A background paper. FAO. 2(5): 21-24.
28. Selim, T. H. (2009). The Egyptian Food Processing Industry: Formalization versus Informalization within the Nation's Food Security Policy. Alfred P. Sloan Industry Studies Association General Industry Studies Annual Conference, Chicago.
29. Okazaki, W. K. (2006). Identification and Assessment of Food Waste Generators in Hawaii. MSc. thesis in Urban and Regional Planning. University of HAWAII. 2: 14-21.
30. Corcoran, E.; Nellemann, C.; Baker, E.; Bos, R.; Osborn, D. and Savelli, H. (2010). Sick Water? The central role of wastewater management in sustainable development. A Rapid Response Assessment. United Nations Environment Programme, UN-HABITAT, GRID-Arendal. www.grida.no .

31. European Parliament, C. (2008). Directive 2008/98/EC of the European Parliament and of the Council of 19 November 2008 on waste and repealing certain Directives.
32. Riemer, J and Kristoffersen, M. (1999). Information on waste management practices. A proposed electronic framework. European Environmental Agency, Copenhagen, Denmark,24(2): 9-15
33. Vigneswaran, S. and Sundaravadivel, M. (2004). Recycle and reuse of domestic wastewater., in Encyclopedia of Life Support Systems (EOLSS). Developed under the Auspices of the UNESCO, Eolss Publishers, Oxford UK.
34. Raschid-Sally, L. and Jayakody, P. (2008). Drivers and characteristics of wastewater agriculture in developing countries: Results from a global assessment, Colombo, Sri Lanka. IWMI Research Report 127, International Water Management Institute, Colombo. 35.
35. Janesiripanich, A. (1995). Waste auditing in rice cracker factory. M.Sc. Thesis, Asian Institute of Technology. Bangkok, Thailand.
36. Orhon, D.; Babuna, F.G. and Insel, G. (2001). Characterization and modeling of denim-processing wastewaters for activated sludge. Journal of Chemical Technology and Biotechnology. 76: 919-931.
37. Jin, B.; Yan, X. Q.; Yu, Q. and Van-Leeuwen H. J. (2002). A comprehensive pilot plant system for fungal biomass protein production and wastewater reclamation, Advances in Environmental Research. 6: 179–189.
38. Mitchell, J. R. (2009). Rice starches: production and properties. In: BeMiller J, Whistler R, editors. Starch chemistry and technology. New York: Academic Press. 569–579.
39. Lawal, O. S.; Lapasin, R.; Bellich, B.; Olayiwola, T. O.; Cesaro, A.; Yoshimura, M. and Nishinari, K. (2011). Rheology and functional properties of starches isolated from five improved rice varieties from West Africa. Food Hydrocolloid 25: 1785–92.
40. Manek, R. V.; Kunle, O. O.; Emeje, M. O.; Builders, P.; Rao, G. V. R.; Lopez, G. P. and Kolling, W. M. (2005). Physical, thermal and sorption profile of starch obtained from *Tacca leontopetaloides*. *Starch - Stärke*, 57 (2): 55-61.
41. Shen, L.; Haufe, J. and Patel, M. K. (2009). Product overview and market projection of emerging bio-based plastics. Final report, Group Science, Technology and Society (STS) Copernicus Institute for Sustainable Development and Innovation. Utrecht University.
42. Debnath, S.; Gayathri, V. G. and Niranjana-Babu, M. (2013). Chemically modified starches and their applications in pharmacy. International Journal of Research in Pharmaceutical and Nano Sciences. 2(3): 332 -344.
43. Sajilata, M. G.; Singhal, R. S. and Kulkarni, P. R. (2006). 'Resistant starch—A Review'. Comprehensive reviews in food science and food safety, 5 (1): 1-17.
44. Tolvanen, P. A. (2013). Development of an environmentally friendly method of starch oxidation by hydrogen peroxide and a complex water-soluble iron catalyst. Doctoral Thesis, Industrial Chemistry and Reaction Engineering, Process Chemistry Centre, Department of Chemical Engineering, Åbo Akademi University.

45. Duval, L.; Menet, L. and Charbonneau, C. (2010). Evaluation of common agricultural policy measures applied to the starch sector Final Report - Agro-Synergie.
46. Fuglie, K. O.; Oates, C. G. and Xie, J. (2006). Root crops, Starch and agro-industrialization in Asia International Potato Center. CIP Social sciences working paper, 3: 1-19.
47. Agrosynergie. (2010). Evaluation of common agricultural policy measures applied to the starch sector. Final report. European commission, DG Agriculture and rural development, 2:17-79.
48. Rosell, C. M. and Marco, C. (2008). Rice in Arendt, E. K and DalBello, F editor. Gluten-Free Cereal Products and Beverages Chapter 4: 81-90.
49. FAO. (2013). Rice market monitor. Trade and markets division. Food and Agriculture Organization of the United Nations.. Volume XVI - Issue No. 1.
50. Antunes, T. M. S. (2002). Permeability of rice cystatin Across Caco-2 Cells 3-7.
51. Cruz, N. D., and Khush, G. S. (2000). Rice grain quality evaluation procedures. First ed. *In 'Aromatic Rices'*. 15-28.
52. Calpe, C. (2006). Rice international commodity profile. FAO. Markets and trade division. 2-22
53. Lai, H. M. (2001). Effect of rice properties and emulsifiers on the quality of rice pasta. *Journal of Science Food Agriculture*. 82: 203-216.
54. Onwulata, C. I.; Smith, P.W.; Konstance, R. P. and Holsinger, V. H. (2001). Incorporation of whey products in extruded corn, potato or rice snacks. *Food Research International Journal*. 34: 679-687.
55. GuJral, H. S.; Haros, M. and Cristina, M. R. (2003). Starch hydrolyzing enzymes for retarding the staling of rice bread. *Cereal Chemistry*. 80(6): 750-754.
56. Lovis, I. J. (2003). Alternatives to wheat flour in baked goods. *Cereal Foods World*, 48 (2): 61-63.
57. GuJral, H. S. and Cristina, M. R. (2004). Functionality of rice flour modified with a microbial trans glutaminase. *Journal of Cereal Science*. 39: 225-230.
58. Sivaramakrishnan, H. P.; Senga, B. and chattopadhyay, p. k. (2004). Rheological properties of rice dough for making rice bread. *Journal of Food Engineering*. 62: 37-45.
59. Takahashi, T.; Miura, M.; Ohisa, N.; Mari, K. and Kobayashi, S. (2005). Heat treatments of milled rice and properties of the flour. *Cereal Chemistry*. 82(2): 228-232.
60. Cameron, D. and Wang, Y. J. (2005). A better understanding of factors that affect the hardness and thickness of long-grain rice. *Cereal Chemistry*. 82(2): 113-119.
61. Heinemann, R. J. B.; Fagundes, P. L.; Pinto, E. A.; Pentead, M. V. C. and Marquez, U. M. L.(2005). Comparative study of nutrient composition of commercial brown parboiled and milled rice from Brazil. *Journal of Food composition Analysis*, 18: 287-296.

62. Anjum, F.M.; Pasha, I.; Bugti, M.A. and Butt, M.S. (2007). Mineral composition of different rice varieties and their milling fractions. *Pakistan Journal of Agricultural Sciences*, 44(2):332-336.
63. Lozaridou, A.; Duta, D.; Papageoriou, M.; Belc, N. and Biliaderis, C. G. (2007). Effects of hydrocolloids on dough rheology and bread quality parameters in gluten-free formulation. *Journal of Food Engineering*. 79: 1033-1047.
64. Yadav, R. B.; Khatkar, B. S. and Yadav, B. S. (2007). Morphological, physical and cooking properties of some Indian rice (*Oryza Sativa L.*) cultivars. *Journal of Agriculture Technology*, 3(2): 203-210.
65. Wanyo, P.; Chomnawang, C. and Siriamornpum, S. (2009). Substitution of wheat flour with rice flour and rice bran in flake products: Effects on chemical, physical and antioxidant properties. *World Applied Sciences Journal*. 7(10): 49-56.
66. Hodzic, Z.; Memisevic, A.; Srabovic, M.; Saletovic, M. and Poljakovic, M. (2009). The influence of total phenols content on antioxidant capacity in the whole grain extracts. *European Journal of Scientific Research*. 28(3): 471-477.
67. Umadevi, M.; Pushpa, R.; Sampathkumar, K. P. and Bhowmik, D. (2012). Rice-traditional medicinal plant in india. *Journal of Pharmacognosy and Phytochemistry*. 1(1):6-12.
68. Juliano, B. O. (1993). Rice in human nutrition. . Food and Nutrition Series 26. FAO, Rome. Italy. International Rice Research Institute, Los Baños, Laguna, Philippines.
69. Puwastien, P.; Raroengwichit, M.; Sungpuag, P. and Judprasong, K. (1999). Thai food composition tables, first edition. Institute of Nutrition, Mahidol University, Thailand.
70. Juliano, B. O. and Villareal, C. P. (1993). Grain quality evaluation of world rice's. International Rice Research Institute, Manila, Philippines.
71. Lumdubwong, N and Seib, P. A. (2000) Rice starch isolation by alkaline protease digestion of wet-milled rice flour. Kansas State University, Department of Grain Science and Industry, Manhattan, KS 66506, U.S.A.
72. Lau, J. Y. (2005). Protein structure database for structural genomics group. M.Sc. Thesis, Biomedical Engineering. University of New Jersey.
73. Young, V. R (2001). Protein and amino acids. In: Present Knowledge in Nutrition. 8th Edition. Bowman BA and Russel RM (eds). International Life Sciences Institute, Washington DC. 5: 43-58.
74. Voet, D. and Voet, J. G. (2004). Biochemistry: biomolecules, mechanisms of enzyme action, and metabolism, 3rd edn, Vol. 1. John Wiley & Sons, Inc., Hoboken.
75. Rondina, M. V.; Beringer, M. and Wintermeyer, W. (2007). 'How ribosomes make peptide bonds,' *Trends in biochemical Sciences*, 32: 20-26.
76. Shewry, P. R and Halford, N. G. (2001). Cereal seed storage proteins: structures, properties and role in grain utilization. *Journal of Experimental Botany*, 53: 370.
77. Landers, P. S. and Hamaker, B. R. (1994). Antigenic properties of albumin, globulin, and protein concentrate fractions from rice bran. *Cereal Chemistry*. 71: 409-411.

78. Marshall, W. E. and Wordsworth, J. I. (1994). Rice science and technology. New York: Marcel Dekker, Inc. 237-259.
79. Janick, J. (2002). Rice: the natural history of rice [online] Available at http://www.hort.purdue.edu/newcrop/tropical/lecture_20/rice.html. (Verified 24 Sep. 2006).
80. Padhye, Y and Salunkhe, D. K (1979) Extraction and characterization of rice proteins. Cereal Chemistry. 56 -389.
81. Shih, F. F and Daigle, K. W (2000). Preparation and characterization of rice protein isolates. Journal of the American Oil Chemists Society, 77 (8): 885-889.
82. Wu, L. S.; Wang, L. D.; Chen, P.W.; Chen, L. J and Tzen, J. T. (1998). Genomic cloning of 18 kDa oleosin and detection of triacylglycerols and oleosin isoforms in maturing rice and postgerminative seedlings. Journal of Biochemistry. 123: 386–391.
83. Zhout, Z.; Blanchard, C.; Helliwell, S. and Robardsa, K. (2002) Fatty acid composition of three rice varieties following storage. Journal of Cereal Science. 1-9.
84. Organisation for Economic Co-operation and Development (2004). Consensus document on compositional considerations for new Varieties of Rice (*Oryza sativa*): Key Food and Feed Nutrients and Anti-nutrients.
85. Juliano, B. O. and Bechtel, D. B. (1985). The rice grain and its gross composition. I n: 2nd ed. Rice Chemistry and Technology, American Association of Cereal Chemists, Inc. St. Paul, Amsterdam. 17–57.
86. El-Hissewy, A. A.; Laila, F. R. and Hanaa, A. D. (2002). Effect of degree of milling on the chemical composition and nutritional value of the milled rice. Egyptian Journal of Agricultural Research. 80(1): 341-353.
87. Adam, K. K. and Liu, R. H. (2002). Antioxidant of activity of grain. Journal of Agricultural and Food Chemistry. 50: 6182-6187.
88. Goffman, F. D. and Bergman, C. J. (2004). Rice kernel phenolic content and its relationship with antiradical efficiency, Journal of the Science of Food and Agriculture. 84: 1235-1240.
89. Tian, S. U.; Nakamura, K. and Kayahara, H. (2004). Analysis of phenolic compound in white rice. Brown rice and germinated brown rice. Journal of Agricultural and Food Chemistry. 52: 4808-4813.
90. Zhou, Z.; Robards, K.; Helliwell, S. and Blanchard, C. (2004). The distribution of phenolic acids in rice. Food Chemistry. 87: 401-406.
91. Dykes, L. and Rooney, L. W. (2007). Phenolic compounds in cereal grain and their health benefits. Cereal Food World. 52(3): 105-111.
92. Liu, R. H. (2007). Whole grain phytochemicals and health. Journal of Cereal Science. 46: 207-219.
93. Liang, J.; Han, B. Z.; Nout, M. J. R. and Hamer, R. J. (2009). Effect of soaking and phytase treatment on phytic acid. Calcium. Iron and zinc in rice fractions. Food Chemistry. 115: 789-794.

94. Shen, Y.; Jin, L.; Xiao, P.; Lu, Y. and Bao, J. (2009). Total phenolics, flavonoids, antioxidant capacity in rice grain and their relation to grain color, size and weight. *Journal of Cereal Science*. 49: 106-111.
95. USDA (1994). Chapter 5. Inspection of milled rice in Rice Inspection Handbook. USDA Fedral. Grain Inspection Service. Washington D. C., U.S.
96. Schramm, R. C. (2010). Value-added processing of rice and rice by-products. Ph.D. thesis, Department of Engineering Science. Louisiana State University, 2 - 67.
97. Food and Agriculture Organization of the United Nations (FAO). (2004). World Rice Research Conference. Tsukuba, 5-7 November 2004.
98. Ministry of Supply and Internal Trade, Egypt (1990) Egyptian standards for broken rice. 330.
99. Doma, M. B. (1970). Chemical and technological studies on rice starch industry in U.A.R. and the utilization of the Wastes. M.Sc. thesis, Faculty of agriculture, Alexandria University. 40-91.
100. Hertrampf, J. W. and Piedad-Pascual, F. (2000). Handbook on Ingredients for Aquaculture Feeds. 351-355.
101. Dahab, D. B. O. (2006). Utilization of different cereal flour mixes in the preparation of some baking products. M.Sc. Thesis, Faculty of Agriculture. Minufia Univirsity. 160.
102. Brault, J. B. (1991). Water treatment handbook. Vol 1 dan 2, 6th Edition Degremont, Lavoisier Publishing, Codex, France.
103. Whistler, R. L. and Paschall, E. F. (1967). Starch chemistry and technology. 2(3): 65-86.
104. Puchongkavarin, H.; Varavinit, S. and Bergthaller, W. (2005). Comparative study of pilot scale Rice starch production by an alkaline and an enzymatic process. *Starch/ Stärke*, 57:134–144.
105. Cornell, H. J. and Hoveling, A. W. (1998). Wheat: chemistry and utilization. *Plant Growth Regulation Journal*. 26(2): 139.
106. Jaczynski, J. (2008). Protein and lipid recovery from food processing by-products using isoelectric solubilization/precipitation. West Virginia University. Food Chemistry Research Developments.
107. Bordwell, F. G. (1988). Equilibrium acidities in dimethyl sulfoxide solution. *Accounts of Chemical Research*. 21:456-463.
108. Heinau, V. and Kirste, B. (1994). Amino acids, institute of chemistry, Department of Biology, Chemistry, Pharmacy, FU Berlin, http://WWW.chemie.fuberlin.de/chemistry/bio/amino-acids_en.html, 15 July.
109. Shyur, L.; Zia, K.. and Chen, C. (1988). Purification and some properties of storage proteins in japonica rice. *Botanical Bulletin of Academia Sinica*. 29.
110. Ju, Z. Y.; Hettiarachchy, N. S. and Rath, N. (2006). Extraction, denaturation and hydrophobic Properties of Rice Flour Proteins. *Journal of Food Science*. 66(2): 229-232 .

111. Swanson, B. G. (1990). Pea and lentil protein extraction and functionality. *Journal of the American Oil Chemists' Society* 67: 276-280.
112. Owusu-Ansah, Y. J. and McCurdy, S. M. (1991). Pea proteins: a review of chemistry, technology of production, and utilization. *Food Reviews International* 7: 103-134.
113. Ford, M. and Fleming, R. (2002). Mechanical solid-liquid separation of livestock manure Literature Review Ridgetown College - University of Guelph. 2–7.
114. Omelia, C. (1998). "Coagulation and sedimentation in lakes, reservoirs and water treatment plants". *Water Science and Technology* 37 (2): 129.
115. Perry, R. H. and Green, D. W. (1997). *Perry's Chemical Engineers' Handbook*, 7th ed., McGraw-Hill, New York. Chapters 18 and 22.
116. Schweitzer, P. A. (1997). *Handbook of separation techniques for chemical engineers*, 3rd ed., McGraw-Hill, New York. (4): 140-156.
117. Sheffield, R.; Barker, J. and Rashash, D. (2000). Solids separation of animal manure. North Carolina State University. Available online at: [res2.agr.ca/initiatives/manurenet/en/man_tech_int.html#Solid/Liquid Separation](http://res2.agr.ca/initiatives/manurenet/en/man_tech_int.html#Solid/Liquid%20Separation).
118. Metcalf, L. and Eddy, H. P. (1991). *Wastewater engineering: treatment, disposal and reuse*, 3d ed., McGraw-Hill, New York.
119. Fleming, R. (1986). Solids-Liquid separation of manure. Ontario Ministry of Agriculture and Food Factsheet. 86-032.
120. Glerum, J. C.; Klomp, G. and Poelma, H. R. (1971). "The separation of solid and liquid parts of pig slurry. In Proc. 1st International Symposium on Livestock Wastes, 345-347. Columbus, Ohio, 19-22 April: American Society of Agricultural Engineers
121. Sneath R.W., Shaw M. and Williams A.G. 1988. 'Centrifugation for separating piggery slurry 1. The performance of a decanting centrifuge'. *Journal of Agricultural Engineering Research*. 39: 181-190.
122. Shutt, J. W.; White, R. K.; Taiganides, E. P. and Mote, C. R. (1975). Evaluation of solids separation devices. In Proc. 3rd International Symposium on Livestock Wastes, 463-467. Urbana Champaign, Illinois, 21-24 April: American Society of Agricultural Engineers.
123. FAO. (1983). *World food security: a Reappraisal of the Concepts and Approaches*. Director General's Report. Rome.
124. Tolera, A. (2008). *Feed resources and feeding management: A manual for feedlot operators and development workers*. Ethiopia Sanitary & Phytosanitary Standards and Livestock & Meat Marketing Program (SPS-LMM) Texas Agricultural Experiment Station (TAES)/Texas A&M University.
125. Ensminger, M. E.; Oldfield, J. E. and Heinemann, W.W., (1990). *Feeds and nutrition*. 2nd ed. Ensminger Publishing Company, California, U.S.A.
126. Aregheore, E. M. and Chimwano, A. M., (1991). Agro-industrial by-products and crop residues in Zambia: Availability, utilisation and potential value in ruminant nutrition. In: *The Complementarity of Feed Resources for Animal Production in Africa*. Proceedings of the Joint Feed Resources Networks Workshop. Eds. Stares, J.E.S., Said, A.N. & Kategile, J.A.,

Gaborone (Botswana) African Feed Resources Network. International Livestock Centre for Africa (ILCA), Addis Ababa (Ethiopia), 223-238.

127. Fadel, J. G. (1999). Quantitative analyses of selected plant by-product feedstuffs, a global perspective. *Animal Feed Science and Technology*. 79: 255–268.

128. Boucqué, C. H. V. and Fiems, L.O., (1988). Vegetable by-products of Agro-Industrial origin. *Livestock Production Science*. 19: 97-135.

129. Wilkinson, J. M. (1988). The feed value of by-products and wastes. In: *World Animal Science, B4 Feed Science, 4. Disciplinary approach*. Ed. Ørskov, E.R. Elsevier Science, Amsterdam, the Netherlands. 313-327.

130. Fouda, S. M. E. (2008). Studies on the food bagasse previous treatment with some transactions chemical and biological. Ph.D. Thesis, Department of Animal Production. Azhar, university.

131. Ministry of Agriculture and Land Reclamation. (2011). Statistics. Economic Affairs Sector, Egypt.

132. Egyptian Starch, Yeast and Detergent Company, Financial Sector. (2008: 2012).

133. APHA, AWWA, WEF, “Standard Methods for the Examination of Water and Wastewater,” 20th Edition, Washington DC, (1998).

134. A.O.A.C. (2000). Official Methods of Analysis of the Association of Official Analytical Chemists, 17th edn (edited by W. Horwitz). Washington, DC. 32. Cereal Foods.

135. Smith, F., Gilles, M. A.; Hamilton, J. K. and Godess, P. A. (1956). Colorimetric method for determination of sugar and related substances. *Anal. Hem.* 26: 350 – 358.

136. Mohr C. F. (1856). Neue massanalytische Bestimmung des Chlors in Verbindungen, *Ann. Chem. Pharm.*, 97: 335-338.

137. Lee, S. Y.; Morr, C. V. and Ha, E. Y. W. (1992). Structural and functional properties of caseinate and whey protein isolate as affected by temperature and pH. *Journal of Food Science*, 57: 1210–1214.

138. Taga, M. S., Miller, E. E. and Pratt, D. E. (1984). Chia seeds as a source of natural lipids antioxidants. *Journal of American Oil Chemistry Society*. 61: 928-931.

139. International Agency for Research on Cancer (2002). Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. In *IARC monographs on the evaluation of carcinogenic risks to humans*, 82: 171–175.

140. European Community (2003). 2003/100/EC. Commission Directive 2003/100/EC of 31 October 2003 amending Annex I to Directive 2002/32/EC of the European Parliament and of the Council on undesirable substances in animal feed (text with EEA relevance). *Official Journal of the European Union L 285*: 33–37.

141. A.O.A.C. (1995). Official Methods of Analysis of the Association of Official Analytical Chemists, 16th ed. Gaithersburg, MD, Method 970.44, sec. 49.2.02.

142. A.O.A.C. (1998). Official methods of analysis. Association of Official Analytical Chemists, 16th edition. Arlington, Virginia.

143. "MAFF" Ministry of Agriculture, Fisheries and Food Department of Agriculture and Fisheries for Scotland, Department of Agriculture for Northern Ireland. (1975). Energy Allowances and Feeding Systems for ruminants Technical Bulletin No 33 Her Majesty's Stationery Office London.
144. Marai, I. F. M.; El-Darawany, A. A.; Fadiel, A. and Abdel-Hafez M. A. M. (2007). Physiological traits as affected by heat stress in sheep—A review. *Small Ruminant Research*; 71: 1–12.
145. N.R.C., (1995). National Requirements of Laboratory Animals. National Research Council Fourth revised Edition. National Academy Press Washington, D.C. 11-79.
146. Greer, J. P. Foerster, J. and Lukens, J. N. (2003). *Wintrobe's Clinical Hematology*, 11th Ed; Publisher: Lippincott Williams & Wilkins Publishers, USA.
147. Emerson, W.W. (1977). Physical properties and structure. In *Soil factors and crop production in semi-arid environment*, ed. J. S. Russel and E. L. Greacen, 78–104. St Lucia, Australia: University of Queensland Press.
148. Armstrong, W. D. and Carr, C.W. (1964). *Physiological chemistry laboratory Directions* 3rd ed. Burges Publishing Co., Minneapolis, Minnesota. 75.
149. Dumas, B. T., Watson, W. A. and Biggs, H. G. (1971). Albumin standards and the measurement of serum albumin with bromocresol green. *Clinica. Chemica. Acta.*; 31: 87-96.
150. Frings, C. S., Fendley, T. W., Dunn, R. T. and Queen, C.A. (1972). Improved determination of total serum lipids by the sulfo-phospho-vanilin reaction. *Clini. Chem.*; 18: 673-674.
151. Watson, D. (1960). A simple method for the determination of serum cholesterol. *Clinica. Chemica. Acta.*; 5: 637- 638.
152. Richmond, W. (1973). Quantitative determination of cholesterol in serum or plasma by enzymatic method. *Clin. Chem.*; 19: 1350.
153. Assmann, G., Jabs, H.U., Kohnert, U., Nolte, W. and Schriewer, H. (1984). Low density lipoprotein – cholesterol method. *Clinica. Chemica. Acta.*; 140: 77–83.
154. Bucolo, G. and David, H. (1973). Quantitative determination of serum triglycerides by the use of enzymes. *Clinica. Chemica. Acta.*; 19: 476-482.
155. Reitman, S. and Frankel, S. (1957). A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*. 28: 56-63.
156. Bauer, J. D. (1982). *Clinical Laboratory Methods*. C. V. Mosby Co., USA, 9th Ed, 580-581. *Biological Trace Element Research Journal*. 34: 225.
157. Andersch, M.A. and Szcypinski, A. J. (1947). A colorimetric method for determination of acid phosphatase from serum. *American Journal of Clinical Pathology*. 17: 571.
158. Stroev, E.A. and Makarova, V.G. (1989). *Laboratory Manual in Biochemistry*, Mir publishers, Moscow, Russia.
159. Patton, C. J. and Crouch, S.R. (1977). Colorimetric determination of urea. *Clinica. Chemica. Acta.*; 49: 464-469.

160. Larsen, K. (1972). Colorimetric method for creatinine determination. *Clin. Chemica. Acta.*; 41: 209-217.
161. Jendrassik, L. and Grof, P. (1938). Vereinfachte photometrische Methoden zur Bestimmung des Bilirubins. *Biochemische Zeitschrift* 297: 81-89.
162. Barham, D. and Trinder, P. (1972). Enzymatic determination of uric acid. *Analyst*; 97: 142-145.
163. Tappel, L and Zalkin, H. (1959). Inhibition of lipid peroxidation in mitochondria by vitamin E. *Arch. Biochem. Biophys.*; 80: 333-336.
164. Statistical Product and Service Solutions "SPSS" (2006). Manual Guide of SPSS version 16.0. SPSS Inc. Chicago, USA.
165. Nasr, M.; Tawfik, A.; Ookawara, S. and Suzuki, M. (2013). Anaerobic biohydrogen production. *BioResources* : 8 (4): 4951-4968.
166. Kumar, K. and Gidde, M. R. (2014). Electrocoagulation for Treatment of Parboiled Rice Waste Water. *The International Journal of Science & Technoledge*. ISSN 2321 – 919X. 2 (4).
167. Tawfik, A., Elsayed, H. and Dessouki, H. (2013). Hydrogen and methane production from starch waste water in a mesophilic anaerobic baffled reactor. Presented at the Seventeenth International Water Technology Conference Proceedings.
168. Harris, J. L. (1986) Influence of gel layer rheology on ultrafiltration flux of wheat starch effluent, *Journal of Membrane Science*, 29: 97-109.
169. Jin, B.; van Leeuwen, H. J.; Patel, B.; Doelle, H.W. and Yu, Q. (1999). Production of fungal protein and glucoamylase by *Rhizopus oligosporus* from starch processing wastewater. *Process Biochem* 34: 59 – 65.
170. Pidgeon, E. F. (2008). The Application of crossflow membrane filtration technology to remediate wheat starch processing wastewater for reuse. Griffith School of Engineering, Griffith University.
171. Kroyer, G.T. (1995). Impact of food processing on the environment– An overview. *Lebensm. Wiss. Technol.* 28: 547–552.
172. Yanagi, C., Sato, M. and Takahara, Y. (1994) Treatment of wheat starch waste water by a membrane combined two phase methane fermentation system, *Desalination*, 98: 161-170.
173. Hein, P. G., Oanh, L. T. K., Viet, N. T. and Lettinga, G. (1999) Closed wastewater system in the tapioca industry in Vietnam, *Water, Science and Technology*, 39: 89-96.
174. Oanh, L. T. K., de Jong, K., Mai, H. N. P., and Viet, N. T., (2001). Removing Suspended Solids from Tapioca Processing Wastewater in Upflow Anaerobic Filter (UAF). *International Conference: Industry and Environment in Vietnam, Ho Chi Minh City, Vietnam*.
175. Mai, H. N. P.; Thai, L. N.; Viet, N. T. and Lettinga, G., (2001) Effect of Organic Loading Rate on Treatment Efficiency for Tapioca Processing Wastewater using UASB. *Proceedings of International Conference on Industry and Environment in Viet Nam, 20-21 April 2001, HCMC, Viet Nam*, 224-233.

176. Mironescu, M. (2011). Investigations on wastewaters at potato processing and starch recovery and characterization. *Journal of Agroalimentary Processes and Technologies*, 2011, 17(2): 134-138.
177. Sharif, M. K. (2009). Rice industrial by-products management for oil extraction and value added products. Doctor of Philosophy thesis, Degree of Doctor of Philosophy of Food Technology, National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan, 206.
178. Li, S.; Chou, T. and Shih, C. (2010). Effects of brown rice, rice bran, and polished rice on colon carcinogenesis in rats. *Food Research International*.
179. Mesomya, W.; Sutthivaiyakit, P.; Cuptapun, Y. and Hengsawadi, D. (2012). Effects of organic rice compared with conventional rice on protein efficiency ratio and pesticide residue in rats. *Maejo International Journal of Science and Technology*. 6(03): 470 – 482.
180. Yang, L.; Chen, J.; Lv, J.; Wu, Q.; Xu, T.; Zhang, H.; Liu, Q. and Yang, H. (2012). Rice protein improves adiposity, body weight and reduces lipids level in rats through modification of triglyceride metabolism. *Lipids Health Dis*, 13: 11-24.
181. Ahmed, F.; Platel, K.; Vishwanatha, S.; Puttaraj, S. and Srinivasan, K. (2007). Improved shelf-life of rice bran by domestic heat processing and assessment of its dietary consumption in experimental rats. *Journal of the Science of Food and Agriculture*. 87(1): 1–175.
182. Um, M. Y.; Ahn, J.; Jung, C. H. and Ha, T. Y. (2013). Cholesterol-lowering effect of rice protein by enhancing fecal excretion of lipids in rats. *Preventive Nutrition and Food Science*. 18(3): 210-213.
183. Zakharov, M. N. (2009). A Thesis, A microrheological study of sickle hemoglobin polymerization. Drexel University.5.
184. Dacie, J.V. and S.N. Lewis. (1984). *Practical hematology*. 6th Edition. Edinburg, Churchill Livingstone.
185. Maton, D.; Hopkins, J.; McLaughlin, Ch. W.; Johnson, S.; Warner, M. Q.; LaHart, D.; Wright, J. D. and Deep, V. K (1997). *Human Biology and Health*. Englewood Cliffs, New Jersey, USA: Prentice Hall.
186. Deng, J. L.; Wei, Q., Zhang, M. H., Wang, Y. Z., and Li, Y. Q., (2001). Study of the effect of alcohol on single human red blood cells using near-infrared laser tweezers Raman spectroscopy.
187. Jonson-Delaney, C. A. (1996). *Exotic companion medicine handbook*. Eastside Avian and Exotic animal medical centre, Kirkland, Washington. 12-29.
188. Jędrzejewska-Szczerska, M. and Gnyba M. (2011). Optical investigation of hematocrit level in human blood. *ACTA PHYSICA POLONICA A*, 120.
189. Research Animal Resources, (2007). “Reference Values for Laboratory Animals: Normal Haematological Values,” Research Animal Resources, University of Minnesota. <http://www.ahc.umn.edu/rar/index.html>.

190. Garba, S. H.; Adelaiye, A. B. and Mshelia, L. Y. (2007). Histopathological and Biochemical changes in the rats kidney following exposure to a pyrethroid based mosquito coil. *Journal of Applied Sciences Research*. 3: 1788- 1793.
191. Shallan, M. A.; El-Beltagi, H. S.; Mona, A. M.; Amera, T. M. and Sohir, N. A. (2010). Effect of amylose content and pre-germinated brown rice on serum blood glucose and lipids in experimental animal. *Australian Journal of Basic and Applied Sciences*, 4(2): 114-121.
192. Shakib, M.; Gabrial, S. and Gabrial, G. (2014). Rice bran oil compared to atorvastatin for treatment of dyslipidemia in patients with type 2 diabetes. *Macedonian Journal of Medical Sciences*: 7(1): 95-102.
193. Delanghe, J.; De Slypere, J. P.; De Buyzere, M.; Robbrecht, J.; Wieme, R. and Vermeulen, A. (1989). Normal reference values for creatine, creatinine, and carnitine are lower in vegetarians. *Clin. Chem.*, 35(8): 1802-3.
194. Shih, C.; Chen, S.; Hou, W. and Cheng, H. (2007). A high-resistance-starch rice diet reduces glycosylated hemoglobin levels and improves the antioxidant status in diabetic rats. *Food Research International*. 40: 842–847.
195. Ghalehkandi, J. G.; Ebrahimnezhad, Y. and Nobar, R. S. (2012). Effect of garlic (*Allium sativum*) aqueous extract on serum values of urea, uric-acid and creatinine compared with chromium chloride in male rats. *Annals of Biological Research*, 3 (9): P 4485 - 4490.
196. Radosavljevic, T.; Mladenovic, D.; Vucevic, D.; Petrovic, J.; Hrncic, D.; Djuric, D.; Loncar, H. and Stanojlovic, O. (2008). Effect of acute lindane and alcohol intoxication on serum concentration of enzymes and fatty acids in rats. *Food and Chemical Toxicology*. 46: 1739-1743.
197. Drotman, R. B. and Lawhorn, G.T. (1978). Serum enzymes as indicators of chemically induced liver damage. *Drug and Chemical Toxicology*. 1: 163-171.
198. Ramazzotto, L. J. and Carlin, R. (1978), Effects of DMSO on SGOT during hypothermia in adrenalectomized rats. *Life Sci*, 22: 329-336.
199. Al-Okbi, S. Y.; Mohamed, D. A.; Hamed, T. E. and Esmail, R. S. H. (2014). Rice Bran Oil and Pumpkin Seed Oil Alleviate Oxidative Injury and Fatty Liver in Rats Fed High Fructose Diet. *Polish Journal of Food and Nutrition Sciences*. 64(2): 127–133.
200. Aly El-Deen, M. I. A. (2009). A study on the biochemical indicator changes produced by Y- Hexachlorocyclohexane application and role of fish oil. M.Sc. Thesis, in Environmental Studies, Institute of Graduate Studies and Research, Alexandria University, Egypt. 60-67.
201. Burstone, M. S. (1959). Histochemical demonstration of acid phosphatase activity in osteoclasts. *Journal of Histochemistry and Cytochemistry*. 7: 39-41.
202. Bull, H.; Murray, P. G.; Thomas, D.; Fraser, A. M. and Nelson, P. N. (2002). Acid phosphatases. *Molecular Pathology*. 55 (2): 65–72.
203. Bradbury, S. P.; Mc kim, J. M. and Coats, J. R. (1987). Physiological responses of rainbow trout, *Salmo gairdneri* to acute fenvalerate intoxication, *Pesticide Biochemistry and Physiology*. 27: 258-260.

204. Araújo, E. J.; Sant'Ana Dd Dde, M.; Molinari, S. L. and de Miranda Neto, M. H. (2005). Hematologic and biochemical parameters of rats subjected to hypoprotic and hibercaloric. *Arquivos de Ciências Veterinárias e Zoologia da UNIPAR.*, 8(2): 139-146.
205. Nelson, D. L. and Cox, M. M. (2000). Bioenergetics and metabolism. In "Lehninger, Principles of Biochemistry" 3rd ed., pp.343-363. Worth Publishing: New York,
206. Joo, I. W.; Ryu, J. H. and Oh, H. J. (2010). The influence of sam-chil-geun (Panax Notoginseng) on the serum lipid levels and inflammations of rats with hyperlipidemia induced by poloxamer - 407. *Yonsei Medical Journal.* 51(4): 504-510.
207. Ononogbu, I. C. (1988). Lipid and lipoproteins: chemistry, methodology, metabolism, biochemical and physiological importance. New Africa Publishing Co. Ltd, Owerri , Nigeria.. 77 – 116.
208. Peter, T. (1992). Barley germination: biochemical changes and hormonal control in Barley: genetics, biochemistry molecular biology and biotechnology. Long Ashton, Bristol B5189AF, G74. B32 C.A.B. International UK., 353.
209. Miriam, J. F. V.; Emile, A. M. D.; Lilian, B. M. T.; Johan, M. V. A. and Anton, C. B. (1995). Influence of dietary retrograded starch on the metabolism of natural steroids and bile acids in rats. *British Journal of Nutrition.* 74: 807-820.
210. Izumi, Y.; Ishibashi, G.; Nakanishi, Y. and Kikunaga, S. (2007). Beneficial effect of 3% milled rice on blood glucose level and serum lipid concentration in spontaneously non-insulin dependent diabetic rats. *Journal of Nutritional Science and Vitaminology.* 53: 400-409.
211. Revilla, E., Maria, C. S., Miramontes, E., Bautista, J., Martinez, A. G., Cremades, O., Cert, R. and Parrad, J. (2009). Nutraceutical composition, antioxidant activity and hypercholesterolemia effect of a water-soluble enzymatic extract from rice bran. *Food Research International.* 42: 387-393.
212. Al-Subhi, F. M. M. and Hejazi, M. A. (2013). Brown Rice as a Healthy Food and Lowering Blood Sugar in Rats. *Journal of American Science.* 9(4): 21-28.
213. Olson, R. E. (1998). "Discovery of the lipoproteins, their role in fat transport and their significance as risk factors". *Journal of Nutrition.* 128 (2): 439S-443S.
214. Brunzell, J. D. (2008). "Consensus statement from the American Diabetes Association and the American College of Cardiology Foundation – Lipoprotein management in patients with cardiometabolic risk" *Diabetes Care,* 31: 811-822.
215. Brown, M. S. and Goldstein, J. S. (1992). Koch's postulates for cholesterol. *Cell* 71(2): 187–188.
216. TTL (Taconic Technical Library). (2008). Hematological clinical chemistry values Sprague Dawley rats. Available from: <http://www.taconic.com/>
217. Mesomya, W.; Sutthivaiyakit, P.; Cuptapun, Y. and Hengsawadi, D. (2009). Effects of organic rice compared with conventional rice on serum lipids in rats. *Kasetsart Journal (Natural Science).* 43: 703 – 708.
218. Cromwell, W. C. and Otvos, J. D. (2004). "Low-density lipoprotein particle number and risk for cardiovascular disease" *Curr Atheroscler Rep.*, 6: 381-387.

219. Ihedioha, J. L.; Noel-Uneke, O. A. and Ihedioha, T. E. (2013). Reference values for serum lipid profile of albino rats (*Rattus norvegicus*) of varied ages and sexes. *Comparative Clinical Pathology*, 22 (1): 93 – 99.
220. Purushothama, S.; Raina, P.L. and Hariharan, K. (1995). Effect of long term feeding of rice bran oil upon lipids and lipoproteins in rats. *Mol. Cell. Biochem.* 146 (1): 63-69.
221. Philip, M. D. (2007). HDL cholesterol, very low levels of ldl cholesterol, and Cardiovascular Events. *New England Journal of Medicine*. 357: 301-1310.
222. Gibbons, G. F.; Wiggins, D.; Brown, A. M. and Hebbachi, A. M. (2004). "Synthesis and function of hepatic very-low-density lipoprotein." *Biochemical Society Transactions*. 32 (1): 59–64.
223. Dashti, M.; Kulik, W.; Hoek, F.; Veerman, E.C.; Peppelenbosch, M. P. and Rezaee, F. (2011). "A phospholipidomic analysis of all defined human plasma lipoproteins." *Sci Rep*. 1: 139.
224. Abuja, P. M. and Albertini, R. (2001). Methods for monitoring oxidative stress, lipid peroxidation, and oxidative resistance of lipoproteins. *Clin. Chim. Acta*, 306: 1-17.
225. Shankaraiah, P. and Reddy, Y. N. (2011). α amylase expressions in Indian type-2 diabetic patients. *Journal of Medical Sciences*. 11 (7): 280-284.
226. Windish, W. W. and Mhatre, N. S. (1965). AMP-activated protein kinase, a metabolic master switch: Possible roles in type 2 diabetes. *American Journal of Physiology*. 277: 1-10.
227. Van der Maarel, M. J. E. C.; Van der Veen, B.; Uitdehaag, J. C. M.; Leemhuis, H. and Dijkhuizen, L. (2002). Properties and applications of starch converting enzymes of the α amylase family. *Journal of Biotechnology*. 94: 137- 155.
228. Gupta, R.; Gigras, P.; Mohapatra, H.; Goswami, V. K. and Chauhan, B. (2003). Microbial α amylases: A biotechnological perspective. *Process Biochem*. 38: 1599-1616.
229. Alexander, R. (1992). Maltodextrins: Production, properties and applications. In: starch hydrolysis products: worldwide technology, products and applications, schenck, F. W. Hebeda, R. E. (eds). VCH Publishers: New York; 233-276.
230. Annison, G. and Topping, D. L. (1994). Nutritional role of resistant starch: chemical structure vs physiological function. *Annual Review of Nutrition*. 14: 297-320.
231. Cheng, H. H.; Chen, H. T.; Tung, Y. C and Shieh, M. J. (1994). Different cereals as carbohydrate sources influence lipid metabolism of hypercholesterolemia rats. *Journal of the Chinese Chemical Society*. 32: 575-588.
232. Kabir, M.; Rizkalla, S. W.; Champ, M.; Lun, J.; Boillot, J. and Bruzzo, F. (1998). Dietary amylase-amylpectin starch content affects glucose and lipid metabolism in adipocytes of normal and diabetic rats. *Journal of Nutrition*. 128: 35-43.
233. Kim, W. K.; Cnung, M. K.; Kang, N. E.; Kim, M. H. and Park, Q. J. (2003). Effect of resistant starch from corn or rice on glucose control. Colonic events and blood lipid concentration in streptozotocin-induced diabetic rats. *Journal of Nutritional Biochemistry*. 14: 166-172.

234. Mohammed, A. T. M. (2010). Biochemical studies on rice. Ph.D. thesis, Department of Agriculture Biochemistry. Faculty of Agriculture. Cairo Univ. Egypt.
235. Purushothama, S.; Raina, P.L.; Hariharan, K. (1995) Effect of long term feeding of rice bran oil upon lipids and lipoproteins in rats. *Molecular and Cellular Biochemistry*.146: 63-69.
- 236.El-Shamei, Z. S.; Gab-Alla, A. A.; Shatta, A. A.; Moussa, E. A. and Rayan, A. M. (2012). Histopathological changes in some organs of male rats fed on genetically modified corn (Ajeeb YG). *Journal of American Science*. 8(10): 684-696.
- 237.Magri, C. J. and Fava, S. (2009). The role of tubular injury in diabetic nephropathy. *European Journal of Internal Medicine*. 20: 551-555.



Appendixes



Appendix 1. a.: Daily Water consumption in rice starch factory (December 2011& January 2012)

Date	Start	End	Vol. of Water (m ³)	Date	Start	End	Vol. of Water (m ³)
December 2011				January 2012			
1. *	129750	129800	50	1.	144246	144706	460
2. *	129800	129853	53	2.	144706	145246	540
3. *	129853	129902	49	3.	145246	145796	550
4. *	129902	129951	49	4.	145796	146352	556
5. *	129951	130001	50	5.	146352	146967	615
6. *	130001	130055	54	6.	146967	147507	540
7. *	130055	130102	47	7.	147507	148073	566
8.	130102	130188	86	8.	148073	148588	515
9.	130188	130547	359	9.	148588	149213	625
10.	130547	131209	662	10.	149213	149818	605
11.	131209	131881	672	11.	149818	150388	570
12.	131881	132431	550	12.	150388	150888	500
13.	132431	133141	710	13.	150888	151493	605
14.	133141	133741	600	14.	151493	152040	547
15.	133741	134453	712	15.	152040	152668	628
16.	134453	135153	700	16.	152668	153226	558
17.	135153	135793	640	17.	153226	153789	563
18.	135793	136508	715	18.	153789	154329	540
19.	136508	137151	643	19.	154329	154929	600
20.	137151	137763	612	20.	154929	155504	575
21.	137763	138444	681	21.	155504	155983	479
22.	138444	139039	595	22.	155983	156283	300
23.	139039	139664	625	23. *	156283	156338	55
24.	139664	140214	550	24. *	156338	156388	50
25.	140214	140764	550	25. *	156388	156437	49
26.	140764	141317	553	26. *	156437	156485	48
27.	141317	142008	691	27. *	156485	156533	48
28.	142008	142715	707	28. *	156533	156583	50
29.	142715	143346	631	29. *	156583	156632	49
30.	143346	143846	500	30. *	156632	156682	50
31.	143846	144246	300	31. *	156682	156732	50
Average	(13658 -1100) / 22 = 570			Average	(11277 – 1000) / 20 = 513		

- Average water consumption =(total exactly daily water consumption exp first and end work day consumption) – (50 x number of working day exp first and end work day) / number of working day exp first and end work day
- *Daily consumption during the production process is stopped

Appendix1. b.: Daily Water consumption in rice starch factory (February 2012& April 2012)

Date	Start	End	Vol. of Water (m ³)	Date	Start	End	Vol. of Water (m ³)
February 2012				April 2012			
1. *	156732	156783	51	1.	171502	171582	80
2. *	156783	156833	50	2.	171582	172262	680
3. *	156833	156883	50	3.	172262	172946	684
4. *	156883	156931	48	4.	172946	173601	655
5. *	156931	156984	53	5.	173601	174273	672
6.	156984	157069	85	6.	174273	174965	692
7.	157069	157557	488	7.	174965	175617	652
8.	157557	158219	662	8.	175617	176209	592
9.	158219	158785	566	9.	176209	176808	599
10.	158785	159408	623	10.	176808	177476	668
11.	159408	160085	677	11.	177476	178101	625
12.	160085	160647	562	12.	178101	178699	598
13.	160647	161229	582	13.	178699	179314	615
14.	161229	161798	569	14.	179314	179989	675
15.	161798	162410	612	15.	179989	180550	561
16.	162410	162988	578	16.	180550	181125	575
17.	162988	163632	644	17.	181125	181699	574
18.	163632	164187	555	18.	181699	182357	658
19.	164187	164767	580	19.	182357	183057	700
20.	164767	165369	602	20.	183057	183662	605
21.	165369	165994	625	21.	183662	184237	575
22.	165994	166635	641	22.	184237	184739	502
23.	166635	167247	612	23.	184739	185039	300
24.	167247	167833	586	24.	185039	185095	56
25.	167833	168491	658	25.	185095	185107	12
26.	168491	169106	615	26.	185107	185147	40
27.	165480	169566	460	27.	185147	185197	50
28.	169566	169942	376	28.	185197	185249	52
Average	(12497 – 1050) / 21 = 545			29.	185249	185299	50
				30.	185299	185348	49
				Average	(13157 – 1050) / 21 = 576		

Appendix1. c.: Daily Water consumption in rice starch factory (July 2012& September 2012)

Date	Start	End	Vol. of Water (m ³)	Date	Start	End	Vol. of Water (m ³)
July 2012				September 2012			
1.	188448	188538	90	1.	204073	204123	50
2.	188538	189163	625	2.	204123	204173	50
3.	189163	189861	698	3.	204173	204221	48
4.	189861	190506	645	4.	204221	204271	50
5.	190506	191161	655	5.	204271	204322	51
6.	191161	191845	684	6.	204322	204369	47
7.	191845	192530	685	7.	204369	204419	50
8.	192530	193208	678	8.	204419	204504	85
9.	193208	193811	603	9.	204504	205054	550
10.	193811	194413	602	10.	205054	205672	618
11.	194413	195014	601	11.	205672	206264	592
12.	195014	195672	658	12.	206264	206839	575
13.	195672	196272	600	13.	206839	207394	555
14.	196272	196914	642	14.	207394	207983	589
15.	196914	197519	605	15.	207983	208588	605
16.	197519	198169	650	16.	208588	209187	599
17.	198169	198872	703	17.	209187	209749	562
18.	198872	199498	626	18.	209749	210301	552
19.	199498	200119	621	19.	210301	210876	575
20.	200119	200748	629	20.	210876	211463	587
21.	200748	201347	599	21.	211463	212060	597
22.	201347	201847	500	22.	212060	212638	578
23.	201847	202075	228	23.	212638	213235	597
24. *	202075	202129	54	24.	213235	213806	571
25. *	202129	202179	50	25.	213806	214390	584
26. *	202179	202226	47	26.	214390	215045	655
27. *	202226	202276	50	27.	215045	215653	608
28. *	202276	202326	50	28.	215653	216246	593
29. *	202326	202375	49	29.	216246	216764	518
30. *	202375	202425	50	30.	216764	217098	334
31. *	202425	202473	48				
Average	(13309 - 1050) / 21 = 583			Average	(12260 - 1050) / 21 = 534		

Appendix1. d.: Daily Water consumption in rice starch factory (October 2012& December 2012)

Date	Start	End	Vol. of Water (m3)	Date	Start	End	Vol. of Water (m3)
October 2012				December 2012			
1.	217098	217398	300	1. *	232125	232175	50
2.	217398	217952	554	2. *	232175	232225	50
3.	217952	218557	605	3. *	232225	232272	47
4.	218557	219209	652	4. *	232272	232320	48
5.	219209	219809	600	5. *	232320	232366	46
6.	219809	220284	475	6. *	232366	232416	50
7.	220284	220934	650	7. *	232416	232471	55
8.	220934	221559	625	8.	232471	232553	82
9.	221559	222208	649	9.	232553	233068	515
10.	222208	222799	591	10.	233068	233670	602
11.	222799	223399	600	11.	233670	234274	604
12.	223399	224049	650	12.	234274	234894	620
13.	224049	224674	625	13.	234894	235504	610
14.	224674	225274	600	14.	235504	236092	588
15.	225274	225839	565	15.	236092	236676	584
16.	225839	226415	576	16.	236676	237262	586
17.	226415	226985	570	17.	237262	237844	582
18.	226985	227554	569	18.	237844	238436	592
19.	227554	228169	615	19.	238436	239036	600
20.	228169	228774	605	20.	239036	239605	569
21.	228774	229374	600	21.	239605	240202	597
22.	229374	229962	588	22.	240202	240795	593
23.	229962	230217	255	23.	240795	241380	585
24. *	230217	230272	55	24.	241380	241964	584
25. *	230272	230324	52	25.	241964	242549	585
26. *	230324	230375	51	26.	242549	243104	555
27. *	230375	230425	50	27.	243104	243666	562
28. *	230425	230475	50	28.	243666	244817	572
29. *	230475	230524	49	29.	244817	245396	579
30. *	230524	230574	50	30.	245396	245905	509
31. *	230574	230622	48	31.	245905	246180	275
Average	(12564 – 1050) / 21=548			Average	(12773 – 1050) /21=558		

Appendix1. e.: Daily Water consumption in rice starch factory (May 2013& June 2013)

Date	Start	End	Vol. of Water (m ³)	Date	Start	End	Vol. of Water (m ³)
May 2013				June 2013			
1.	252248	252295	47	1.	265855	266031	176
2.	252295	252344	49	2.	266031	266543	512
3.	252344	252395	51	3.	266543	267168	625
4.	252395	252445	50	4.	267168	267844	676
5.	252445	252496	51	5.	267844	268453	609
6.	252496	252546	50	6.	268453	269052	599
7.	252546	252598	52	7.	269052	269652	600
8.	252598	252687	89	8.	269652	270241	589
9.	252687	253236	549	9.	270241	270825	584
10.	253236	253851	615	10.	270825	271414	589
11.	253851	254475	624	11.	271414	272005	591
12.	254475	255059	584	12.	272005	272617	612
13.	255059	255675	616	13.	272617	273222	605
14.	255675	256290	615	14.	273222	273810	588
15.	256290	256885	595	15.	273810	274400	590
16.	256885	257435	550	16.	274400	274977	577
17.	257435	258005	570	17.	274977	275610	633
18.	258005	258595	590	18.	275610	276183	573
19.	258595	259184	589	19.	276183	276807	624
20.	259184	259759	575	20.	276807	277400	593
21.	259759	260350	591	21.	277400	278012	612
22.	260350	260969	619	22.	278012	278589	577
23.	260969	261536	567	23.	278589	279111	522
24.	261536	262124	588	24.	279111	279435	324
25.	262124	262725	601	25. *	279435	279493	58
26.	262725	263323	598	26. *	279493	279545	52
27.	263323	263869	546	27. *	279545	279595	50
28.	263869	264460	591	28. *	279595	279642	47
29.	264460	265043	583	29. *	279642	279690	48
30.	265043	265542	499	30. *	279690	279740	50
31.	265542	265855	313				
Average	(12855 – 1100) /22=534			Average	(13080 – 1100) / 22=544		

Appendix1. f.: Daily Water consumption in rice starch factory (November 2013& March 2014)

Date	Start	End	Vol. of Water (m3)	Date	Start	End	ol. of Water (m3)
November 2013				March 2013			
1.	285900	285947	47	1.	308173	308339	166
2.	285947	285998	51	2	308339	308879	540
3.	285998	286049	51	3	308879	309504	625
4.	286049	286099	50	4	309504	310110	606
5.	286099	286150	51	5	310110	310710	600
6.	286150	286199	49	6	310110	311298	588
7.	286199	286251	52	7	311298	311998	700
8.	286251	286330	79	8	311998	312586	588
9.	286330	286889	559	9	312586	313180	594
10.	286889	287494	605	10	313180	313780	600
11.	287494	288098	604	11	313780	314359	579
12.	288098	288698	600	12	314359	314962	603
13.	288698	289285	587	13	314962	315554	592
14.	289285	289873	588	14	315554	316225	671
15.	289873	290468	595	15	316225	316769	544
16.	290468	291018	550	16	316769	317338	569
17.	291018	291608	590	17	317338	317922	584
18.	291608	292188	580	18	317922	318573	651
19.	292188	292787	599	19	318573	319273	700
20.	292787	293362	575	20	319273	319954	681
21.	293362	293976	614	21	319954	320540	586
22.	293976	294591	615	22	320540	321117	577
23.	294591	295172	581	23	321117	321649	532
24.	295172	295758	586	24	321649	321948	299
25.	295758	296359	601	25	321948	322003	55
26.	296359	296938	579	26	322003	322056	53
27.	296938	297525	587	27	322056	322105	49
28.	297525	298135	610	28	322105	322153	48
29.	298135	298750	615	29	322153	322203	50
30.	298750	299073	323	30	322203	322253	50
				31	322253	322303	50
Average	(12822 -- 1050) /21=541			Average	(13310 -- 1100) / 22=555		

Appendix2. a.: Flow rate of wet cleaning wastewater from discharge point

Date	No.	Vol. of Water (litre)	Duration Time(sec.)	Flow rate (L/sec.)
10/12/2011	1	15.000	6.5	2.31
	2	15.500	6.8	2.28
	3	14.400	6.4	2.25
18/2/2012	1	16.100	7.0	2.30
	2	15.000	6.5	2.31
	3	16.000	7.0	2.29
18/9/2012	1	15.800	6.9	2.29
	2	16.000	6.9	2.32
	3	15.500	6.8	2.28
15/6/2013	1	15.800	6.9	2.29
	2	15.700	6.8	2.31
	3	16.800	7.4	2.27

Average Flow rate = 2.291 L/sec

The average working time = 80 min/day

Wastewater discharged per day= 11 m³/day

Appendix2. b.: The flow rate of soaking wastewater discharging from discharge point

Date	No.	Vol. of Water (litre)	Duration Time(sec.)	Flow rate (L/sec.)
10/12/2011	1	17.700	6.5	2.72
	2	18.200	6.8	2.69
	3	16.000	5.9	2.71
18/2/2012	1	17.800	6.6	2.70
	2	15.000	5.5	2.71
	3	18.000	6.6	2.70
18/9/2012	1	15.000	5.5	2.73
	2	17.800	6.6	2.68
	3	16.100	6.0	2.70
15/6/2013	1	17.400	6.4	2.74
	2	16.100	5.9	2.71
	3	15.000	5.5	2.71

Average Flow rate for basin =2.708 L/sec , The average working time of basin 20min/day

Number of soaking basin per day 4 , Soaking wastewater discharged per day= 13 m³/day

Appendix2. c.: Flow rate of clean water discharging from the first centrifuge process using plastic bag and stopwatch

Date	No.	Vol. of Water (litre)	Duration Time(sec.)	Flow rate (L/sec.)
25/12/2011	1	18.200	3.5	5.20
	2	22.400	4.3	5.21
	3	26.000	5.0	5.20
15/7/2012	1	23.500	4.5	5.22
	2	20.250	3.9	5.19
	3	17.600	3.4	5.18
20/12/2012	1	23.800	4.6	5.17
	2	21.400	4.1	5.22
	3	16.000	3.1	5.16
9/6/2013	1	22.800	4.4	5.18
	2	23.800	4.5	5.29
	3	22.300	4.3	5.19

Average Flow rate = 5.2 L/sec, The average working time on 16 hour/day, Clean water discharged per day= 300 m³/day

Appendix2. d.: Flow rate of wastewater discharging from the first centrifuge process using plastic bag and stopwatch

Date	No.	Vol. of Water (litre)	Duration Time(sec.)	Flow rate (L/sec.)
22/12/2011	1	15.900	3.0	5.3
	2	21.000	4.0	5.25
	3	24.100	4.6	5.24
15/2/2012	1	23.400	4.5	5.20
	2	21.400	4.1	5.22
	3	23.800	4.5	5.29
18/9/2012	1	22.800	4.4	5.18
	2	22.300	4.3	5.19
	3	21.400	4.1	5.22
15/5/2013	1	24.200	4.6	5.26
	2	18.200	3.5	5.20
	3	21.000	4.0	5.25

Average Flow rate = 5.233 l/sec, The average working time 8 hour/day , waste water discharged per day= 150 m³/day

Appendix3. a.: Chemical composition of broken rice (grade zero)

Date	Crude Protein	Fat	Ash	Crude Fiber	Total Carbohydrate
24/11/2011	8.65	0.699	1.92	1.52	87.211
25/2/2012	8.80	0.769	1.71	1.43	87.291
10/7/2012	8.62	0.749	1.84	1.51	87.281
20/12/2012	8.79	0.743	1.85	1.44	87.177
12/5/2013	8.75	0.745	1.81	1.51	87.185
Average	8.72	0.74	1.83	1.48	87.23

* As dry weight

* Available carbohydrate by difference

Moisture content of grade zero was 11.89%, 11.95 %, 11.90%, 11.84% and 11.89 respectively

Appendix3. b.: Chemical composition of broken rice (grade one)

Date	Crude Protein	Fat	Ash	Crude Fiber	Total Carbohydrate
24/11/2011	8.82	0.696	1.95	1.57	86.964
25/2/2012	8.78	0.869	1.87	1.46	87.021
10/7/2012	8.94	0.763	1.91	1.39	86.997
20/12/2012	8.72	0.872	1.83	1.46	87.118
12/5/2013	8.75	0.725	2.05	1.62	86.855
Average	8.80	0.79	1.92	1.50	86.99

* As dry weight

* Available carbohydrate by difference

Moisture content of grade zero was 11.85%, 11.78%, 11.67%, 11.75% and 11.90 respectively

Appendix4. a.: Characteristics of Wet cleaning wastewater

Date	pH	T.S	T.SS	T.DS	COD	BOD	Temp °C
22/12/2011	6.23	32105	21299	10806	19095	12600	24
06/07/2012	6.00	32484	21726	10758	19325	12779	27
11/10/2012	6.84	29114	20290	8824	17489	11737	26
11/12/2012	6.68	29878	20699	9179	18000	12000	24
15/05/2013	6.58	28807	19869	8938	17304	11594	26
15/3/2014	6.56	27860	19232	8628	16789	11290	25
Average	6.48	30041	20519.3	9522.16	18000	12000	25.33

Appendix4. b.: Chemical analysis of Wet cleaning wastewater

Date	T.S	Crude protein	Fat	Crude Fiber	Ash	Total Carbohydrate	Sodium chloride
22/12/2011	3.21	0.668	0.149	0.221	0.190	1.982	0.069
06/07/2012	2.91	0.654	0.151	0.198	0.187	1.720	0.065
11/10/2012	2.90	0.612	0.139	0.189	0.205	1.755	0.076
11/12/2012	2.98	0.619	0.152	0.215	0.205	1.789	0.051
15/05/2013	3.11	0.648	0.158	0.213	0.201	1.890	0.065
15/3/2014	2.89	0.641	0.134	0.200	0.208	1.707	0.064
Average	3.00	0.640	0.147	0.206	0.199	1.808	0.065

Appendix4. c.: Characteristics of soaking wastewater

Date	PH	T.S	T.S.S	T.D.S	COD	BOD	Temp °C
18/1/2012	4.14	44178	28845	15333	23999	16125	23
6/4/2012	4.60	44430	29067	15363	24100	16283	24
18/9/2012	4.30	40800	26212	14588	21995	14562	25
16/10/2012	4.62	45800	29354	16446	24924	16495	24.5
15/12/2012	4.48	46548	29779	16769	25678	17059	23
10/6/2013	4.70	46559	29795	16764	25704	17072	24
Average	4.47	44719	28842	15877	24400	16.266	23.9

Appendix4. d.: Chemical composition of soaking wastewater

Date	T.S	Crude protein	Fat	Crude Fiber	Ash	Total Carbohydrate	Sodium chloride
18/1/2012	4.418	1.612	0.089	0.319	0.291	2.107	0.085
6/4/2012	4.443	1.504	0.088	0.302	0.301	2.248	0.095
18/9/2012	4.080	1.425	0.099	0.297	0.301	1.958	0.099
16/10/2012	4.580	1.529	0.098	0.298	0.275	2.38	0.115
15/12/2012	4.655	1.651	0.090	0.286	0.289	2.339	0.103
10/6/2013	4.656	1.669	0.088	0.298	0.271	2.330	0.097
Average	4.472	1.565	0.092	0.300	0.288	2.227	0.099

Appendix4. e.: Characteristics of first centrifuge wastewater

Date	PH	T.S	T.S.S	T.D.S	COD	BOD	Temp °C
9/2/2012	9.68	7540	3170	4370	5290	3550	28
5/4/2012	9.68	8199	3375	4824	5899	3925	30
12/7/2012	9.63	8055	3362	4693	5790	3845	32
24/9/2012	9.88	8645	3195	5450	6066	4140	31
21/12/2012	9.40	8002	3383	4619	5740	3790	31
25/5/2012	9.39	8819	3225	5594	6315	4150	31
Average	9.61	8210	3285	4925	5850	3900	30.5

Appendix4. f.: Chemical composition of first centrifuge wastewater

Date	T.S	Crude protein	Fat	Crude Fiber	Ash	Total Carbohydrate	Sodium chloride
9/2/2012	0.754	0.262	0.010	0.049	0.034	0.399	0.012
5/4/2012	0.819	0.346	0.012	0.043	0.039	0.379	0.014
12/7/2012	0.805	0.324	0.009	0.052	0.040	0.380	0.015
24/9/2012	0.864	0.391	0.010	0.059	0.046	0.358	0.014
21/12/2012	0.800	0.340	0.010	0.049	0.038	0.363	0.015
25/5/2012	0.881	0.419	0.009	0.042	0.049	0.362	0.014.
Average	0.821	0.347	0.010	0.049	0.041	0.374	0.014

Appendix4. g.: Characteristics of dilute screen milling waste water

Date	PH	T.S	T.S.S	T.D.S	COD	BOD	Temp °C
23/1/2012	7,5	58113	56056	2057	29206	19158	25
1/3/2012	7,73	54539	52610	1929	27410	17980	25.5
24/10/2012	8,15	57648	55654	2030	28991	19017	24.5
1/1/2013	7,38	54547	52602	1945	27414	17983	25.5
25/6/2013	8.00	56218	54229	1989	28254	18534	23.5
1/12/2013	8.04	56207	54223	1984	28249	18530	26
Average	7.8	56218	54229	1989	28254	18534	25

Appendix4. f.: Chemical composition of dilute screen milling wastewater

Date	T.S	Crude protein	Fat	Crude Fiber	Ash	Total Carbohydrate	Sodium chloride
23/1/2012	5.811	0.654	0.039	0.599	0.402	4.117	0.105
1/3/2012	5.454	0.645	0.042	0.590	0.402	3.775	0.114
24/10/2012	5.765	0.641	0.042	0.596	0.411	4.075	0.115
1/1/2013	5.455	0.646	0.042	0.585	0.390	3.792	0.099
25/6/2013	5.622	0.651	0.033	0.591	0.396	3.951	0.113
1/12/2013	5.621	0.639	0.036	0.579	0.393	3.974	0.120
Average	5.622	0.646	0.039	0.590	0.399	3.947	0.111

Appendix4. h. Characteristics of concentrate screen milling waste water

Date	PH	T.S	T.S.S	T.D.S	COD	BOD	Temp °C
23/1/2012	7,5	244122	236937	7485	64668	32774	25
1/3/2012	7,73	229111	222368	6743	60692	30759	25.5
24/10/2012	8,15	242324	235239	7463	64192	32533	24.5
1/1/2013	7,38	229145	222337	6808	60701	30764	25.5
25/6/2013	8.00	236164	229214	6672	62560	31706	23.5
1/12/2013	8.04	236118	229189	6529	62548	31700	26
Average	7.8	236164	229214	6950	62560	31706	25

Appendix4. i.: Chemical composition of concentrate screen milling wastewater

Date	T.S	Crude protein	Fat	Crude Fiber	Ash	Total Carbohydrate	Sodium chloride
23/1/2012	24.412	2.749	0.157	3.477	1.99	16.039	0.105
1/3/2012	22,911	2.717	0.176	3.458	1.79	14.770	0.114
24/10/2012	24.232	2.700	0.171	3.503	2.03	15.828	0.115
1/1/2013	22.915	2.721	0.173	3.452	1.85	14.719	0.099
25/6/2013	23.616	2.742	0.136	3.499	1.87	15.369	0.113
1/12/2013	23.612	2.691	0.147	3.491	1.95	15.333	0.120
	236164	2.72	0.160	3.480	1.91	15.343	0.111

Appendix5. Egyptian Environmental Legal Requirements for Industrial Wastewater

Parameter (mg/L) unless otherwise noted Law4/94	Discharge Coastal Environment Law 93/62	Discharge to Sewer System (as modified by Decree 44/2000)	Law 48/82: Discharge into :			
			Underground Reservoir & Nile Branches/Canal s Nile	Main Stream	Drains	
					Municipa l	Industri al
BOD (5day, 20 deg.)	60	<600	20	30	60	60
COD	100	<1100	30	40	80	100
PH	6-9	6-9.5	6-9	6-9	6-9	6-9
Oil & Grease	15	<100	5	5	10	10
Temperature (deg.	10C>avg. temp of receiving body	<43	35	35	35	35
Total Suspended Solids	60	<800	30	30	50	60
Settable Solids	—	8cm³/l (10min) 15 cm³/l (30 min)	—	20	—	—
PO₄	5	-----	—	1	---	10
Total phosphorus	-----	25	-----	----	-----	-----
Fluoride	1	----	0.5	0.5	----	0.5
Total Dissolved Solids	2000	----	800	1200	2000	2000



Arabic Summary



بصورة طبيعية في جميع مجموعات المعاملة بينما لوحظ ظهور بعض التجمعات البروتينية و سماكة في الأنابيب في مجموعة المعاملة ١٠ %.

وقد أوصت الدراسة بما يأتي:

- أهمية تقليل كمية المياه الخام المستخدمة في صناعة نشا الأرز وانخفاض مياه الصرف الصناعي للمحافظة على مصادر المياه و خفض تكاليف الإنتاج.
- استخدام المستخلصات العضوية من مياه صرف مصنع نشا الأرز والتي تحتوي على نسبة عالية من البروتين كإضافات لتغذية الحيوانات لرفع إيرادات الشركة .

أولاً: التأثير على معدل استهلاك الغذاء و الماء و نسبة الزيادة في الوزن إلى وزن الجسم و القيمة الغذائية

تبين من نتائج الدراسة عدم وجود فروق معنوية في كمية المياه المستهلكة و في كفاءة التحول الغذائي بينما كان هناك فروق معنوية سالبة في كميات الغذاء المستهلك و فروق معنوية موجبة في معدل الزيادة في الوزن في الحيوانات المعاملة.

ثانياً: التأثير على خواص الدم:

تبين من نتائج الدراسة

- عدم وجود فروق معنوية في معظم المتغيرات الهيماتولوجية بين المجموعة الضابطة و مجاميع المعاملة ٥ ، ١٠ ، ٢٠% بينما ظهرت فروق معنوية موجبة في عدد كرات الدم البيضاء
- عدم وجود فروق معنوية في مستوى البروتين الكلي ، الجلوبيولين ، الألبومين ، الجلوكوز ، اللييدات الكلية ، الليوبروتين عالي الكثافة و في نشاط إنزيمات الفوسفاتيز القاعدي ، الألفا أميليز و أسبرتيت أمينو ترانس فيريز.
- ظهرت فروق معنوية سالبة في مستوى البيليروبين ، الكولسترول ، الجلوسيدات الثلاثية ، الليوبروتين منخفض الكثافة ، الليوبروتين شديد الانخفاض في الكثافة ، الفوسفاتيز الحامضي و الثيوباربيتوريك
- ظهرت فروق معنوية موجبة في نشاط إنزيم الأنين أمينو ترانس فيريز بين المجموعة الضابطة و مجاميع المعاملة ٥ و ٢٠% بينما لم تظهر فروق معنوية بين المجموعة الضابطة و مجموعة المعاملة ١٠%

ثالثاً: التأثير على الوزن النسبي للأعضاء

تبين عدم وجود فروق معنوية في نسب وزن الأعضاء الداخلية بالجسم (الكبد – الكلى). بينما ظهرت فروق معنوية سالبة في نسبة وزن القلب إلى الوزن الكلي

رابعاً: التأثير على أنسجة الأعضاء الداخلية (الكبد – الكلى)

الفحص الميكروسكوبي لأنسجة الكبد و الكلى لم يظهر اختلافات جوهرية في خلايا الكبد و الكلى بين المجموعة الضابطة و مجموعات المعاملة حيث أشار الفحص الميكروسكوبي لخلايا الكبد أن الخلايا كانت طبيعية ولم يظهر بها أي تلفيات و ظهر الوريد المركزي بصورة عادية ولم يظهر به أي التهابات أو تقرحات بينما ظهر زيادة قليلة في خلايا كوبر و تمدد في بعض الجيوب كما ظهرت إتهابات بسيطة في مجموعة المعاملة ١٠%. بينما أشار الفحص الميكروسكوبي لخلايا الكلى إلى ظهور كبيبات الكلى بصورة طبيعية ولم يظهر بها أي ضمور في الأنابيب أو الخلايا وظهرت الأنسجة الخارجية و الداخلية

الكربوهيدرات و البروتينات و انخفاض محتواها من الدهون وكلوريد الصوديوم . وقد تم دراسة خواص تلك المياه من حيث محتواها من المادة الصلبة الكلية ، والمواد العالقة والذائبة ، وقياس كمية الأكسجين الحيوي و الكيميائي المستهلك وتبين ارتفاع محتوى تلك المياه من المواد العالقة وارتفاع قيمة الأكسجين الحيوي و الكيميائي المستهلك.

تم إجراء تجارب لاستخلاص المادة العضوية الذائبة والعالقة في مياه الصرف الصناعي الناتجة عن مصنع النشا باستخدام نقطة التعادل الكهربائي ودراسة الزمن الأنسب لترسيب تلك المواد بالجاذبية وقد تبين أن أنسب درجة حموضة لترسيب أكبر كمية من البروتينات هي ٤.٥ و تبين أن أفضل زمن لترسيب أكبر كمية من المواد العالقة هو ١٨٠ دقيقة ثم تم فصل الراسب بواسطة آلة الطرد المركزي. وقد كانت نسبة البروتينات المفصولة ٧٠% من البروتينات الكلية بينما وصلت نسبة المواد العالقة ٩٥% من المواد العالقة الكلية.

تم تجفيف الراسب بواسطة البخار ثم تم تقدير محتواه من الرطوبة ، الكربوهيدرات ، البروتينات ، الدهون ، الألياف والرماد وكذلك تم تقدير محتواه من الفينولات والأفلاتوكسينات وقد تبين أن الرطوبة (١٢ %) ، الكربوهيدرات (٦٠.٥ %) ، البروتينات (١٦.٨%) ، الدهون (١.١٨%) ، الألياف (٣.٣٨%) والرماد (٦.١٢%) وكانت الفينولات (١.٤٩ مجم / جم) وتبين خلو الناتج من الأفلاتوكسينات.

للتعرف على القيمة الغذائية للناتج المستخلص المجفف تم إجراء تجربة على الجرزان معمليا حيث كان تصميم التجربة استخدام ٤٠ من ذكور الجرزان من سلالة الألبينو (٢٠٠ ± ٥٠) جرام تم وضعها في أقفاص و تهيئة الظروف المناسبة من ماء وغذاء وتركت الحيوانات للتأقلم على البيئة لمدة ٥ أيام وقسمت إلى ٤ مجموعات عشوائيا : المجموعة الأولى وتمثل المجموعة الضابطة ، المجموعة الثانية تم إضافة المنتج الثانوي الجديد المستخلص من مياه الصرف الصناعي إلى العليقة القياسية المستخدمة في تغذية المجموعة الضابطة بنسبة ٥% ، المجموعة الثالثة تم إضافة المنتج المستخلص بنسبة ١٠%، المجموعة الرابعة تم إضافة المنتج المستخلص بنسبة ٢٠% وقد إستمرت التجربة لمدة ٨ أسابيع تم خلالها تقدير وزن الجسم ومعدل استهلاك الغذاء و الماء أسبوعيا كذلك تم تقدير معدلات النمو وكفاءة التحول الغذائي وفي نهاية التجربة تم ذبح الجرزان وتم جمع الدم من الشريان الأورطي وتم فصل ١ ملي تم إضافة الهيبارين إليه لإجراء الدراسات الهيماتولوجية. وتم جمع الدم في أنابيب وترك ليتجلط ثم تم استخدام الطرد المركزي ١٠٠٠ لفة لفصل السيرم ثم تم حفظه على درجة -٢٠ في الثلجة ، وفي السيرم تم تقدير مستوى الجلوكوز ومستوى الدهون ومشتقاتها ووظائف الكبد والكلية و إنزيم الألفا أميليز وحامض الثيوباربيتوريك ثم تم تحليل النتائج إحصائيا وكانت كالاتي:

الملخص العربي

الشركة المصرية للنشا و الخميرة و المنظفات إحدى الشركات التابعة للشركة القابضة للصناعات الغذائية ويقع المركز الرئيسى للشركة في منطقة السيوف بشرق محافظة الإسكندرية بجمهورية مصر العربية. تضم الشركة ٤ مصانع لإنتاج المنظفات و الخميرة و المواد الكيميائية و المواد المساعدة و مصنع النشا. والسعة القصوى لمصنع النشا هي ٣٥-٣٠ طن من النشا و النشا المعدل يوميا. و المنتجات الرئيسية هي أنواع مختلفة من النشا الصناعي، نشا الطعام و المواد النشوية و النشا المعدل الذى يستخدم فى الصناعات مثل صناعة المنسوجات. تنتج الشركة النشا من كسر الأرز عن طريق نقع كسر الأرز في الماء المضاف إليه محلول الصودا الكاوية المخفف داخل أحواض ثم يتم فصل النشا عن طريق آلات طرد مركزي ثم يجفف بواسطة البخار.

الهدف من هذه الدراسة هو خفض تكاليف الإنتاج من خلال تخفيض كمية استهلاك المياه في العمليات الصناعية . ودراسة إمكانية إعادة تدوير المياه ورفع عائدات الشركة باستخدام المواد العضوية المنفصلة من مياه الصرف الصناعى كإضافات لتغذية الحيوانات.

يستهلك مصنع النشا كميات كبيرة من الماء في مراحل الإنتاج المختلفة ، تم تقدير تلك الكميات المستهلكة يوميا خلال مدة الدراسة وتبين أن متوسط الاستهلاك اليومي هو ٥٥.٥ م^٣ / طن كسر أرز وذلك أعلى ٤.٥ مرة من المعدل العالمي. وتبين من الدراسة أن الفاقد الأعلى من المياه كان في مرحلة الطرد المركزي والتي كان متوسط الاستهلاك اليومي لها ٣٢٥ م^٣ / يوم منها ٣٠٠ م^٣ عبارة عن مياه نظيفة تستخدم لتبريد آلة الطرد المركزي أثناء توقف تغذية الفراز بمحلول النشا (لين النشا) والتي وصلت إلى ١٦ ساعة يوميا خلال مدة الدراسة وهذه المياه يمكن إعادة تدويرها لتقليل كمية المياه المستهلكة يوميا بالإضافة إلى وجود استهلاك عالي من المياه في عملية التنظيف الرطب و الغسيل نتيجة استخدام طرق بدائية وعدم استخدام مدافع الغسيل التي تعمل بضغط الهواء وكذلك وجود تسرب للمياه من أحواض النقع وصمامات التحكم وعدم وجود مصائد للبخار على خطوط البخار.

تم استخدام درجتين من كسر الأرز خلال فترة الدراسة (صفر – واحد) تبين من تحليلهما كيميائيا عدم وجود اختلافات جوهرية بين الدرجتين ولكن الفروق بينهم هي فروق فيزيائية فقط (حجم الحبيبة وكمية الشوائب)

تم أخذ عينات من مياه الصرف الناتجة عن مراحل الإنتاج المختلفة وتقدير محتواها من الكربوهيدرات ، البروتينات ، الدهون ، الألياف ، كلوريد الصوديوم و الرماد وتبين ارتفاع محتوى تلك المياه من المادة العضوية وخاصة



الملخص العربي



لجنة الإشراف

د/ صباح جابر البنا

أستاذ مساعد الكيمياء الحيوية
قسم الدراسات البيئية
معهد الدراسات العليا و البحوث
جامعة الإسكندرية

د/ مرفت أمين عبد القوي

أستاذ مساعد الهندسة الكيميائية
قسم الدراسات البيئية
معهد الدراسات العليا و البحوث
جامعة الإسكندرية

د/عاطف عبد المجيد يوسف

مدير عام مصنع النشا
الشركة المصرية للنشا و الخميرة و المنظفات
مدرس الكيمياء الحيوية – جامعة فاروس



جامعة الإسكندرية
معهد الدراسات العليا و البحوث
قسم الدراسات البيئية



الاستفادة من مخلفات صناعة نشا الأرز في الشركة المصرية للنشا و الخميرة و المنظفات

رسالة مقدمة إلى
معهد الدراسات العليا و البحوث - جامعة الإسكندرية
ضمن متطلبات درجة الماجستير

في
الدراسات البيئية

مقدمة من

أشرف سعيد محمد رجب

بكالوريوس زراعة - قسم تكنولوجيا الأغذية

قسم الدراسات البيئية
معهد الدراسات العليا و البحوث
جامعة الإسكندرية

٢٠١٥