

SUMMARY AND CONCLUSION

I- Summary

Magnetic Resonance Imaging (MRI) is one of the most rapidly advancing diagnostic imaging tools today. Various types of acoustic noise are produced during the operation of MR systems. The problems associated with acoustic noise for patients and healthcare workers include simple annoyance, difficulties in verbal communication, heightened anxiety, temporary hearing loss, and potential permanent hearing impairment. Acoustic noise may pose a particular hazard to specific patient groups who may be at increased risk.

This study aimed to investigate the potential MRI sound adverse effects on some biophysical (dielectric properties) and biochemical (blood glucose level, malondialdehyde and superoxide dismutase in the brain) parameters of mice. Exploring there was correlation between MRI sound power and the severity of these parameters.

In this work the following materials and methods were used:

- The actual sound arise from MRI system (closed type 1.5T) were recorded using microphone attached to laptop, then saved on mp3 device which then connected to loud speaker. The noise source intensity was adjusted to be 90, 100 and 121 dB at the loud speaker using sound level meter device.
- Anechoic chamber was designed and performed to insulate the sound. Illumination and ventilation were taken into account upon design. All the exposures were performed inside it.
- The animals used were 40 mice divided to 4 groups each of ten mice, classified as the following:
 - **Control group (GpA):** remained in the anechoic chamber for two hours daily for a week without exposure to any noise.
 - **Experimental group (GpB1):** exposed to sound intensity of 121 dB for 2 hours daily for a week.
 - **Experimental group (GpB2):** exposed to sound intensity of its *actual value* 100 dB for 2 hours daily for a week.
 - **Experimental group (GpB3):** exposed to sound intensity of 90 dB for 2 hours daily for a week.

After exposure period, mice were sacrificed and blood samples were withdrawn to determine:

- Total and differential white blood cells (WBCs) count.
- Blood glucose level.
- Brain tissues dissected, and divided into two portions, one portion for the dielectric measurements and the other portion was homogenized for determination of:
 - Brain superoxide dismutase activity to indicate the oxidative stress state.

- Brain malondialdehyde concentration to assess the lipid peroxidation state in brain cell membrane.

The results of the present work are:

- 1- Significant increase in blood glucose level in all experimental groups as compared with control group.
- 2- Significant increase in total blood WBCs and blood platelets count in all experimental groups as compared with control group.
- 3- Significant decrease in hematology markers (Hb, Ht, RBCs, MCV, MCH and MCHC) in all experimental groups as compared with control group
- 4- Significant decrease in brain superoxide dismutase activity in all experimental groups as compared with control group.
- 5- Significant increase in brain malondialdehyde concentration in all experimental groups as compared with control group.
- 6- Changes in the dielectric properties (conductivity and permittivity) in all experimental groups as compared with control group, and the degree of depressed center and the maximum values of the real conductivity and permittivity depend on the noise intensities.

II. Conclusions:

Depending on the present work results, we can conclude that acoustic noise that produced during the operation of MR systems have direct and clear effect on:

- The biochemical parameters (MDA levels, superoxide dismutase activity, and blood glucose level).
- Blood glucose level.
- Total WBCs count.
- Oxidative stress state.
- The dielectric properties of brain tissue.

Recommendation:

The simplest and least expensive means of preventing problems associated with acoustic noise during MR procedures is to encourage the routine use of disposable earplugs or headphones. Earplugs, when properly used, can abate noise by 10 to 30 dB, which is usually an adequate amount of sound attenuation for the MR environment. The use of earplugs typically provides a sufficient decrease in acoustic noise that, in turn, is capable of preventing hearing problems.

REFERENCES

1. Paley MNJ, Wilkinson ID, Beek EV, Lee KJ, Wild JM, Fischele S, et al. Magnetic resonance imaging: basic principles. In: Grainger RG, Allison D, Adam A, eds. *Diagnostic Radiology, a textbook of Medical Imaging.*, London: Chun hill Livingstone, 2001; 4th ed:101.
2. Hedeem RA, Edelstein WA. Characterization and prediction of gradient acoustic noise in MR imagers. *Magn Reson Med* 1997; 37: 7-10.
3. Mansfield P, Glover PM, Beaumont J. Sound generation in gradient coil structures for MRI. *Magn Reson Med* 1998; 39: 539-50.
4. Shellock FG, Ziarati M, Atkinson D, Chen DY. Determination of gradient magnetic field-induced acoustic noise associated with the use of echo planar and three-dimensional, fast spin echo techniques. *J Magn Reson Imaging* 1998; 8:1154-7.
5. Ravicz ME, Melcher JR, Kiang NY. Acoustic noise during functional magnetic resonance imaging. *J Acoust Soc Am* 2000;108: 1683-96.
6. Shah NJ, Steinhoff S, Mirzazade S, Zafiris O, Grosse-Ruyken ML, Jancke L, et al . The effect of sequence repeat time on auditory cortex stimulation during phonetic discrimination. *Neuroimage* 2000; 12: 100-08.
7. Baumgart F, Gaschler-Markefski B, Tempelmann C, Woldorff MG, Tegeler C, Schindler F, et al. Masking of acoustic stimuli by the gradient noise alters the fMRI detectable activation in a particular human auditory cortex. *MAGMA* 1996; 4: p185.
8. Shah NJ, Jancke L, Grosse-Ruycken ML, Pose S, Muller G HW. On the influence of acoustic masking noise in fMRI on the auditory cortex during phonetic discrimination. In: *Proceedings of the 5th Annual Meeting of ISMRM, Vancouver, Canada 1997*; p 340.
9. Elliott MR, Bowtell RW, Morris PG. The effect of scanner sound in visual, motor, and auditory functional MRI. *Magn Reson Med* 1999; 41: 1230-35.
10. Cowan J P. Educating the public on environmental and recreational noise exposure. In *Handbook of Environmental Acoustics*. New York 1994; 14-20.
11. Talbott E, Thompson SJ. Health effects from environmental noise exposure. In: *Introduction to Environmental Epidemiology*. New York: Lewis Publishers 1995; 209-19.
12. Job RFS. The influence of subjective reactions to noise on health effects of the noise. *Environment International* 1996; 22(1):93-104.
13. Westman JC, Walters JR. Noise and stress: A comprehensive approach. *Environmental Health Perspectives* 1981;41:291-309.

14. Stansfield SA. Noise, noise sensitivity and psychiatric disorder. In: Epidemiological and psychophysiological studies. Psychological Medicine Monograph Supplement (Cambridge) 1992; 22:1- 44.
15. Health and Welfare Canada. Federal-Provincial Advisory Committee on Environmental and Occupational Health National Guidelines for Environmental Noise Control Procedures and Concepts for the Drafting of Environmental Noise Regulations/By-laws in Canada. Ottawa, Ontario: Minister of Supply and Services Canada 1989.
16. Ohinata Y, Yamasoba T, Schacht J, Miller JM. Glutathione limits noise-induced hearing loss. *Hear Res* 2000; 146: 28–34.
17. Ohlemiller KK, Dugan LL. Elevation of reactive oxygen species following ischemia-reperfusion in Mouse cochlea observed in vivo. *Audiol. Neurootol* 1999;4:219-28.
18. Yamane H, Nakai Y, Takayama M, Iguchi H, Nakagawa T, Kojima A .Appearance of free radicals in guinea pigs inner ear after noise-induced acoustic trauma. *Eur Arch Otorhinolarygol* 1995;252:504-8.
19. Yamasoba T, Harris C, Shoji F, Lee RJ, Nuttal AL, Miller JM .Influence of intense sound exposure on glutathione synthesis in the cochlea. *Brain Res* 1998; 804:72-8.
20. Nielsen F, Mikkelsen BB, Nielsen JB, Andersen HR, Grandjean P. Plasma malondialdehyde as biomarker for oxidative stress: reference interval and effects of life-style factors. *Clin Chem* 1997; 43: 1209–14.
21. Predy VR, Reilly ME, Mantle D, Peters TJ. Oxidative damage in liver disease. *JIFCC*1998; 10: 16-20.
22. Weiss JM, Simson PG. Neurochemical mechanisms underlying stress-induced depression. In: Field T, McCabe P, Schneiderman N, editors. *Stress and coping*. Hillsdale: Lawrence Erlbaum 1985; 93-116.
23. Ravindran R, Devi SR, Samson J, Senthilvelan M. Noise- Stress-Induced Brain Neurotransmitter Changes and the Effect of *Ocimum sanctum* (Linn) Treatment in Albino Rats. *J Pharmacol Sci* 2005; (98): 354 -60.
24. Pfaff J, Stecker M. Loudness level and frequency content of noise in the animal house. *Laboratory Animals* 1976;10:111- 17.
25. Milligan SR, Sales GD, Khirnykh K. Sound levels in rooms housing laboratory animals: an uncontrolled daily variable. *Physiology and Behavior* 1993;53:1067- 76.
26. Baldwin AL, Primeau R, Johnson W. Effect of noise on the morphology of the intestinal mucosa in laboratory rats. *Contemporary Topics of Laboratory Animal Science* 2006;45 (1): 74-82.
27. Bies DA, Hansen CH. *Engineering noise control: theory and practice*. London Spon 2003; 3rd ed.

28. Pignatelli D, Maia M, Castro AR, Conceicao MM, Vivier J, Defaye G. Chronic stress effects on the rat adrenal cortex. *Endocr Res* 2000;26:537-44.
29. Mayer EA, Naliboff BD, Chang L. Basic pathophysiologic mechanisms in inflammatory bowel syndrome. *Digestive Diseases* 2001;19: 212-18.
30. Baldwin AL, Bell IR. Effect of noise on microvascular integrity in laboratory rats. *Journal of American Assoc. for Laboratory Animal Science* 2007;46 (1): 58-65.
31. Nuttall AL. Sound-induced cochlear ischemia/hypoxia as a mechanism of hearing loss. *Noise Health* 1999;2 :17-32.
32. Brummett RE, Talbot JM, Charuhas P. Potential hearing loss resulting from MR imaging. *Radiology* 1988;169:539-40.
33. Thorne PR, Ameratunga SN, Stewart J, Reid N, Williams W, Purdy SC et al. Epidemiology of noise-induced hearing loss in New Zealand. *New Zealand Medical Journal* 2008;121(1280): 33-44.
34. Quirk ME, Letendre AJ, Ciottoni RA, Lingley JF. Anxiety in patients undergoing MR imaging. *Radiology* 1989;170(2):463-6.
35. Reeves MJ, Brandreth M, Whitby EH, Hart AR, Paley MN, Griffiths PD et al. Neonatal cochlear function: measurement after exposure to acoustic noise during in utero MR imaging. *Radiology* 2010.
36. Bowtell RW, Mansfield PM. Quiet transverse gradient coils: Lorentz force balancing designs using geometric similitude. *Magn Reson Med* 1995;34:494-97.
37. Hurwitz R, Lane SR, Bell RA, Brant-Zawadzki MN. Acoustic analysis of gradient-coil noise in MR imaging. *Radiology* 1989;173:545-48.
38. Schmitter S, Bock M. Acoustic noise-optimized verse pulses. *Magn Reson Med*. 2010;64:1446-52.
39. Peelle JE, Eason RJ, Schmitter S, Schwarzbauer C, Davis MH. Evaluating an acoustically quiet EPI sequence for use in fMRI studies of speech and auditory processing. *Neuroimage*. 2010;52:1410-19.
40. Tomasi DG, Ernst T. Echo planar imaging at 4 Tesla with minimal acoustic noise. *J Magn Reson Imag* 2003;128-130.
41. Ulmer JL, Biswal BB, Mark LP, Mathews VP, Prost RW, Millen SJ, et al. Acoustic echo planar scanner noise and pure tone hearing thresholds: the effects of sequence repetition times and acoustic noise rates. *J Comp Assist Tomogr* 1998;22:480-86.
42. Zapp J, et al. Sinusoidal echo-planar imaging with parallel acquisition technique for reduced acoustic noise in auditory fMRI. *J Magn Reson Imag*. 2012;36:581-88.
43. Hattori Y, Fukatsu H, Ishigaki T. Measurement and evaluation of the acoustic noise of a 3 Tesla MR scanner. *Nagoya J Med Sci*. 2007;69:23-8.

44. U.S. Department of Health and Human Services, Food and Drug Administration, Center for Devices and Radiological Health, Radiological Devices Branch, Division of Reproductive, Abdominal, and Radiological Devices, Office of Device Evaluation. Guidance for Industry and FDA Staff. Criteria for significant Risk Investigations of Magnetic Resonance Diagnostic Devices, July 14, 2003.
45. International Commission on Non-Ionizing Radiation Protection (ICNIRP) Statement, Medical magnetic resonance procedures: protection of patients. *Health Physics* 2004;87:197-216.
46. Roschmann P. Human auditory system response to pulsed radio frequency energy in RF coils for magnetic resonance at 2.4 to 170 MHz. *Magn Reson Med* 1991;21:197-215.
47. Moelker A, Pattynama PM. Acoustic noise concerns in functional magnetic resonance imaging. *Human Brain Mapping* 2003;20:123-41.
48. Miller LE, Keller AM. Regulation of occupational noise. In: Harris CM, ed. *Handbook of noise control*. New York: McGraw-Hill, 1979; pp.1-16.
49. Moelker A, et al. Verbal communication in MR environments: effect of MR system acoustic noise on understanding. *Radiology* 2004;232:107-113.
50. Guyer P J. Fundamentals of acoustics. *CED* 2009; 1-30.
51. Thompson K, Stewart M, Rodriguez J. Biophysics & Bio-imaging. In: *Wave Phenomenon In Hearing (Topic 3)*. University Centenary LA 2008;3:1- 12.
52. Ray FE. Industrial Noise Series: In: *Fundamentals of Environmental Sound* 2010;1:1-16.
53. Laver J. *Acoustic Noise Primer*. 1995; 1.3 Ed.
54. Ho SY, Kveton JF. Acoustic neuroma assessment and management. *Otolaryngol Clin North Am* 2002; 35:393-404.
55. Spoelhof GD. When to suspect acoustic neuroma. *Am Fam Physician* 1995; 52: 1768-74.
56. Asturias C. *The Sound and the Furious*;2001.
57. Mingfeng Li, R Brent, Lim CT, Jing-Huei L. In Situ Active Control of Noise in a 4-Tesla MRI Scanner. *J Magn Reson Imaging* 2011; 34(3): 662-9.
58. McJury M, Shellock FG. Acoustic noise and MR procedures. *Journal of Magnetic Resonance Imaging* 2000; 12: 37-45.
59. Bunder J R, Editor. *Glossary of Nonlethal Weapons Terms*. USAF Academy, Colorado: USAF Institute for National Security Studies, INSS Occasional 1999; Paper 15.
60. Hammerschlag PE, Scheer AA, Fishman A. Revision Stapedectomy: A Review of 308 Cases. *The American Laryngological, Rhinological and Otological Society*. New York. *Laryngoscope* 1998; Vol 108:1794-1800.

61. Berger EH, Franks RJ, Stephenson RM, Merry JC. The Noise Manual. OSHA Technical Manual (OTM) 2003; SecIII: Ch 5.
62. Appler JM, Goodrich LV. Connecting the ear to the brain: Molecular mechanisms of auditory circuit assembly. *Prog Neurobiol* 2011;93(4):488-508.
63. Mårtensson J. Improved analysis of MRI tractography data: Group comparisons of parameters along fibre tracks. *Sciences, Lund University Lund, Sweden* 2010;1-27.
64. Weir J, Abraham PH. MRI. In: *Aslide atlas of imaging of Human Anatomy*. London, Mosby year Book Europe 1994;1st ed.
65. Hart BL, Benzel EC, Ford CC. *Fundamentals of Neuroimaging*. Philadelphia: W. B. Saunders. USA 1997.
66. Schild H. *MRI made easy*. Berlin Schering AG 1990.
67. Brook W C, Kant C. *MRI in Practice UK* 2001, 2nd.
68. Jenkins JPR: Glossary of MRI term. In: Sutton D. *A text book of radiology and imagin*. New York: Churchill Livingstone 1998, 6th ed.
69. Barbara I, Carter. *Computed Tomography of the Head and Neck*. New York, Churchill Livingstone 1985; vol 5.
70. James H. BEATY. MD. American Academy of orthopedic surgery, orthopedic knowledge update. *Journal of the American Medical* 1999; 442.
71. Kuang W, Nelson SO. Low-frequency dielectric dispersion from ion permeability of membranes. *J Colloid. Interface Sci* 1997;193:242-9.
72. Schwan HP. *Electrical Properties of Tissue and Cell Suspensions in Advances in Biological and Medical Physics*. Academic Press 1957; 5: 205 -10.
73. Foster KR, Schwan Ed, Bourne JR. Dielectric properties of tissues and biological material: a critical review, *Biomedical Engineering* 1989;17: 25-104.
74. Foster KR, Joseph D. *Dielectric properties of tissues. The Biomedical engineering Handbook: Second Edition*. Bronzino Boca Raton: CRC press LLC 2000.
75. Luukkonen J, Hakulinen P, Maki-Paakkanen J, Juutilainen J, Naarala J. Enhancement of chemically induced reactive oxygen species production and DNA damage in human SH-SY5Y neuroblastoma cells by 872 MHz radiofrequency radiation. *Mutat Res* 2009; 6621: 54-8.
76. Barry H. Free radicals, antioxidants, and human disease curiosity, cause, or consequence? 1994;1-5.
77. Sarafian TA, Bredesen DE. Is apoptosis mediated by reactive oxygen species? *Free Radic Res* 1994; 20:1-6.
78. Samson J, Sheeladevi R, Ravindran R. Oxidative stress in brain and antioxidant activity of *Ocimum sanctum* in noise exposure. *Neurotoxicology* 2007;28:679-85.

-
79. Sant'anna ID, de Sousa EB, de Moraes AV, Loures DL, Mesquita ET, da Nobrega AC. Cardiac function during mental stress: cholinergic modulation with pyridostigmine in healthy subjects. *Clin Sci (Lond)* 2003;105:161-5.
 80. Liu JK, Mori A. Stress, aging, brain oxidative damage. *Neurochem Res* 1999;1479-97.
 81. Brooker, Robert J. *Genetics: analysis and principles*. McGraw-Hill Science. ISBN 2011; 4th ed.
 82. Henderson D, Bielefeld EC, Harris KC, Hu BH. The role of oxidative stress in noise-induced hearing loss. *Ear & Hearing* 2006; 27 (1):1-19.
 83. Babisch W. Stress hormones in the research on cardiovascular effects of noise. *Noise Health* 2003; 5:1-11.
 84. Ising H, Kruppa B. Health effects caused by noise: Evidence in the literature from the past 25 years. *Noise Health* 2004; 6: 5-13.
 85. Spreng M. Central nervous system activation by noise. *Noise Health* 2000a; 2: 49-58.
 86. Niedernhofer JL, Daniels JS, Rouzer AC, Greene ER, Marnett JL. Malondialdehyde, a Product of Lipid Peroxidation, Is Mutagenic in Human Cells. *The Journal of Biological Chemistry* 2003;278.
 87. Del Rio D, Stewart AJ, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis.* 2005;15:316-28.
 88. Armas-Padilla MC, Armas-Hernández MJ, Sosa-Canache B, Cammarata R, Pacheco B, Guerrero J, et al. Nitric oxide and malondialdehyde in human hypertension. *Am J Ther* 2007;14: 172-6.
 89. Liochev SI, Fridovich I. The role of [O.sub.2] in the production of [HO.sup.*]: in vitro and in vivo. *Free Radic Biol Med* 1994;16:29-33.
 90. Derekoy FS, Dundar Y, Aslan R, Canjal A. "Influence of noise exposure on antioxidant system and TEOAFs in rabbits". *Eur. Arch. Otorhinolaryngol* 2001; 258:518-22.
 91. Antonyuk SV, Seetharaman SV, Whitson LJ, Taylor AB, Holloway SP, Strange RW, et al. "Structures of the G85R variant of SOD1 in familial amyotrophic lateral sclerosis". *J. Biol. Chem* 2008;283(23):16169-77.
 92. Antonyuk SV, Strange RW, Marklund SL, Hasnain SS. "The structure of human extracellular copper-zinc superoxide dismutase at 1.7 Å resolution: insights into heparin and collagen binding". *J. Mol. Biol.* 2009; 388 (2): 310-26.
 93. Li Y, Huang TT, Carlson EJ, Melov S, Ursell PC, Olson JL, et al. "Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase". *Nat. Genet.* 1995;11(4):376-81.

94. Peter H, Löffler G, Petrides EP. Biochemie and Pathobiochemie (Springer-Lehrbuch) (German Edition). Berlin: Springer ISBN 2006;p 123.
95. Ersoy A, Koc ER, Sahin S, Duzgun U , Acar B, Ilhan A. Possible effects of rosuvastatin on noise induced oxidative stress in rat brain. *Noise Health* 2014;16: p18-25.
96. Spreng M. Possible health effects of noise induced cortisol increase. *Noise Health* 2000b; 2: 59–64.
97. Marieb NE. Human anatomy & physiology. 2004; 6thed (97):1946- 53.
98. Goldman L, Schafer AI, eds. Cecil Medicine. Philadelphia, Pa: Saunders Elsevier; 2011;24th ed (161).
99. Dacie JV, Lewis MS. Practical Hematology.S" ed.Churchill Livingstone 1995; 98-118.
100. McPherson RA, Pincus MR. Henry's. Clinical Diagnosis and Management by Laboratory Methods. Pa: WB Saunders, Philadelphia 2007; 21sted: 461-2.
101. Lederbogen F, Baranyai R, Gilles M, Menart-Houtermans B, Tschoepe D, Deuschle M. Effect of mental and physical stress on platelet activation markers in depressed patients and healthy subjects: a pilot study.*Psychiatry Research* 2009;127: 55-64.
102. Roohi N, Irfan S, Bashir T. Haematologica and Hormonal Responses of Traffic Policemen. *Pakistan Journal of Science* 2009; Vol. 61(1):20-25.
103. Medical Research Institute/Ethical Regulation for conducting research on human subjects and animals 2011. (<http://www.apa.org/science/anguide.html>).
104. Chang TY, Liu CS, Young LH, Wang VS, Jian SE, Bao BY.Noise frequency components and the prevalence of hypertension in workers. *Sci Total Environ* 2012; 416: 89-96.
105. Blumenreich MS. The white blood cells and differential count.In: The History, Physical and Laboratory Examination. Walker HK, Hall JW. 3rd ed. Butterworths, Boston.1990;724-7.
106. McMillin JM. Blood glucose. In: The History, Physical and Laboratory Examination. Walker HK, Hall JW (eds). 3rd Ed. Butterworths, Boston. 1990;pp 622-5.
107. Gabriely S, Corthout E. The dielectric properties of biological tissues: I. Literature Survey. *Phys Med Biol* 1996;41:2231-49.
108. Draper HH, Polensek L, Hadley M, Megirr LG. Urinary malondialdehyde as an indicator of lipid peroxidation in the diet and in the tissues. *Lipids* 1984;19: 836-43.
109. Sun YI, Oberley LW, Ying I.A simple method for clinical assay of superoxide dismutase. *Clin Chem* 1988; 34:497-500.
110. Yumita N ,Nishigaki R, Umemurea K ,Umemura S. Synergistic effect of ultra sound and hemato porphyrin on sarcoma 180. *J Pn J cancer Res.*1990; 81: 304-8.

111. Asamiki S, Takashiy H, Takashima S. Dielectric properties of mouse lymphocytes and erythrocytes. *Biochemical et Biophysics Acta*. 1988;1010:49-55.
112. Burtis C, Ashwood ER, Bruns DE. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics*. 4th ed. Elsevier Saunders Company, St Louis. 2006;869-72.
113. Draper HH, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol*. 1990; 186:421-5.
114. Marklund S, Marklund G. Involvement of superoxide anion radical in autoxidation of pyrogallol and a convenient assay for SOD. *Eur. J. Biochem* 1974;47:469-74.
115. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurements with Folin-phenol reagent. *J. Biol. Chem.* 1951;193:265-70.
116. Roper D, Layton M, Lewis SM. Investigation of the hereditary haemolytic anaemias: membrane and enzyme abnormalities. In: Lewis SM, Bain BJ, Bates I eds. *Dacie and Lewis practical haematology*. 4th ed. Harcourt Publisher Limited, London, UK 2001; 10: 167-98
117. Goldman AM, Gossman WE, Friedlander PC. Reduction of sound levels with antinnoise in MR imaging. *Radiology* 1989;173:549-50.
118. Shellock FG, Morisoli SM, Ziarati M. Measurement of acoustic noise during MR imaging: evaluation of six "worst-case" pulse sequences. *Radiology* 1994;191:91-93.
119. McJury M, Blug A, Joerger C, Condon B, Wyper D. Acoustic noise levels during magnetic resonance imaging scanning at 1.5 T. *Br J Radiol* 1994;67:413-15.
120. McJury MJ. Acoustic noise levels generated during high field MR imaging. *Clin Radiol* 1995;50:331-34.
121. Counter SA, Olofsson A, Grahn, Borg E. MRI acoustic noise: sound pressure and frequency analysis. *J Magn Reson Imaging* 1997;7: 606-11.
122. Cho ZH, Park SH, Kim JH, Chung SC, Chung ST, Chung JY et al. Analysis of acoustic noise in MRI. *Magn Reson Imaging* 1997;15: 815- 22.
123. Glover P, Hykin J, Gowland P, Wright J, Johnson I, Mansfield P. An assessment of the intrauterine sound intensity level during obstetric echo-planar magnetic resonance imaging. *Br J Radiol* 1995; 68: 1090-94.
124. Karakoc Y, Orum M H, Kilic M, Dursun M F, Erdogan S. Effects of 85 dB Noise Exposure on Macro and Trace Element Levels in Plasma and Brain Areas of Rats. *Biol Trace Elem Res* 2013; 154:134-39
125. Abumaria N, Yin B, Zhang L, Li XY, Chen T, Descalzi G, et al. Effects of elevation of brain magnesium on fear conditioning, fear extinction, and synaptic plasticity in the infralimbic prefrontal cortex and lateral amygdala. *J Neurosci* 2011;31(42):14871- 81

126. Dempsy MF, Condoa B. Thermal injuries associated with MRI. *Clini Radio* 2001; 56:457- 65.
127. Knopp MV, Metzsnor R, Brix G, van Kaick G. Safety considerations to avoid current-induced skin burns in MRI procedure. (German) *Radiologic* 1998; 38: 759- 63.
128. Aslan M, Kosecik M, Horoz M, Selek S, Celik H, Erel O. Assessment of paraoxonase and arylesterase activities in patients with iron deficiency anemia. *Atherosclerosis*. 2007;191:397-402.
129. Altindag O, Gur A, Calgan N, Soran N, Celik H, Selek S. Paraoxonase and arylesterase activities in fibromyalgia. *Redox Rep*. 2007;12:134- 8.
130. Le Baron FN. Characteristics, metabolism and function role of polyunsaturated fatty acids. In: Devlin TM. *Textbook of Biochemistry*. New York: Wiley's Medical Publication, 1982:479- 85.
131. Liewei S, Duke U, Durham NC. A Review of Dielectric Properties of Normal and Malignant Breast Tissue. *IEEE Southeastcon*. 2002;3:451-6.
132. Joines W, Dhenxing Y, Jirtle R. The measured electrical properties of normal and malignant human tissues from 50 to 900 MHz. *Med. Phys*. 1994;21:547-50.
133. Kanitz E, Otten W, Tuchscherer M. Central and peripheral effects of repeated noise stress on hypothalamic- pituitary-adrenocortical axis in pigs. *Livest Prod Sci* 2005; 94: 213- 24.
134. De La Fuente J, Díaz MT, Ibáñez M, De Chavarri EG. Physiological response of rabbits to heat, cold, noise and mixing in the context of transport. *Anim Welf* 2007; 16:41-47
135. Rewerska A, Pawelczyk M, Rajkowska E, Politanski P, Sliwiska-Kowalska M. Evaluating D-methionine dose to attenuate oxidative stress-mediated hearing loss following overexposure to noise. *Eur Arch Otorhinolaryngol* 2013;270:1513- 20
136. Marita G K, Kandregula S, Lakshmana N R. Acute noise exposure effect on certain hematological parameters in tailors: a pilot study. *Int J Biol Med Res*. 2012;3(2): 1529- 31.
137. Cheng L, Wang SH, Chen QC, Liao XM. Moderate noise induced cognition impairment of mice and its underlying mechanisms. *Physiol Behav* 2011;104:981-8.
138. Babisch W, Fromme H, Beyer A, Ising H. Increased catecholamine levels in urine in subjects exposed to road traffic noise: The role of stress hormones in noise research. *Environ Int* 2001;26:475-81.
139. Selander J, Bluhm G, Theorell T, Pershagen G, Babisch W, Seiffert I, et al. Saliva cortisol and exposure to aircraft noise in six european countries. *Environ Health Perspect* 2009a;117:1713-17
140. Concha Barrientos M, Campbell-Lendrum D, Steenland K. Occupational Noise. *Environmental Burden of Disease Series, No. 9*. World Health Organization. Protection of the Human Environment. Geneva. WHO 2004.

141. Sembulingam K., Sembulingam P, Namasivayam A. Effect of acute noise stress on acetylcholinesterase activity in discrete areas of rat brain . *Indian Journal of Medical Sciences* 2003;57(11):487-92.
142. Kaltenbach JA, Czaja JM, Kaplan CR. Changes in the tonotopic map of the dorsal cochlear nucleus following induction of cochlear lesions by exposure to intense sound. *Hear Res* 1992;59:213-23.
143. Scarfiotti C, Fabris F, Cestaro B, Giuliani A. Free radicals, atherosclerosis, ageing, and related dysmetabolic pathologies: pathological and clinical aspects. *Eur J Cancer Prev* 1997;6:31-6.
144. Manikandan S, Devi RS. Antioxidant property of alpha-asarone against noise-stress-induced changes in different regions of rat brain. *Pharmacol Res* 2005; 52:467-74.
145. Manikandan S, Srikumar R, Jeya Parthasarathy N, Sheela Devi R. Protective effect of *Acorus calamus* LINN on free radical scavengers and lipid peroxidation in discrete regions of brain against noise stress exposed rat. *Biol Pharm Bull* 2005;28: 2327-30.
146. Hammadi S, Ajlan S, Shindi A, Yousof A. During Magnetic Assault Oxidative Examination Imaging Resonance. *Bas J Surg* 2011; 49-53.
147. Aslan M, Sabuncu T, Kocyigit A, Celik H, Selek S. Relationship between total oxidant status and severity of diabetic nephropathy in type 2 diabetic patients. *Nutr Metab Cardiovasc Dis.* 2007;17:734- 40.
148. Tas F, Hansel H, Belce A, Ilvan S, Argon A, Camlica H, et al. Oxidative stress in breast cancer. *Med Oncol* 2005;22:11-5.
149. Manfredini V, Lazzaretti LL, Griebeler IH, Santin AP, Brandao VD, Wagner S, et al. Blood antioxidant parameters in sickle cell anemia patients in steady state. *J Natl Med Assoc* 2008;100:897-902.
150. Minuz P, Patrignani P, Gaino S, Degan M, Menapace L, Tommasoli R, et al. Increased Oxidative Stress and Platelet Activation in Patients With Hypertension and Renovascular Disease. *Circulation.* 2002;106:2800- 05.
151. Surapaneni KM, Venkataramana G. Status of lipid peroxidation, glutathione, ascorbic acid, vitamin E and antioxidant enzymes in patients with osteoarthritis. *Indian J Med Sci.* 2007;61:9-14.
152. Aslan M, Horoz M, Nazligul Y, Bolukbas C., Bolukbas F.F., Selek S, et al. Serum paraoxonase and arylesterase activities for the evaluation of patients with chronic hepatitis. *Int J Clin Pract* 2008;62:1050-55.
153. Srikumar R, Parthasarathy NJ, Manikandan S, Narayanan GS, Sheeladevi R. Effect of Triphala on oxidative stress and on cell-mediated immune response against noise stress in rats. *Mol Cell Biochem* 2006;283:67-74.
154. Ilhan A, Gurel A, Armutcu F, Kamisli S, Iraz M, Akyol O, et al. Ginkgo biloba prevents mobile phone-induced oxidative stress in rat brain. *Clin Chim Acta* 2004;340:153-62.

155. Fetoni AR, De Bartolo P, Eramo SL, Rolesi R, Paciello F, Bergamini C, et al. Noise-induced hearing loss (NIHL) as a target of oxidative stress-mediated damage: cochlear and cortical responses after an increase in antioxidant defense. *J Neurosci* 2013;33(9):4011-23.
156. Naziroglu M. TRPM2 cation channels, oxidative stress and neurological diseases: Where are we now? *Neurochem Res* 2011; 36:355-66.
157. Karakoc Y, Turhan S, Yildirim EA, Mengi M, Yurdakos E. Neuropeptide Y alters stress-induced changes in trace element concentrations of brain in chronically immobilized rats. *J Trace Elem Exp Med* 2004;17(4):283-90.
158. Naziroglu M. Molecular role of catalase on oxidative stress-induced Ca²⁺ signaling and TRP cation channel activation in nervous system. *J Recept Signal Transduct* 2012;32(3):134-41.
159. Sørensen M, Andersen ZJ, Nordsborg RB, Becker T, Tjønneland A, Overvad K, et al. Long-term exposure to road traffic noise and incident diabetes: A cohort study. *Environ Health Perspect* 2013;121:217-22.
160. Samson J, Wiktorek-Smagur A, Politanski P, Rajkowska E, Pawlaczyk-Luszczynska M, Dudarewicz A, Sha S-H, Schacht J, Sliwinska-Kowalska M. Noise-induced time dependent changes in oxidative stress in the mouse cochlea and attenuation by D-methionine. *Neuroscience* 2008;152:146-150.
161. Mcfadden SL, Ohmiller KK, Ding D, Shero M, Salvi RJ. The influence of superoxide dismutase and glutathione peroxidase deficiencies on noise-induced hearing loss in mice. *Noise Health* 2001;331:49-64.
162. Emily G, Mark J. Noise exposure, music, and animals in the Laboratory. A commentary based on laboratory animal refinement and enrichment forum (LAREF) discussions. *Applied Animals Welfare Science Journal* 2006;9(4):327-32.
163. Marth E, Gallasch E, Fueger GF, Mose JR. Aircraft noise: Changes in biochemical parameters. Source: *Zentralbl Bakteriol Mikrobiol Hyg B* 1988;185(4-5):498-508.
164. Kight RC, Swaddle PJ. *Ecology Letters*. 2011;14:1052-61.
165. Zardooz H, Zahedi S, Gharib NM K. Effect of chronic psychological stress on insulin release from rat isolated pancreatic islets. *Life Sciences* 2005;1(79):57-62.
166. Van de Kar LD, Blair ML. Forebrain pathways mediating stress induced hormone secretion. *Front Neuroendocrinol*. 1999;20(1):1-48.
167. Yamada F, Inoue S, Saitoh T, Tanaka K, Satoh S, Takamura Y. Glucoregulatory hormones in the immobilization stress induced increase of plasma glucose in fasted and fed rats. *Endocrinology*. 1993;132(5):2199-205.
168. Teague CR, Dhabhar FS, Barton RH, Beckwith-Hall B, Powell J, Cobain M, Singer B, McEwen BS, Lindon JC, Nicholson JK, et al. Metabonomic studies on the physiological effects of acute and chronic psychological stress in Sprague-Dawley rats. *J Proteome Res*. 2007;6(6):2080-93.

169. Qureshi F, Alam J, Ahmad M, Sheraz G. Effect of examination stress on blood cell parameters of students in a Pakistani Medical College. 1. *Ayub Med. Coil.* 2002;14:20-22.
170. Maes M, Planken V M, Gastel VA, Spanaki VM. Influence of academic stress on hematological measurements in subjectively healthy volunteers. *Psychiatry Res.* 1998;80:201-12.
171. Baum A, Grunberg G. Measurement of stress hormones. In *Measuring stress.* Cohen S, Kessler RC, Gordon L, eds. Oxford University Press, New York, 1995; pp175-192.
172. Boikovi: Hygienic characteristics of working conditions and health status of workers engaged in the production of benzene and xylenes at a petroleum refinery. *Gig Tr Prof Zabol* 1970;14:23-5.
173. Kapaki EN, Varelas NP, Syrigou AN, Spanaki MV, Andreadou E, Kakami EA, Papageorgiou CT. Blood Lead Levels of Traffic and Gasoline-exposed professionals in the city of Athens. *Archives of Environmental Health* 1998; 53: 287-91
174. Sinisalo J, Paronen J, Mattila KJ, Syrjala M, Alfthan G, Palosuo T et al. Relation of inflammation to vascular function in patients with coronary heart disease. *Atherosclerosis* 2000;149:403-11.
175. Zheng KCh, Ariizumi M. Modulations of Immune functions and oxidative status induced by noise stress. *J Occup Health* 2007;49:32-8.
176. Hioshi K, Kanehisa M. Effects of mental arithmetic stress on blood cell counts and immune system. *Environ Health and Prevent Med.* 1996;(1) 2:76- 9.
177. Sabahi A.R, Moradi I. The Effects of Noise Exposure on Rat's Hematologic Parameters and Red Cell Indices. *Iran J Med Sci* 2002;27(2):85- 6.
178. Aguas AP, Esaguy N, Grande N, Castro AP, Castelo Branco NA. Effect low frequency noise exposure on BALB/c mice splenic lymphocytes, *Aviat Space Environ Med.* 1999;70:31-128.
179. Hoffman R, Benz EJ, Shattil SJ, Furie B, Cohen HJ, eds. *Hematology: Basic Principles and Practice.* 4th ed. Philadelphia, Pa: Churchill Livingstone 2005;2676-77.
180. Nakanishi N, Yoshida H, Okamoto M, Matsuo Y, Suzuki K, Tatara K Association of alcohol consumption with white blood cell count: a study of Japanese male office workers. *J Intern Med* 2003;253(3):367-74.
181. Jern C, Wadenvik H, Mark H, Hallgren J, Jern S. Haematological changes during acute mental stress. *Br J Haematol.* 1989;71:153-6.
182. Brydon L, Magid K, Steptoe A. Platelets, coronary heart disease, and stress. *Brain Behav Immun.* 2006;20:113-19.

183. Patterson SM, Gottdiener JS, Hecht G, Vargot S, Krantz DS. Effects of acute mental stress on serum lipids: Mediating effects of plasma volume. *Psychosom Med.* 1993;55:525-32.
184. Koulunis M, Liu Y, Hallstrom K, Socolovsky M. Negative Autoregulation by Fas Stabilizes Adult Erythropoiesis and Accelerates Its Stress Response. *PLoS One* 2011;6(7):e21192.
185. Pandey KB, Rizvi SI. Biomarkers of Oxidative Stress in Red Blood Cells. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub. 2011;1-5.

الملخص العربي

التصوير بالرنين المغناطيسي من الوسائل التشخيصية الهامة ولكن أثناء التصوير يتعرض المريض لضوضاء صادرة من جهاز الرنين المغناطيسي ناتجة من التغير الشديد والسريع في المجال المغناطيسي للجهاز وهذا قد يؤدي بدوره إلى أضرار صحية للمريض.

• الضوضاء: تعرف الضوضاء بأنها خليط متناثر من الأصوات ذات استمرارية غير مرغوب فيها وتشكل خطر كبير علي صحة الإنسان. يمكن قياس الضوضاء بطرق فيزيائية يُعبر عنها بالديسيبل. وقد وجد أن التعرض إلي ضوضاء بمستوي ٨٥ ديسيبل أو أكثر لفترة قد يؤدي إلي فقدان السمع الدائم. هناك الكثير من التأثيرات الصحية الناجمة عن تلك الضوضاء منها التأثيرات السمعية المباشرة ويندرج تحتها ضعف السمع أو فقدانه بشكل كامل ومنها التأثيرات الغير سمعية مثل ارتفاع ضغط الدم، امراض الجهاز الدوري، اضطرابات النوم و الاضطرابات النفسية والعصبية.

تأثيرات الضوضاء السمعية التي تتولد من جهاز الرنين المغناطيسي خلال فترة التصوير الغير مرغوبة لها تأثيرات جانبية هي تأثيرات صحية مضادة تصنف دائما طبقا لنوع الضوضاء. يعمل على سهولة التأثير بهذه الزيادة من مستوى الضوضاء إلى الأمراض وعوامل مضاعفة يعمل مشاكل في القلب وأمراض أخرى. وبيان بان هناك مؤشرات معقولة بارتباط عادي بين التعرض للضوضاء وتأثيرها على الصحة كما موجود في دراستنا.

الإعتبرات الأساسية:

تم عرض الأسس الفيزيائية للصوت وكيفية قياس الضوضاء ومدى تأثيره على الأذن البشرية وعرض أيضا الأسس الفيزيائية لجهاز الرنين المغناطيسي ومصدر الضوضاء لهذا الجهاز. كما تم توضيح خاصية العزل الكهربائي للأنسجة ومدى تأثيرها بالضوضاء.

هدف الرسالة:

تهدف هذه الدراسة إلى بيان التأثيرات الغير سمعية للصوت الصادر أثناء التصوير بالرنين المغناطيسي على بعض الخواص الفيزيائية (العزل الكهربائي) لنسيج مخ الفأر وبعض التغيرات الكيميائية الحيوية مثل نسبة السكر في الدم، ومستوى بعض الأنزيمات المضادة للأكسدة مثل superoxide dismutase، malondialdehyde في مخ الفئران. وكشف الارتباط بين شدة الصوت الصادر أثناء التصوير بالرنين المغناطيسي وهذه المتغيرات في دم ومخ الفئران.

مادة البحث والطرق المستخدمة في البحث:

تم تسجيل الصوت الفعلي الصادر من جهاز الرنين المغناطيسي (من النوع المغلق 1.5T) أثناء التصوير لمدة 27 دقيقة بواسطة مسجل متصل بجهاز كمبيوتر محمول ثم نقل مصدر الصوت على ناقل ذاكرة وتوصيلها بمكبر للصوت متصل بسماعات عالية القدرة. تم ضبط مصدر الصوت على مسافة عشرة سنتيمترات من مركز السماعة لتكون قوة الضوضاء ٩٠ أو ١٠٠ أو ١٢١ ديسيبل باستعمال جهاز قياس مستوى الصوت (Sound level Meter). تم استخدام مجموعة من برامج الكمبيوتر لتحليل مصادر الضوضاء والتعرف على التردد الرئيسي

تم تصميم وتصنيع غرف معزولة صوتيا من الخشب مبطنه بعوازل للصوت والمراعى فيها توصيل مصادر للإضاءة والتهوية مع مراعاة منع تسريب الصوت منها أو إليها وتم إجراء التعريض لكل المجموعات داخل هذه الغرف.

في هذه الرسالة تم استخدام ٤٠ فأر تم تقسيمها إلي أربع مجموعات كالاتي

- المجموعة الأولى: المجموعة الضابطة. تم وضعهم في حجرة التعرض لمدة ساعتان يوميا لمدة أسبوع دون ضوضاء.
- المجموعة الثانية: يتم وضعهم في حجرة التعرض حيث يتعرض الفئران في هذه المجموعة إلى ضوضاء شدتها ١٢١ ديسيبل لمدة ساعتان يوميا لمدة أسبوع.

- المجموعة الثالثة: يتم وضعهم في حجرة التعرض حيث يتعرض الفئران في هذه المجموعة إلى ضوء شدتها ١٠٠ ديسيبيل لمدة ساعتان يوميا لمدة أسبوع.
 - المجموعة الرابعة: يتم وضعهم في حجرة التعرض حيث يتعرض الفئران في هذه المجموعة إلى ضوء شدتها ٩٠ ديسيبيل لمدة ساعتان يوميا لمدة أسبوع.
- وبعد انتهاء فترة التعريض تم إجراء الخطوات الآتية:
- تجميع عينات من دم الفئران لدراسة كل من :
 - ١- نسبة الجلوكوز بالدم كيميائيا.
 - ٢- العد الكلي لكرات الدم البيضاء والحمراء وكذلك الصفائح الدموية بصورة رقمية.
 - ذبح الفئران وأخذ عينات من المخ لعمل الآتي:
 - ١- استخدام جزء من مخ الفئران وتم تحضيرها لقياس:
 - أ. إنزيم السوبر أكسيد ديسميوتيز (Superoxide dismutase) وذلك لمعرفة مدى تكوين الأصول الحرة في جسم الفئران المعرضة.
 - ب. ماده المالوندايالاهايد (Malondialdhd) والتي تعطى انطبعا مباشرة عن مدى أكسدة الدهون المكونة لخلايا المخ.
 - ٢- الجزء الآخر من المخ لدراسة العزل الكهربائي.

النتائج:

- حدوث ارتفاع ملحوظ لنسبة الجلوكوز في دم الفئران المعرضة عن تلك التي لم تتعرض نهائيا للضوء.
- حدوث ارتفاع ملحوظ للعد الكلي لكرات الدم البيضاء و الصفائح الدموية، وانخفاض ملحوظ في عدد كرات الدم الحمراء لجميع الفئران المعرضة للضوء.
- ارتفاع ملحوظ لمستوى تركيز المالوندايالاهايد للفئران المعرضة عن تلك التي لم تتعرض نهائيا للضوء.
- حدوث انخفاض ملحوظ لنسبة إنزيم السوبر أكسيد ديسميوتيز للمجموعات التي تعرضت للضوء عن تلك التي لم تتعرض نهائيا للضوء.
- تغير في خواص العزل الكهربائي لنسيج المخ للمجموعات التي تعرضت للضوء عن تلك التي لم تتعرض نهائيا للضوء.

مناقشة النتائج:

تفسيرا لهذه النتائج تمت مقارنة النتائج التي حصلنا عليها مع دراسات وأبحاث علمية سابقة تتعلق بنفس الموضوع، كما تم ذكر بعض الأبحاث التي أظهرت نتائج مشابهة. وقد عرضت المناقشة بأسلوب علمي سليم و تعد مناقشة وافية لجميع أوجه البحث ونتائجه.

الاستنتاجات:

- نستنتج من النتائج السابقة ان الضوء الصادرة من جهاز الرنين المغناطيسي أثناء التشخيص لها تأثير ضار ومباشر في المدى الصوتي تمت الدراسة علي:
- ١- نسبة الجلوكوز بالدم مما قد يعرض الكائن الحي للإصابة بمرض السكري.
 - ٢- العد الكلي لكرات الدم البيضاء يتأثر مما يؤدي إلى اضطرابات في خط الدفاع الداخلي للجسم وهو المناعة.
 - ٣- نسبة تواجد المواد المؤكسدة بالجسم.
 - ٤- أثبتت الدراسة أن هناك تغيرات في خاصية العزل الكهربائي لأنسجة المخ مما يدل على مدى تأثير الضوء الناتجة من جهاز الرنين المغناطيسي المستخدم على الكثير من العمليات الحيوية بالجسم.

دراسة بعض التأثيرات البيوفيزيائية والبيوكيميائية للصوت الناتج من جهاز الرنين
المغناطيسي للتصوير: باستخدام الفرن كنموذج تجريبي

رسالة

مقدمة إلى معهد البحوث الطبية- جامعة الإسكندرية
إيفاء جزئيا لشروط الحصول على درجة

الماجستير

فى

الفيزياء الحيوية الطبية

مقدمة من

رمزية مصطفى شاهين

بكالوريوس تكنولوجيا طبية فى تقنيات الأشعة
جامعة بغداد، ٢٠٠٢

معهد البحوث الطبية
جامعة الإسكندرية
٢٠١٤

دراسة بعض التأثيرات البيوفيزيائية والبيوكيميائية للصوت الناتج من جهاز الرنين
المغناطيسي للتصوير: باستخدام الفرن كنموذج تجريبي

مقدمة من

رمزية مصطفى شاهين

بكالوريوس تكنولوجيا طبية فى تقنيات الأشعة

جامعة بغداد

٢٠٠٢

للحصول على درجة

الماجستير

فى

الفيزياء الحيوية الطبية

موافقون

لجنة المناقشة والحكم على الرسالة

.....

أ.د. طارق عثمان النمر
أستاذ متفرغ قسم الفيزياء الحيوية
كلية العلوم
جامعة طنطا

.....

أ.د. يوسف سليم يوسف
أستاذ متفرغ قسم الفيزياء الحيوية
معهد البحوث الطبية
جامعة الإسكندرية

.....

أ.د. ثناء إبراهيم شلبي
أستاذ قسم الفيزياء الحيوية
معهد البحوث الطبية
جامعة الإسكندرية

.....

أ.د. سهير محمود الخولى
أستاذ قسم الفيزياء الحيوية
معهد البحوث الطبية
جامعة الإسكندرية

السادة المشرفون

.....

أ.د/ يوسف سليم يوسف

أستاذ متفرغ قسم الفيزياء الحيوية

معهد البحوث الطبية

جامعة الإسكندرية

.....

أ.د/ ثناء إبراهيم شلبي

أستاذ قسم الفيزياء الحيوية

معهد البحوث الطبية

جامعة الإسكندرية

.....

د/ بثينة فؤاد محمود

زميل بقسم الكيمياء الحيوية

معهد البحوث الطبية

جامعة الإسكندرية