

RECOMMENDATIONS

From this study it was recommended that:

Central venous to arterial carbon dioxide gradient may be considered in any futural alterations in the protocol of detecting postoperative complications and mortality in patients undergoing cardiac surgery under cardiopulmonary bypass, putting in mind its simplicity and easy applicability.

LIMITATIONS

Before interpreting ΔPCO_2 , some precautions have to be taken. Numerous potential causes of error of measurements must be eliminated: incorrect sample container, contaminated sample by air or venous blood or catheter fluid.

REFERENCES

1. Rees K, Bearnek-stanley M, Burke M, Ebrahim S. Hypothermia to reduce neurological damage following coronary artery bypass surgery. *Cochrane Database syst Rev* 2001; 1:2138-50.
2. Murphy GJ, Reeves BC, Rogers CA, Rizvi SI, Culliford L, Angelini GD. Increased mortality, postoperative morbidity, and cost after red blood cell transfusion in patients having cardiac surgery. *Circulation* 2007; 116:2544-52.
3. Rossaint J1, Berger C, Van Aken H, Scheld HH, Zahn PK, Rukosujew A, Zarbock A. Cardiopulmonary bypass during cardiac surgery modulates systemic inflammation by affecting different steps of the leukocyte recruitment cascade. *PLoS One* 2012; 7(9):45738-49.
4. Biagioli B1, Borrelli E, Maccherini M, Bellomo G, Lisi G, Giomarelli P, Sani G, Toscano M. Reduction of oxidative stress does not affect recovery of myocardial function: warm continuous versus cold intermittent blood cardioplegia. *Heart Surg* 1997; 77(5):465-73.
5. Salhiyyah K1, Elsobky S, Raja S, Attia R, Brazier J, Cooper GJ. A clinical and economic evaluation of fast-track recovery after cardiac surgery. *Heart Surg Forum* 2011; 14(6):330-4.
6. Kuiper AN, Trof RJ, Groeneveld AB. Mixed venous O₂ saturation and fluid responsiveness after cardiac or major vascular surgery. *Cardiothorac surg* 2013; 8:189-97.
7. Caille V, Squara P. Oxygen uptake-to-delivery relationship: a way to assess adequate flow. *Crit Care* 2006; 10:54-5.
8. Van Beest PA, Brander L, Jansen SP, Rommes JH, Kuiper MA, Spronk PE. Cumulative lactate. *Ann Intensive Care* 2013; 6:3-6.
9. Gersak B, Sutlic Z. Aortic and mitral valve surgery on the beating heart is lowering cardiopulmonary bypass and aortic cross clamp time. *Heart Surg Forum* 2002; 5(2):182-6.
10. 10 - Laver RD1, Wiersema UF1, Bersten AD. Echocardiographic estimation of mean pulmonary artery pressure in critically ill patients. *Crit Care Ultrasound J* 2014; 6(1):9-20.
11. Cioccarl L, Baur HR, Berger D, Wiegand J, Takala J, Merz TM. Hemodynamic assessment of critically ill patients using a miniaturized transesophageal echocardiography probe. *Crit Care* 2013; 17(3):121-31.
12. Imren Y, Tasoglu I, Oktar GL, et al. The importance of transesophageal echocardiography in diagnosis of pericardial tamponade after cardiac surgery. *J Card Surg* 2008; 23:450-8.

References

13. McKenney PA, Apstein CS, Mendes LA, et al. Increased left ventricular diastolic chamber stiffness immediately after coronary artery bypass surgery. *J Am Coll Cardiol* 1994; 24:1189-98.
14. Hottinger DG, Beeb DS, Kozhimannil T, Prielipp RC et al. Sodium nitroprusside in 2014: A clinical concepts review. *J Anaesthesiol Clin Pharmacol* 2014; 30(4):462-71.
15. Varon JI, Marik PE. Perioperative hypertension management. *Vasc Health Risk Manag* 2008; 4(3):615-27.
16. Rivera A, Montoya E, Varon J. Intravenous clevidipine for management of hypertension. *Integr Blood pressure Control* 2010; 3:105-11.
17. Kuttilla K, Niinikoski J. Peripheral perfusion after cardiac surgery. *Crit Care Med* 1989; 17:217-24.
18. Denault A1, Deschamps A, Tardif JC, Lambert J, Perrault L. Pulmonary hypertension in cardiac surgery. *Curr Cardiol Rev* 2010; 6(1):1-14.
19. Christakis GT, Fremes SE, Koch JP, et al. Determinants of low systemic vascular resistance during cardiopulmonary bypass. *Ann Thorac Surg* 1994; 58(4):1040-9.
20. Rees K, Baranek-Stanley M, Burke M, Ebrahim S. Hypothermia to reduce neurological damage following coronary artery bypass surgery. *Cochrane Database Syst Rev* 2001; (1):CD002138.
21. Kobayashi M1, Massiello A, Karimov JH, Van Wagoner DR, Fukamachi K. Cardiac autonomic nerve stimulation in the treatment of heart failure. *Ann Thorac Surg* 2013; 96(1):339-45.
22. De Baker D, Creteur J, Silva E, Vincent JL. Effects of dopamine, norepinephrine, and epinephrine on the splanchnic circulation in septic shock: which is best?. *Crit Care Med* 2003; 31(6):1659-67.
23. Amsallem E, Kasparian C, Haddour G, Boissel JP. Phosphodiesterase III inhibitors for heart failure. *Cochrane Database Syst Rev* 2005; 1:CD002230.
24. Morgan BP, Muci A, Lu PP, et al. Discovery of omecamtiv mecarbil the first, selective, small molecule activator of cardiac myosin. *ACS Med Chem Lett* 2010; 1(19):472-7.
25. Haghjoo M, Heidarali M, Nikfarjam S, et al. Very late effects of postoperative atrial fibrillation on outcome of coronary artery bypass graft surgery. *Res Cardiovasc Med* 2012; 1(1):23-7.
26. Regenbogen SE1, Lancaster RT, Lipsitz SR, Greenberg CC, Hutter MM, Gawande AA. Does the Surgical Apgar Score measure intraoperative performance. *Ann Surg* 2008; 248(2):320-8.

References

27. Yokoyama Y, Chaitman BR, Hardison RM, et al. Association between new electrocardiographic abnormalities after coronary revascularization and five-year cardiac mortality in BARI randomized and registry patients. *Am J Cardiol* 2000; 86(8):819-24.
28. Muehlschlegel JD, Perry TE, Liu KY, Nascimben L, Fox AA, Collard CD, Avery EG, Aranki SF, D'Ambra MN, Shernan SK, Body SC. CABG Genomics Investigators. Troponin is superior to electrocardiogram and creatinine kinase MB for predicting clinically significant myocardial injury after coronary artery bypass grafting. *Eur Heart J* 2009; 30(13):1574-83.
29. Jimenez Rivera JJ, Iribarren JL, Raya JM, Nassar I, Lorente L, Perez R, Brouard M, Lorenzo JM, Garrido P, Barrios Y, Diaz M, Alarco B, Martinez R, Mora ML. Factors associated with excessive bleeding in cardiopulmonary bypass patients: a nested case-control study. *J Cardiothorac Surg* 2007; 10(2):17-8.
30. Ballotta A, Saleh HZ, El Baghdady HW, et al. Comparison of early platelet activation in patients undergoing on-pump versus off-pump coronary artery bypass surgery. *J Thorac Cardiovasc Surg* 2007; 134(1):132-8. .
31. Ker K, Edwards P, Perel P, Shakur H, Roberts I. Effect of tranexamic acid on surgical bleeding: systematic review and cumulative meta-analysis. *BMJ* 2012; 344:e3054.
32. Henry D1, Carless P, Fergusson D, Laupacis A. The safety of aprotinin and lysine-derived antifibrinolytic drugs in cardiac surgery: a meta-analysis. *CMAJ* 2009; 180(2):183-93.
33. Bednar F, Osmancik P, Vanek T, et al. Platelet activity and aspirin efficacy after off-pump compared with on-pump coronary artery bypass surgery: results from the prospective randomized trial PRAGUE 11-Coronary Artery Bypass and Reactivity of Thrombocytes (CABARET). *J Thorac Cardiovasc Surg* 2008; 136(4):1054-60.
34. Zimmermann N, Kurt M, Wenk A, et al. Is cardiopulmonary bypass a reason for aspirin resistance after coronary artery bypass grafting? *Eur J Cardiothorac Surg* 2005; 27(4):606-10.
35. Zimmermann N, Gams E, Hohlfeld T. Aspirin in coronary artery bypass surgery: new aspects of and alternatives for an old antithrombotic agent. *Eur J Cardiothorac Surg* 2008; 34(1):93-108.
36. Oz BS, Iyem H, Akay HT, Yildirim V, Karabacak K, Bolcal C, Demirkiliç U, Tatar H. Preservation of pleural integrity during coronary artery bypass surgery affects respiratory functions and postoperative pain: a prospective study. *Can Respir J* 2006; 13(3):145-9.
37. Machuca TN, Collaud S, Mercier O, et al. Outcomes of intraoperative extracorporeal membrane oxygenation versus cardiopulmonary bypass for lung transplantation. *J Thorac Cardiovasc Surg* 2014; 523(14):1819-24.

References

38. Ramsey J. The Respiratory, renal, and hepatic systems: Effects of cardiac surgery and cardiopulmonary bypass. In: *Cardiopulmonary Bypass*, Mora, CT (Eds), Springer, New York 1995; 5:147-52.
39. Engoren M, Buderer NF, Zacharias A. Long-term survival and health status after prolonged mechanical ventilation after cardiac surgery. *Crit Care Med* 2000; 28:2742-9.
40. Vidal Melo MF, Musch G, Kaczka DW. Pulmonary pathophysiology and lung mechanics in anesthesiology: a case-based overview. *Anesthesiol Clin* 2012; 30(4):759-84.
41. Trachte AL, Lobato EB, Urdaneta F, et al. Oral sildenafil reduces pulmonary hypertension after cardiac surgery. *Ann Thorac Surg* 2005; 79:194-201.
42. Lee JE, Hillier SC, Knoderer CA. Use of sildenafil to facilitate weaning from inhaled nitric oxide in children with pulmonary hypertension following surgery for congenital heart disease. *J Intensive Care Med* 2008; 23(5):329-34.
43. Di Tommaso L1, Iannelli G, Monaco M, Mottola M, De Amicis V, Spampinato N. A single-stage procedure for carotid endarterectomy and myocardial revascularization: early and late results. *Tex Heart Inst J* 2005; 32(3):271-6.
44. Smulter N, Lingehal HC, Gustafson Y, Et al. Delirium after cardiac surgery: incidence and risk factors. *Interact Cardiovasc Thorac Surg* 2013; 17(5):790-6.
45. Walzer T, Herrmann M, Wallesch CW. Neuropsychological disorders after coronary bypass surgery. *J Neurol Neurosurg Psychiatry* 1997; 62(6):644-8.
46. Cartin-Ceba R, Kashiouris M, Plataki M, Kor DJ, Gajic O, Casey ET. Risk factors for development of acute kidney injury in critically ill patients: a systematic review and meta-analysis of observational studies. *Crit Care Res Pract.* 2012; 6(9):101-3.
47. Safaie N, Chaichi P, Habibzadeh A, Nasiri B. Postoperative outcomes in patients with chronic renal failure undergoing coronary artery bypass grafting in Madani heart center: 2000-2010. *J Cardiovasc Thorac Res* 2011; 3 (2):53-6.
48. Kinsey GR, Okusa MD. Pathogenesis of acute kidney injury: foundation for clinical practice. *Am J Kidney Dis* 2011; 58(2):291-301.
49. Kuiper AN, Trof RJ, Groeneveld AB. Mixed venous O₂ saturation and fluid responsiveness after cardiac or major vascular surgery. *Cardiothorac Surg* 2013; 8:189-97.
50. Caille V, Squara P. Oxygen uptake-to-delivery relationship: a way to assess adequate flow. *Crit Care* 2006; 10:54-5.
51. Van Beest PA, Brander L, Jansen SP, Rommes JH, Kuiper MA, Spronk PE. Cumulative lactate. *Ann Intensive Care* 2013; 6:3-6.

References

52. Cheitfetz IM, Kern FH, Schiman SR, et al. Serum lactates correlate with mortality after operations for complex congenital heart disease. *Ann Thorac Surg* 1997; 64:735-8.
53. Bakker J, Nijsten MW, Jansen TC. Clinical use of lactate monitoring in critically ill patients. *Ann Intensive care* 2013; 3:3-12.
54. Hein OV, Ohring R, Schilling A, Oellerich M, Armstrong VW, Kox WI, Spies C. N-acetylcysteine decreases lactate signal intensities in liver tissue and improves liver function in septic shock patients, as shown by magnetic resonance spectroscopy: extended case report. *Crit Care* 2004; 2:66-71.
55. Handy J. The origin and interpretation of hyperlactemia during low oxygen delivery state. *Crit Care* 2007; 11:104-5.
56. Pittman J, Gupta KJ. Cardiac output monitoring: Will new technologies replace the pulmonart artery catheter. *Yearbook of Intensive Care and Emerg Med* 2003; 481-98.
57. Troskot R, Simurina T, Zizac M, Majstorovic K, Marinac I, Markovcic SI. Prognostic value of venoarterial carbon dioxide gradient in patients with severe sepsis and septic shock. *Croat Med J* 2010; 16:501-8.
58. Cuschieri J, Hays G, Rivers EP. Arterial-venous carbon dioxide gradient as an indicator of cardiac index: A comparison between the mixed and central circulation. *Crit Care Med* 1998; 26:A56-64.
59. Vincent JL, Rhodes A, Perel A, Martin Gs, Della Rocca G, Vallet B, et al. Clinical review: update on hemodynamic monitoring. *Crit Care* 2011; 15:229-36.
60. Mecher CE, Rackow EC, Astiz ME, et al. Venous hyper-carbia associated with severe sepsis and systemic hypoperfusion. *Crit Care Med* 1990; 18:585-9.
61. Bakker J, Vincent JL, Gris P, et al. Venous-arterial carbon dioxide gradient in human septic shock. *Chest* 1992; 101:509-15.
62. Teboul JL, Merat A, Lenique F, Berton C, Value of the venous-arterial PCO₂ gradient to reflect the oxygen supply to demand in humans: effect of dobutamine. *Crit Care Med* 1998; 26:1007-10.
63. Lamia B, Monnet X, Teboul JL. Meaning of veno-arterial PCO₂ difference in circulatory shock. *Minerva Anesthesiol* 2006; 72:597-604.
64. Herve P, Simonneau G, Girard P, Cerrina J, Mathieu M, Duroux P. Hypercapnic acidosis induced by nutrition in mechanically ventilated patients: glucose versus fat. *Crit Care Med* 1985; 13(7):537-40.
65. De Somer F, Mulholland JW, Bryan MR, Aloisio T, Van Nooten GJ, Ranucci M. O₂ delivery and CO₂ production during cardiopulmonary bypass as determinants of acute kidney injury. *Crit Care* 2011; 15(4):192-9.
66. Jensen FB. Red blood cell ph, the bohe effect, and other oxygenation-linked phenomena in blood O₂ and CO₂ transport. *Acta physiol Scand* 2004; 182:215-27.

References

67. West J. Gas transport to the periphery: how gase moved to the peripheral tissue? In: West JB, editor. Respiratory physiology. The essentials. 4th ed. Baltimore: Williams & Wilkins 1990; 12:69-85.
68. Cavaliere F, Giovannini I, Chiarla C, Conti G, Pennisi MA, Montini L, et al. Comparison of two methods to assess blood CO₂ equilibration curve in mechanically ventilated patients. *Respir Physiol Neurobiol* 2005; 146:77-83.
69. Jensen FB. Comparative analysis of autoxidation of haemoglobin. *J Exp Biol* 2001; 204:2029-33
70. McHardy GJ. The relationship between the differences in pressure and content of carbon dioxide in arterial and venous blood. *Clin Sci* 1967; 32:299-309.
71. Troškot R1, Šimurina T, Žižak M, Majstorović K, Marinac I, Mrakovčić-Šutić I. Prognostic value of venoarterial carbon dioxide gradient in patients with severe sepsis and septic shock. *Croat Med J* 2010; 51(6):501-8.
72. Groeneveld AB, Vermeij CG, Thijs LG. Arterial and mixed venous blood acid-base balance during hypoperfusion with incremental positive end-expiratory pressure in the pig. *Anesth Analg* 1991; 73:576-82.
73. Futier E1, Robin E, Jabaudon M, Guerin R, Petit A, Bazin JE, Constantin JM, Vallet B. Central venous O₂ saturation and venous-to-arterial CO₂ difference as complementary tools for goal-directed therapy during high-risk surgery. *Crit Care* 2010; 14(5):193-9.
74. Ospina-Tascón GA, Bautista-Rincón DF, Umaña M, Tafur JD, Gutiérrez A, García AF, Bermúdez W, Granados M, Arango-Dávila C, Hernández G. Persistently high venous-to-arterial carbon dioxide differences during early resuscitation are associated with poor outcomes in septic shock. *Crit Care* 2013; 17(6):294-301.
75. Dubin A, Murias G, Estenssoro E, Canales H, Badie J, Pozo M, Sottile JP, Barán M, Pálizas F, Laporte M. Intramucosal-arterial PCO₂ gap fails to reflect intestinal dysoxia in hypoxic hypoxia. *Crit Care* 2002; 6(6):514-20.
76. Adroge HJ, Rashad MN, Gorin AB, et al. Assessing acid-base status in circulatory failure. Difference between arterial and central venous blood. *N Engl J Med* 1989; 320:1312-6.
77. Van der Linden P, Rausin I, Deltell A, Bekrar Y, Gilbert E, Bakker J, et al. Detection of tissue hypoxia by arteriovenous gradient for PCO₂ and pH in anesthetized dogs during progressive hemorrhage. *Anesth Analg* 1995; 80:269-75.
78. De Backer D, Creteur J, Preiser JC, Dubois MJ, Vincent JL. Microvascular blood flow is altered in patients with sepsis. *Am J Respir Crit Care Med* 2002; 166:98-104.
79. Fink MP. Cytopathic hypoxia. Mitochondrial dysfunction as mechanism contributing to organ dysfunction in sepsis. *Crit Care Clin* 2001; 17:219-37.
80. Wendon JA, Harrison PM, Keays R, et al. Arterial-venous Ph difference and tissue hypoxia in patients with fulminant hepatic failure. *Crit Care Med* 1991; 19:1362-4.

References

81. Neviere R, Chagon JL, Teboul JL, Vallet B, Wattel F. Small intestine intramucosal PCO₂ and microvascular blood flow during hypoxic and ischemic hypoxia. *Crit Care Med* 2002; 30:379-84.
82. Dubin A, Murias G, Estenssoro E, Canales H, Badie J, Pozo M, et al. Intramucosal-arterial PCO₂ gap fails to reflect intestinal dysoxia in hypoxic hypoxia. *Crit Care* 2002; 6:514-20.
83. Vallet B, Teboul JL, Cain S, Curtis S. Venous-arterial CO₂ difference during regional ischemia or hypoxic hypoxia. *J Appl Physiol* 2000; 89:1317-21.
84. Gutierrez G. A mathematical model of tissue-blood carbon dioxide exchange during hypoxia. *Am J Respir Crit Care Med* 2004; 169:525-33.
85. De Backer D, Creteur J, Vincent JL. Perioperative optimization and right heart catheterization: what technique in which patient?. *Crit Care* 2003; 7(3):201-2.
86. Cecconi M, Corredor C, Arulkumaran N, Abuella G, Ball J, Grounds RM, Hamilton M, Rhodes A. Clinical review: Goal-directed therapy-what is the evidence in surgical patients? The effect on different risk groups. *Crit Care* 2013; 17(2):209-16.
87. Jones AE, Brown MD, Trzeciak S, Shapiro NI, Garrett JS, Heffner AC, Kline JA; Emergency Medicine Shock Research Network investigators. The effect of a quantitative resuscitation strategy on mortality in patients with sepsis: a meta-analysis. *Crit Care Med* 2008; 36(10):2734-9.
88. Cohen IL, Sheikh FM, Perkins RJ, Feustel PJ, Foster ED. Effect of hemorrhagic shock and reperfusion on respiratory quotient in swine. *Crit Care Med* 1995; 23:545-52.
89. Mekontso-Dessap A, Caselain V, Anguel N, Bahloul M, Schavvliege F, Richard C, et al. Combination of venous-arterial PCO₂ difference with arteriovenous O₂ content difference to detect anaerobic metabolism in patients. *Intensive Care Med* 2002; 28:272-7.
90. Crapo RO. Arterial blood gases: quality assessment. In: Tobin MJ, editor. *Principle and practice of intensive care monitoring*. New York: MC Graw-Hill 1998; 7:107-22.
91. d'Ortho MP, Delclaux C, Zerah F, Herigault R, Adnot S, Harf A. Use of glass capillaries avoids the time changes in high blood PCO₂ observed with plastic syringes. *Chest* 2001; 120:165-4.
92. Richard C, Monnet X, Teboul JL. Pulmonary artery catheter monitoring in 2011. *Curr Opin Crit Care* 2011; 17:296:302.
93. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, et al. Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 2008; 36:296-327.
94. Cuschier J, Rivers EP, Donnino MW, Katilius M, Jacobsen G, Nguyen HB, et al. Central venous-arterial carbon dioxide difference as an indicator of cardiac index. *Intensive Care Med* 2005; 31(6):818-22.

References

95. Vallee F, Vallet B, Mathe O, Parraguette J, Mari A, Silva S, et al. Central venous-to-arterial carbon dioxide difference: an additional target for goal-directed therapy in septic shock. *Intensive Care Med* 2008; 34:2218-25.
96. Futier E, Robin E, Jabaudon M, Guerin R, Petit A, Bazin JE, et al. Central venous O2 saturation and venous-to-arterial CO2 difference as complementary tools for goal-directed therapy during high risk surgery. *Crit Care* 2010; 14:193-7.
97. Monnet X, Julien F, Ait-Hamou N, Lequoy M, Gosset C, Jozwiak M, et al. Lactate and veno-arterial carbon dioxide difference/arterial-venous oxygen difference ratio, but not central venous oxygen saturation, predict increase in oxygen consumption in fluid responders. *Crit Care Med* 2013; 41(6):1412-20.
98. Ziauddin A, Kashmiri, Mudassir et al. Incidence of complications after cardiac surgery single unit. *Pakistan Heart* 2008; 41:418-26.
99. Giovanni P, Alessandro D, Luca R et al. Postoperative arrhythmias after cardiac surgery: Incidence, Risk factors and therapeutic management. *Cardiology Research and practice* 2014; 1:1-15.
100. Vivek R, Joan I, Richard D et al. Predictors of low cardiac output syndrome after coronary bypass. *The journal of thoracic and cardiovascular surgery* 1996; 112:38-51.
101. Khilji SA, Khan AH. Acute renal failure after cardiopulmonary bypass surgery. *J Ayub Med Coll Abbottabad* 2004; 16(3):25-8.
102. Ieva N, Donata R, Vilma K, et al. Incidence and risk factors of early delirium after cardiac surgery. *BioMed Research International* 2013; 10:115-9.
103. Jenifer Goodwin. Post-op Delerium's Toll on mental function may linger. *Health day news* 2012; 4:42-6.
104. Charles W, Christopher A, Joseph E et al. Cardio-pulmonary bypass management and neurologic outcomes; An Evidence-Based appraisal of current practices. *Anesth Analg* 2006 ; 103:21-37.
105. Pezzella AT, Ferraris VA, Lancey RA. Care of the adult cardiac surgery patients . *Curr Probl Surg* 2004; 41(6):526-74.
106. James Roxburg. National Adult Cardiac Surgery Audit Report. Annual Report 2010-2011; 6:21-5.
107. Jan Bakker, M.D.; Jean-Louis Vincent, M.D., Ph.d, F.C.C.P.; Philippe Gris, M.D.; Marc Leon, M.D.; Michel Coffernils, M.D.; and Robert J, M.D. Veno-arterial Carbon Dioxide Gradient in Human Septic Shock. *Chest* 1992; 101:509-15.
108. Jihad Mallet, Florent Pepy, Malcolm Lemzye, Gaelle Gasan, Nicolas Vangrunderbeeck, Laurent Tronchon, Benoit Vallet and Didier Thevenin, Central venous-to-arterial carbon dioxide partial pressure difference in early resuscitation from septic shock. *Eur J Anaesthesiol* 2014; 31:371-380.

References

109. Ariza M, Gothard JW, Macnaughton P, et al. Blood lactate and mixed venous-arterial PCO₂ gradient as indices of poor peripheral perfusion following cardiopulmonary bypass surgery. *Intensive Care Med* 1991; 17:320.
110. Martin Dres, Xavier Monnet, Jean-Louis Teboul. Hemodynamic management of cardiovascular failure by using PCO₂ venous-arterial difference. *J Clin Monit Comput* 2012; 26:367-74.

المخلص العربي

معظم جراحات القلب المفتوح تجري حالياً باستخدام ماكينته تجاوز القلب والرئتين، مما يجعل المريض عرضة لمجموعه مختلفه من مضاعفات ما بعد الجراحه التي تنتج عن انخفاض تدفق الدم في الدوره الدمويه الدقيقه للقلب والمخ وباقي أعضاء الجسم والتي قد تؤدي إلي قصور في أداء أعضاء الجسم، لذلك فان التعرف علي وسائل للتنبؤ بحدوث المضاعفات أو الوفاة يعتبر موضوع هام.

الفرق في نسبة ثاني أكسيد الكربون بين الدم الوريدي المركزي والدم الشرياني يمكن استخدامه كدلاله علي فاعليه الدم الوريدي علي إزالة ثاني أكسيد الكربون الذي تنتجه الأنسجه. حتي الآن فان فارق نسبة ثاني أكسيد الكربون بين الدم الوريدي المركزي والدم الشرياني تم بحثه فقط في حالات الصدمه التسمييه. لكن لم ترد تقارير حول سلوكه وعلاقته بالنتائج السريره ما بعد جراحات القلب المفتوح.

وكان هدفنا من هذا العمل هو تقييم القيمه التنبؤيه لفارق نسبة ثاني أكسيد الكربون بين الدم الوريدي المركزي والدم الشرياني لمريض جراحات القلب المفتوح وقد أجريت هذه الدراسة علي ٧٠ مريضاً الذين خضعوا لعمليات القلب المفتوح باستخدام ماكينته تجاوز القلب والرئتين وقد كانت أعمارهم تتراوح بين ١٨ و ٦٠ عاماً وقد تم سحب عينات وريديه وشريانيه معا بعد مرور ٣ ساعات ثم ٦ ساعات ثم ١٢ ساعه من الحجز بوحده عناية القلب المفتوح. وتم قياس نسبة ثاني أكسيد الكربون واللكتات ويتم ملاحظه معدل حدوث المضاعفات والوفيات خلال ٢٨ يوم.

أظهرت نتائج هذه الدراسة ما يلي:

- كان متوسط فارق نسبة ثاني أكسيد الكربون بعد ٣ ساعات ٧,١٣ مم زئبق بينما بعد مرور ٦ ساعات كان ٥,٧ مم زئبق وبعد ١٢ ساعه كان ٥,٥٩ مم زئبق.
- المرضى الذين كان لديهم فارق نسبة ثاني أكسيد الكربون ≥ 6 مم زئبق بعد ٣ ساعات يمثلون ٤١,٤ %، بعد مرور ٦ ساعات كانوا ٦١,٤ % وبعد ١٢ ساعه كانوا ٦٤,٣ %.
- كان متوسط اللكتات بعد ٣ ساعات ٢,٥٨ مليمول/لتر وبعد ٦ ساعات كان ١,٨ مليمول / لتر وبعد ١٢ ساعه كان ١,٥ مليمول / لتر.
- مستوي اللكتات كان اعلي بشكل ملحوظ في المرضى الذين لديهم فارق نسبة ثاني أكسيد الكربون اعلي من ٦ مم زئبق عن باقي المرضى.
- كان هناك انخفاض ملحوظ إحصائياً في مستوي اللكتات في المرضى الذي انخفض فيهم فارق نسبة ثاني أكسيد الكربون إلي المعدل الطبيعي ما بين الساعه الثالثه بعد العمليات والساعه السادسه بعد العمليات.
- كان معدل حدوث المضاعفات اعلي بشكل ملحوظ إحصائياً في المرضى الذين لديهم فارق نسبة ثاني أكسيد الكربون اعلي من ٦ مم زئبق.
- اختلال ضربات القلب كان أكثر المضاعفات حدوثاً حيث مثل ٢٢,٩ % من إجمالي المرضى في الدراسة، تلاه تأخر الإفاقة بنسبة ١٧,١ % ثم الهذيان بنسبة ١١,٤ % ثم فشل عضلة القلب بنسبة ١٠ % ثم فشل كلوي حاد بنسبة ٨,٦ %.
- كان معدل حدوث الوفاة خلال ٢٨ يوم من الجراحه ٤,٣ %.
- كانت القيمه الفاطعه لفارق نسبة ثاني أكسيد الكربون ٦ مم زئبق والتي عندها يمكن التنبؤ بحدوث مضاعفات بحساسيه وخصوصيه عاليه بعد ٣ ساعات، ٦ ساعات و ١٢ ساعه.



جامعة الإسكندرية
كلية الطب
قسم الطب الحرج

دراسة حول القيمة التنبؤية لفارق نسبة ثاني أكسيد الكربون بين الدم الوريدي المركزي والدم الشرياني في مرضى جراحات القلب المفتوح

رسالة مقدمة

لقسم الطب الحرج - كلية الطب - جامعة الإسكندرية
ضمن متطلبات درجة

الماجستير

في

الطب الحرج

من

تامر عاطف عبدالحميد عبدالحميد

بكالوريوس الطب والجراحة، ٢٠٠٦

كلية الطب، جامعة الإسكندرية

٢٠١٥



جامعة الإسكندرية
كلية الطب
قسم الطب الحرج

دراسة حول القيمة التنبؤية لفارق نسبة ثاني أكسيد الكربون بين الدم الوريدي المركزي والدم الشرياني في مرضى جراحات القلب المفتوح

رسالة مقدمة من

تامر عاطف عبدالحميد عبدالحميد

للحصول على درجة

الماجستير

في

الطب الحرج

التوقيع

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د/ سلوى شعبان شعراوي
أستاذ التخدير والعناية المركزة الجراحية
قسم التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

أ.د/ صلاح عبد الفتاح محمد
أستاذ التخدير والعناية المركزة الجراحية
قسم التخدير والعناية المركزة الجراحية
كلية الطب
جامعة قناة السويس

أ.م.د/ عمرو عبد الله المرسي
أستاذ مساعد الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الإسكندرية

لجنة الإشراف

موافقون

أ.د/ سلوى شعبان شعراوى

أستاذ التخدير والعناية المركزة الجراحية
قسم التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

.....

أ.م.د/ تامر عبدالله حلمى

أستاذ مساعد الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الإسكندرية

.....