

CONCLUSION

- CT is the primary imaging modality for diagnosis and follow-up of patients with known or suspected primary or secondary neoplastic conditions of the peritoneum and mesentery.
- Secondary peritoneal tumors are much more common than primary, however primary peritoneal tumor should be in the differential diagnosis in presence of peritoneal soft tissue masses with no detected primary non peritoneal tumor.
- There are other tumor like pathological conditions affect the peritoneum and mesentery as granulomatous peritonitis, sclerosing mesenteritis, inflammatory pseudotumor and endometriosis, these lesions could also manifest as peritoneal soft tissue with or without ascites.
- In case of primary non-peritoneal tumor associated with unexplained ascites, the peritoneum should be inspected carefully to look for peritoneal metastasis.
- Even with resection of the primary tumor and clear surgical bed with no residual or recurrent masses, periodic follow up CECT is still recommended to detect any denovo peritoneal deposits even if small ones.
- Correlation of imaging findings with pathological analysis is essential to reach an appropriate diagnosis and initiate management.

REFERENCES

1. Patela CM, Sahdevb A, Reznicka RH. CT, MRI and PET imaging in peritoneal malignancy. *Cancer Imaging* 2011;11:123–39.
2. Tirkes T, Sandrasegaran K, Patel AA, Hollar MA, Tejada JG, Tann M, et al. Peritoneal and retroperitoneal anatomy and its relevance for cross-sectional imaging. *Radiographics* 2012; 32:437– 51.
3. Battifora H, McCaughey WTE. Development, anatomy, and function of the serosal membranes. In: Rosai J, (ed). *Tumors of the serosal membranes*. Washington, DC: Armed Forces Institute of Pathology; 1995. pp. 1–7.
4. Moore KL. The coelomic cavities and mesenteries. In: Moore KL, (ed). *The developing human: clinically oriented embryology*. 3rded. Philadelphia, Pa: Saunders; 1982. pp. 303–17.
5. Jeong YJ, Kim S, Kwak SW, Lee NK, Lee JW, Kim KI, et al. Neoplastic and nonneoplastic conditions of serosal membrane origin: CT findings. *Radiographics*. 2008;28(3):801-18.
6. Kim S, Kim TU, Lee JW, Lee TH, Lee SH, Jeon TY, et al. The perihepatic space: comprehensive anatomy and CT features of pathologic conditions. *Radiographics* 2007;27(1): 129– 43.
7. DeMeo JH, Fulcher AS, Austin RF. Anatomic CT demonstration of the peritoneal spaces, ligaments, and mesenteries: normal and pathologic processes. *Radiographics* 1995;15(4):755–70.
8. Oliphant M, Berne AS, Meyers MA. The subperitoneal space of the abdomen and pelvis: planes of continuity. *Am J Roentgenol* 1996;167(6): 1433–9.
9. Yoo E, Kim JH, Kim MJ, Yu JS, Chung JJ, Yoo HS, et al. Greater and lesser omenta: normal anatomy and pathologic processes. *Radiographics* 2007;27(3):707–20.
10. Charnsangavej C, DuBrow RA, Varma DG, Herron DH, Robinson TJ, Whitley NO. CT of the mesocolon: part 1 anatomic considerations. *Radiographics* 1993;13(5):1035– 45.
11. Takeyama N, Gokan T, Ohgiya Y, Satoh S, Hashizume T, Hataya K, et al. CT of internal hernias. *Radiographics* 2005;25(4):997–1015.
12. Hewitt MJ, Hall GD, Wilkinson N, Perren TJ, Lane G, Spencer JA.. Image-guided biopsy in women with breast cancer presenting with peritoneal carcinomatosis. *Int J Gynecol Cancer* 2006; 16(1): 108–10.
13. Hewitt MJ, Anderson K, Hall GD, Weston M, Hutson R, Wilkinson N, et al. Women with peritoneal carcinomatosis of unknown origin: efficacy of image-guided biopsy to determine site-specific diagnosis. *BJOG* 2007; 114: 46–50.

References

14. Pannu HK, Bristow RE, Montz FJ, Fishman EK. Multidetector CT of peritoneal carcinomatosis from ovarian cancer. *Radiographics* 2003; 23: 687–701.
15. Pannu HK, Horton KM, Fishman EK. Thin section dual-phase multidetector-row computed tomography detection of peritoneal metastases in gynecologic cancers. *J Comput Assist Tomogr* 2003; 27: 333–40.
16. Coakley FV, Choi PH, Gougoutas CA, Pothuri B, Venkatraman E, Chi D, et al. Peritoneal metastases: detection with spiral CT in patients with ovarian cancer. *Radiology* 2002; 223: 495–9.
17. Franiel T, Diederichs G, Engelken F, Elgeti T, Rost J, Rogalla P. Multi-detector CT in peritoneal carcinomatosis: diagnostic role of thin slices and multiplanar reconstructions. *Abdom Imaging* 2009; 34: 49–54.
18. Marin D, Catalano C, Baski M, Di Martino M, Geiger D, Di Giorgio A, et al. 64-Section multi-detector row CT in the preoperative diagnosis of peritoneal carcinomatosis: correlation with histopathological findings. *Abdom Imaging* 2010; 35: 694–700.
19. de Bree E, Koops W, Kröger R, van Ruth S, Witkamp AJ, Zoetmulder FA. Peritoneal carcinomatosis from colorectal or appendiceal origin: correlation of preoperative CT with intraoperative findings and evaluation of inter observer agreement. *J Surg Oncol* 2004; 86: 64–73.
20. Low RN. MR imaging of the peritoneal spread of malignancy. *Abdom Imaging* 2007; 32: 267–83.
21. Akisik FM, Sandrasegaran K, Aisen AM, Lin C, Lall C. Abdominal MR imaging at 3.0 T. *Radiographics* 2007;27(5):1433-44.
22. Levy A, Medjhouli A, Caramella C, Zareski E, Berges O, Chargari C, et al. Interest of diffusion-weighted echo-planar MR imaging and apparent diffusion coefficient mapping in gynecological malignancies: a review. *J Magn Reson Imaging* 2011;33(5):1020–7.
23. Satoh Y, Ichikawa T, Motosugi U, Kimura K, Sou H, Sano K, et al. Diagnosis of peritoneal dissemination: comparison of 18F-FDG PET/CT, diffusion-weighted MRI, and contrast-enhanced MDCT. *Am J Roentgenol* 2011;196(2): 447–53.
24. Fujii S, Matsusue E, Kanasaki Y, Kanamori Y, Nakanishi J, Sugihara S, et al. Detection of peritoneal dissemination in gynecological malignancy: evaluation by diffusion-weighted MR imaging. *Eur Radiol* 2008; 18: 18–23.
25. Low RN. Diffusion-weighted MR imaging for whole body metastatic disease and lymphadenopathy. *Magn Reson Imaging Clin N Am* 2009; 17: 245-61.
26. Pannu HK, Bristow RE, Cohade C, Fishman EK, Wahl RL. PET-CT in recurrent ovarian cancer: initial observations. *Radiographics* 2004;24:209–23.

References

27. Tanaka T, Kawai Y, Kanai M, Taki Y, Nakamoto Y, Takabayashi A. Usefulness of FDG-positron emission tomography in diagnosing peritoneal recurrence of colorectal cancer. *Am J Surg* 2002;184: 433–6.
28. Turlakow A, Yeung HW, Salmon AS, Macapinlac HA, Larson SM. Peritoneal carcinomatosis: role of (18)F-FDG PET. *J Nucl Med* 2003;44:1407–12.
29. Levy AD, Arnáiz J, Shaw JC, Sobin LH. Primary peritoneal tumors: imaging features with pathologic correlation. *Radiographics* 2008;28:583–607.
30. Meyers MA, Oliphant M, Berne AS, Feldberg MA. The peritoneal ligaments and mesenteries: Pathways of intra-abdominal spread of disease. *Radiology* 1987; 163: 595–604.
31. Oliphant M, Berne A, Meyers MA. Subperitoneal spread of intra-abdominal disease. In: Myers MA, (ed). *Computed tomography of the gastrointestinal tract including the peritoneal cavity and mesentery*. New York: Springer Verlag; 1986. pp. 95–137.
32. Meyers MA. Distribution of intraabdominal malignant seedling: Dependency on dynamics of flow of ascetic fluid. *AJR* 1973; 119: 198–206.
33. Goffinet DR, Castellino RA, Kim H, Dorfman RF, Fuks Z, Rosenberg SA, et al. Staging laparotomies in unselected patients with non-Hodgkin's lymphoma. *Cancer* 1973; 32: 672–81.
34. Whitley NO, Bohlman ME, Baker LP. CT patterns of mesenteric disease. *J Comput Assist Tomogr* 1982;6: 490–6.
35. Mueller PR, Ferrucci JT Jr, Harbin WP, Kirkpatrick RH, Simeone JF, Wittenberg J. Appearance of lymphomatous involvement of the mesentery by ultrasonography and body computer tomography: the 'sandwich sign'. *Radiology* 1980; 134: 467–73.
36. Kawashima A, Fishman EK, Kuhlman JE, Schuchter LM. CT of malignant melanoma: Patterns of small bowel and mesenteric involvement. *J Comput Assist Tomogr* 1991; 15(4): 570–4.
37. Chu DZ, Lang NP, Thompson C, Osteen PK, Westbrook KC. Peritoneal carcinomatosis in nongynecologic malignancy: a prospective study of prognostic factors. *Cancer* 1989;63:364–7.
38. Sadeghi B, Arvieux C, Glehen O, Beaujard AC, Rivoire M, Baulieux J, et al. Peritoneal carcinomatosis from non-gynecologic malignancies: results of the EVOCAPE 1 multicentric prospective study. *Cancer* 2000 ;88:358–63.
39. Hanbidge AE, Lynch D, Wilson SR. US of the peritoneum. *Radiographics* 2003;23:663–85.
40. Yeh HC. Ultrasonography of peritoneal tumors. *Radiology* 1979;133:419–24.
41. Walkey MM, Friedman AC, Sohotra P, Radecki PD. CT manifestations of peritoneal carcinomatosis. *AJR Am J Roentgenol* 1988;150:1035–41.

References

42. Jacquet P, Jelinek JS, Steves MA, Sugarbaker PH. Evaluation of computed tomography in patients with peritoneal carcinomatosis. *Cancer* 1993;72: 1631–6.
43. Hamrick-Turner JE, Chiechi MV, Abbitt PL, Ros PR. Neoplastic and inflammatory processes of the peritoneum, omentum, and mesentery: diagnosis with CT. *Radiographics* 1992;12:1051–68.
44. Levy AD, Shaw JC, Sobin LH. Secondary Tumors and Tumor like Lesions of the Peritoneal Cavity: Imaging Features with Pathologic Correlation. *RadioGraphics* 2009; 29:347–73.
45. Moran BJ, Cecil TD. The etiology, clinical presentation, and management of pseudomyxoma peritonei. *Surg Oncol Clin N Am* 2003;12:585–603.
46. Fenoglio-Preiser CM. *Gastrointestinal pathology: an atlas and text*. 3rded. Philadelphia, Pa: Wolters Kluwer/Lippincott Williams & Wilkins; 2008. p.1296.
47. Bradley RF, Stewart JH 4th, Russell GB, Levine EA, Geisinger KR. Pseudomyxoma peritonei of appendiceal origin: a clinicopathologic analysis of 101 patients uniformly treated at a single institution, with literature review. *Am J Surg Pathol* 2006; 30: 551–9.
48. Capella C, Solcia G, Sobin LH, Arnold R, et al. Adenocarcinoma of the appendix. In: Aaltonen LA, Hamilton SR, (eds). *World Health Organization classification of tumors: pathology and genetics of tumors of the digestive system*. Lyon, France: IARC; 2000. pp. 95–102.
49. Ronnett BM, Zahn CM, Kurman RJ, Kass ME, Sugarbaker PH, Shmookler BM. Disseminated peritoneal adenomucinosis and peritoneal mucinous carcinomatosis: a clinicopathologic analysis of 109 cases with emphasis on distinguishing pathologic features, site of origin, prognosis, and relationship to “pseudomyxoma peritonei.” *Am J Surg Pathol* 1995; 19:1390–408.
50. Pai RK, Longacre TA. Pseudomyxoma peritonei syndrome: classification of appendiceal mucinous tumors. In: Ceelen WP, (ed). *Peritoneal carcinomatosis: a multidisciplinary approach*. New York, NY: Springer; 2007. pp. 71–107.
51. Seidman JD, Elsayed AM, Sobin LH, Tavassoli FA. Association of mucinous tumors of the ovary and appendix: a clinicopathologic study of 25 cases. *Am J Surg Pathol* 1993;17:22–34.
52. Young RH, Gilks CB, Scully RE. Mucinous tumors of the appendix associated with mucinous tumors of the ovary and pseudomyxoma peritonei: a clinicopathological analysis of 22 cases supporting an origin in the appendix. *Am J Surg Pathol* 1991;15: 415–29.
53. Hellmer H. Die konturen des rechten leberlappens beim ascites. *Acta Radiol* 1942;23:533–40.

References

54. Miller DL, Udelsman R, Sugarbaker PH. Calcification of pseudomyxoma peritonei following intraperitoneal chemotherapy: CT demonstration. *J Comput Assist Tomogr* 1985;9:1123–4.
55. Sulkin TV, O'Neill H, Amin AI, Moran B. CT in pseudomyxoma peritonei: a review of 17 cases. *Clin Radiol* 2002;57:608–13.
56. Seshul MB, Coulam CM. Pseudomyxoma peritonei: computed tomography and sonography. *AJR Am J Roentgenol* 1981;136:803–6.
57. Young RH. Pseudomyxoma peritonei and selected other aspects of the spread of appendiceal neoplasms. *Semin Diagn Pathol* 2004;21:134–50.
58. Nador RG, Cesarman E, Chadburn A, Dawson DB, Ansari MQ, Sald J, et al. Primary effusion lymphoma: a distinct clinicopathologic entity associated with the Kaposi's sarcoma-associated herpes virus. *Blood* 1996;88:645–56.
59. Kim Y, Cho O, Song S, Lee H, Rhim H, Koh B. Peritoneal lymphomatosis: CT findings. *Abdom Imaging* 1998;23:87–90.
60. Lynch MA, Cho KC, Jeffrey RB Jr, Alterman DD, Federle MP. CT of peritoneal lymphomatosis. *AJR Am J Roentgenol* 1988;151:713–5.
61. Ferrozzi F, Tognini G, Mulonzia NW, Bova D, Pavone P. Primary effusion lymphomas in AIDS: CT findings in two cases. *Eur Radiol* 2001;11: 623–5.
62. Morassut S, Vaccher E, Balestreri L, Gloghini A, Gaidano G, Volpe R, et al. HIV associated human herpesvirus 8-positive primary lymphomatous effusions: radiologic findings in six patients. *Radiology* 1997;205:459–63.
63. Banks PM, Warnke RA. Primary effusion lymphoma. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, (eds). *World Health Organization classification of tumors: pathology and genetics of tumors of the haematopoietic and lymphoid tissue*. Lyon, France: IARC; 2001. pp. 179–80.
64. Zanarini D, Sugarbaker PH. Extremity soft tissue sarcoma with metastases to abdominopelvic surfaces. *J Surg Oncol* 1997;64:68–73.
65. DeMatteo RP, Lewis JJ, Leung D, Mudan SS, Woodruff JM, Brennan MF. Two hundred gastrointestinal stromal tumors: recurrence patterns and prognostic factors for survival. *Ann Surg* 2000;231: 51–8.
66. Levy AD. Imaging and staging of gastrointestinal stromal tumors. In: Jankowski JAZ, (ed). *Gastrointestinal oncology: a critical multidisciplinary team approach*. Oxford, England: Blackwell; 2008. pp. 240–4.
67. Attanoos RL, Gibbs AR. Pathology of malignant mesothelioma. *Histopathology* 1997;30:403–18.
68. Churg A, Cagle PT, Roggli VL. Tumors of the serosal membranes. *Atlas of tumor pathology*. Silver Spring, Md: ARP; 2006. pp 107–9.

References

69. Roggli VL, Sharma A, Butnor KJ, Sporn T, Vollmer RT. Malignant mesothelioma and occupational exposure to asbestos: a clinicopathological correlation of 1445 cases. *Ultrastruct Pathol* 2002;26:55–65.
70. Metintas M, Hillerdal G, Metintas S. Malignant mesothelioma due to environmental exposure to erionite: follow-up of a Turkish emigrant cohort. *Eur Respir J* 1999;13:523–6.
71. Guest PJ, Reznick RH, Selleslag D, Geraghty R, Slevin M. Peritoneal mesothelioma: the role of computed tomography in diagnosis and follow up. *Clin Radiol* 1992;45:79–84.
72. Puvaneswary M, Chen S, Proietto T. Peritoneal mesothelioma: CT and MRI findings. *Australas Radiol* 2002;46:91–6.
73. Ros PR, Yuschok TJ, Buck JL, Shekitka KM, Kaude JV. Peritoneal mesothelioma: radiologic appearances correlated with histology. *Acta Radiol* 1991;32:355–8.
74. Whitley NO, Brenner DE, Antman KH, Grant D, Aisner J. CT of peritoneal mesothelioma: analysis of eight cases. *AJR Am J Roentgenol* 1982;138:531–5.
75. Lovell FA, Cranston PE. Well-differentiated papillary mesothelioma of the peritoneum. *AJR Am J Roentgenol* 1990;155:1245–6.
76. Mok SC, Schorge JO, Welch WR, et al. Peritoneal tumors. In: Tavassoli FA, Devilee P, (eds). *Pathology and genetics of tumors of the breast and female genital organs*. Lyon, France: IARC; 2003. pp. 197–202.
77. Mayall FG, Gibbs AR. The histology and immunohistochemistry of small cell mesothelioma. *Histopathology* 1992;20:47–51.
78. González-Moreno S, Yan H, Alcorn KW, Sugarbaker PH. Malignant transformation of “benign” cystic mesothelioma of the peritoneum. *J Surg Oncol* 2002;79:243–51.
79. Weiss SW, Tavassoli FA. Multicystic mesothelioma: an analysis of pathologic findings and biologic behavior in 37 cases. *Am J Surg Pathol* 1988;12:737–46.
80. Ross MJ, Welch WR, Scully RE. Multilocular peritoneal inclusion cysts (so-called cystic mesotheliomas). *Cancer* 1989;64:1336–46.
81. Kim JS, Lee HJ, Woo SK, Lee TS. Peritoneal inclusion cysts and their relationship to the ovaries: evaluation with sonography. *Radiology* 1997;204:481–4.
82. Wasnik AP, Menias CO, Platt JF, Lalchandani UR, Bedi DG, Elsayes KM. Multimodality imaging of ovarian cystic lesions: Review with an imaging based algorithmic approach. *World J Radiol* 2013; 5(3): 113–25.
83. Schorge JO, Muto MG, Lee SJ, Huang LW, Welch WR, Bell DA, et al. BRCA1-related papillary serous carcinoma of the peritoneum has a unique molecular pathogenesis. *Cancer Res* 2000;60:1361–4.

References

84. Truong LD, Maccato ML, Awalt H, Cagle PT, Schwartz MR, Kaplan AL. Serous surface carcinoma of the peritoneum: a clinicopathologic study of 22 cases. *Hum Pathol* 1990;21:99–110.
85. Shmueli E, Leider-Trejo L, Schwartz I, Aderka D, Inbar M. Primary papillary serous carcinoma of the peritoneum in a man. *Ann Oncol* 2001;12:563–7.
86. Zhou J, Iwasa Y, Konishi I, Kan N, Kannagi R, Kobashi Y, et al. Papillary serous carcinoma of the peritoneum in women: a clinicopathologic and immunohistochemical study. *Cancer* 1995;76:429–36.
87. Chiou SY, Sheu MH, Wang JH, Chang CY. Peritoneal serous papillary carcinoma: a reappraisal of CT imaging features and literature review. *Abdom Imaging* 2003;28:815–9.
88. Furukawa T, Ueda J, Takahashi S, Higashino K, Shimura K, Tsujimura T, et al. Peritoneal serous papillary carcinoma: radiological appearance. *Abdom Imaging* 1999;24:78–81.
89. Morita H, Aoki J, Taketomi A, Sato N, Endo K. Serous surface papillary carcinoma of the peritoneum: clinical, radiologic, and pathologic findings in 11 patients. *AJR Am J Roentgenol* 2004;183:923–8.
90. Stafford-Johnson DB, Bree RL, Francis IR, Korobkin M. CT appearance of primary papillary serous carcinoma of the peritoneum. *AJR Am J Roentgenol* 1998;171:687–9.
92. Kushner BH, LaQuaglia MP, Wollner N, Meyers PA, Lindsley KL, Ghavimi F, et al. Desmoplastic small round-cell tumor: prolonged progression-free survival with aggressive multimodality therapy. *J Clin Oncol* 1996;14:1526–31.
93. Quaglia MP, Brennan MF. The clinical approach to desmoplastic small round cell tumor. *Surg Oncol* 2000;9:77–81.
94. Pickhardt PJ, Fisher AJ, Balfe DM, Dehner LP, Huettner PC. Desmoplastic small round cell tumor of the abdomen: radiologic-histopathologic correlation. *Radiology* 1999;210:633–8.
95. Bellah R, Suzuki-Bordalo L, Brecher E, Ginsberg JP, Maris J, Pawel BR. Desmoplastic small round cell tumor in the abdomen and pelvis: report of CT findings in 11 affected children and young adults. *AJR Am J Roentgenol* 2005;184:1910–4.
96. Sabaté JM, Torrubia S, Roson N, Matias-Guiu X, Gómez A. Intra-abdominal desmoplastic small round-cell tumor: a rare cause of peritoneal malignancy in young people. *Eur Radiol* 2000;10:817–9.
97. Lae ME, Roche PC, Jin L, Lloyd RV, Nascimento AG. Desmoplastic small round cell tumor: a clinicopathologic, immunohistochemical, and molecular study of 32 tumors. *Am J Surg Pathol* 2002;26:823–35.
98. Gerald WL, Ladanyi M, de Alava E, Cuatrecasas M, Kushner BH, LaQuaglia MP, et al. Clinical, pathologic, and molecular spectrum of tumors associated with t(11;22)(p13;q12): desmoplastic small round-cell tumor and its variants. *J Clin Oncol* 1998;16:3028–36.

References

99. Levy AD, Rimola J, Mehrotra AK, Sobin LH. From the archives of the AFIP: Benign fibrous tumors and tumor like lesions of the mesentery: radiologic-pathologic correlation. *Radiographics* 2006; 26:245–64.
100. Deering S, Miller B, Kopelman JN, Reed M. Recurrent leiomyomatosis peritonealis disseminata exacerbated by in vitro fertilization. *Am J Obstet Gynecol* 2000;182:725–6.
101. Lausen I, Jensen OJ, Andersen E, Lindahl F. Disseminated peritoneal leiomyomatosis with malignant change in a male. *Virchows Arch A Pathol Anat Histopathol* 1990;417:173–5.
102. Nguyen GK. Disseminated leiomyomatosis peritonealis: report of a case in a postmenopausal woman. *Can J Surg* 1993;36:46–8.
103. Fulcher AS, Szucs RA. Leiomyomatosis peritonealis disseminata complicated by sarcomatous transformation and ovarian torsion: presentation of two cases and review of the literature. *Abdom Imaging* 1998;23:640–4.
104. Papadatos D, Taourel P, Bret PM. CT of leiomyomatosis peritonealis disseminata mimicking peritoneal carcinomatosis. *AJR Am J Roentgenol* 1996;167:475–6.
105. Abulafia O, Angel C, Sherer DM, Fultz PJ, Bonfiglio TA, DuBeshter B. Computed tomography of leiomyomatosis peritonealis disseminata with malignant transformation. *Am J Obstet Gynecol* 1993;169:52–4.
106. Sheth S, Horton KM, Garland MR, Fishman EK. Mesenteric Neoplasms: CT Appearances of Primary and Secondary Tumors and Differential Diagnosis. *RadioGraphics* 2003; 23:457–73.
107. Jones IT, Jagelman DG, Fazio VW, Lavery IC, Weakley FL, McGannon E. Desmoid tumors in familial polyposis coli. *Ann Surg* 1986; 204:94–7.
108. Brooks AP, Reznick RH, Nugent K, Farmer KC, Thomson JP, Phillips RK. CT appearances of desmoid tumors in familial adenomatous polyposis: further observations. *Clin Radiol* 1994; 49:601–7.
109. Einstein DM, Tagliabue JR, Desai RK. Abdominal desmoids: CT findings in 25 patients. *AJR Am J Roentgenol* 1991; 157:275–9.
110. Casillas J, Sais GJ, Greve JL, Iparraguirre MC, Morillo G. Imaging of intra- and extraabdominal desmoid tumors. *Radiographics* 1991; 11:959–68.
111. Kawashima A, Goldman SM, Fishman EK, Kuhlman JE, Onitsuka H, Fukuya T, et al. CT of intraabdominal desmoid tumors: is the tumor different in patients with Gardner's disease?. *AJR Am J Roentgenol* 1994; 162:339–42.
112. Pickhardt PJ, Bhalla S. Primary neoplasms of peritoneal and sub-peritoneal origin: CT findings. *Radiographics* 2005;25:983-95.
113. Park IS, Kye BH, Kim HS, Kim HJ, Cho HM, Yoo C. Primary mesenteric carcinoid tumor. *J Korean Surg Soc* 2013; 84(2): 114–7.

References

114. Mindelzun RE, Jeffrey RB Jr, Lane MJ, Silverman PM. The misty mesentery on CT: differential diagnosis. *AJR Am J Roentgenol* 1996; 167:61–5.
115. Bernardino ME, Jing BS, Wallace S. Computed tomography diagnosis of mesenteric masses. *AJR Am J Roentgenol* 1979; 132:33–6.
116. Wallace S, Ajani JA, Charnsangavej C, DuBrow R, Yang DJ, Chuang VP, et al. Carcinoid tumors: imaging procedures and interventional radiology. *World J Surg* 1996; 20:147–56.
117. Pantongrag-Brown L, Buetow PC, Carr NJ, Lichtenstein JE, Buck JL. Calcification and fibrosis in mesenteric carcinoid tumor: CT findings and pathologic correlation. *AJR Am J Roentgenol* 1995; 164:387–91.
118. Burke AP, Thomas RM, Elsayed AM, Sobin LH. Carcinoids of the jejunum and ileum: an immunohistochemical and clinicopathologic study of 167 cases. *Cancer* 1997; 79:1086–93.
119. Godwin JD. Carcinoid tumors: an analysis of 2,837 cases. *Cancer* 1975; 36:560–9.
120. Buck JL, Sobin LH. Carcinoids of the gastrointestinal tract. *RadioGraphics* 1990; 10:1081–95.
121. Woodard PK, Feldman JM, Paine SS, Baker ME. Midgut carcinoid tumors: CT findings and biochemical profiles. *J Comput Assist Tomogr* 1995; 19:400–5.
122. Silverman PM, Baker ME, Cooper C, Kelvin FM. Computed tomography of mesenteric disease. *Radiographics* 1987; 7:309–20.
123. McDermott VG, Low VH, Keogan MT, Lawrence JA, Paulson EK. Malignant melanoma metastatic to the gastrointestinal tract. *AJR Am J Roentgenol* 1996; 166:809–13.
124. Kawashima A, Fishman EK, Kuhlman JE, Schuchter LM. CT of malignant melanoma: patterns of small bowel and mesenteric involvement. *J Comput Assist Tomogr* 1991; 15:570–4.
125. Hardy SM. The sandwich sign. *Radiology* 2003;226(3):651–2.
126. Ayantunde AA, Parsons SL. Pattern and prognostic factors in patients with malignant ascites: a retrospective study. *Ann Oncol* 2007;18:945–9.
127. Le O. Patterns of peritoneal spread of tumor in the abdomen and pelvis. *World J Radiol* 2013; 5(3): 106–12.
128. Agarwal A, Yeh BM, Breiman RS, Qayyum A, Coakley FV. Peritoneal calcification: causes and distinguishing features on CT. *AJR Am J Roentgenol* 2004; 182(2):441-5.
129. Sugarbaker PH, Ronnett BM, Archer A, Averbach AM, Bland R, Chang D, et al. Pseudomyxoma peritonei syndrome. *Adv Surg*1996;30:233–80.
130. Healy JC. Detection of peritoneal metastasis. *Cancer imaging* 2001;1(2):4–12.

References

131. Çinar HG, Eksioğlu AS, Ucan B, Tasci-Yıldız Y, Uner C, Pala Akdoğan M, et al. Extensive peritoneal lymphomatosis in three pediatric cases with burkitt lymphoma: computed tomography findings. Poster, Presented In: European Conference of Radiology, March 6-10; Austria, Vienna; 2014.
132. Ha HK, Jung JI, Lee MS, Choi BG, Lee MG, Kim YH, et al. CT differentiation of tuberculous peritonitis and peritoneal carcinomatosis. *AJR Am J Roentgenol.* 1996;167(3):743-8.
133. Burke AP, Sobin LH, Shekitka KM, Federspiel BH, Helwig EB. Intra-abdominal fibromatosis: a pathologic analysis of 130 tumors with comparison of clinical subgroups. *Am J Surg Pathol* 1990;14(4):335–41.
134. Kim SJ, Kim HH, Kim YH, Hwang SH, Lee HS, Park do J, et al. Peritoneal metastasis: detection with 16- or 64 -detector row CT in patients undergoing surgery for gastric cancer. *Radiology* 2009; 253(2): 407-15.
135. Rubini G, Altini C, Notaristefano A, Merenda N, Rubini D, Ianora AA, et al. Role of 18F-FDG PET/CT in diagnosing peritoneal carcinomatosis in the restaging of patient with ovarian cancer as compared to contrast enhanced CT and tumor marker Ca-125. *Rev Esp Med Nucl Imagen Mol* 2014;33(1):22-7.
136. Spencer JA, Swift SE, Wilkinson N, Boon AP, Lane G, Perren TJ. Peritoneal Carcinomatosis: Image-guided Peritoneal Core Biopsy for Tumor Type and Patient Care. *Radiology* 2001; 221:173–7.

المُلخَص العَرَبِي

الغشاء البيروتوني هو غشاء مصلي معقد وكبير، يعتبر مكان شائع للأورام. أورام البيروتون الأولية أقل تواترا بكثير من الأورام الثانوية.

الرؤية الواضحة للغشاء البيروتوني و المساريقي بواسطة الأشعة المقطعية أدت الى زيادة عظيمة في معرفة التكوين الطبيعي للغشاء البيروتوني، المساريقي وأ ربطته.

الهدف من هذه الدراسة هو دراسة دور الأشعة المقطعية في تقييم أورام الغشاء البيروتوني والمساريقي مع ربط نتائج الأشعة المقطعية بنتائج التحليل الباثولوجي للعينات، وشملت هذه الدراسة ثلاثين حالة، تراوحت أعمارهم بين (٥-٧٣ سنة) بمتوسط عمر (٤٤.٨ سنة)، وشملت ١٧ حالة من الاناث (٥٦.٧%) و ١٣ حالة من الذكور (٤٣.٣%).

الأورام الثانوية كانت اكثر شيوعا من الأولية حيث كانت الأورام الثانوية ثلاثة وعشرين حالة (٧٦.٧%) و الأورام الأولية كانت سبع حالات (٢٣.٣%)، و لقد وجد أن أكثر المصادر لأورام الغشاء البيروتوني الثانوية هي أورام المبيضين يليها أورام الجهاز الهضمي.

وجد أيضا في هذه الدراسة أن الثرب الأكبر هو أكثر جزء في الغشاء البيروتوني تأثرا بالأورام الثانوية و أن أكثر مظهر له في الأشعة المقطعية هو الكعكة الثربية.

أظهر الربط بين تشخيص الأشعة متعددة المقاطع ونتائج تحليل العينات أن الأشعة المقطعية كانت قادرة على اكتشاف الأورام البيروتونية وكانت قادرة على التشخيص الصحيح و تحديد نوع الورم في ستة وعشرين حالة (٨٦.٦%).

خلال هذه الدراسة لم تكن الأشعة المقطعية مفيدة فقط في اكتشاف و تشخيص الأورام البيروتونية بلأضافة الى التشخيص المرحلي ولكن كانت أيضا مفيدة في أخذ عينات التحليل الباثولوجي.

و بهذا يكون الربط بين نتائج الأشعة المقطعية و التاريخ المرضي للحالة مع التحليل الباثولوجي للعينات أساسي من أجل التشخيص الدقيق للحالات و بدء العلاج المناسب.



جامعة الإسكندرية
كلية الطب
قسم الأشعة التشخيصية والتدخلية

دور الأشعة متعددة المقاطع في تقييم أورام الغشاء البيروتوني والمساريقي

رسالة مقدمة

لقسم الأشعة التشخيصية والتدخلية - كلية الطب - جامعة الإسكندرية
ضمن متطلبات درجة

الماجستير

فى

الأشعة التشخيصية والتدخلية

من

داليا عبد رب النبي محمد عبد الهادي
بكالوريوس الطب والجراحة، ٢٠٠٩
كلية الطب، جامعة الإسكندرية

[٢٠١٥]



جامعة الإسكندرية
كلية الطب
قسم الأشعة التشخيصية والتدخلية

دور الأشعة متعددة المقاطع في تقييم أورام الغشاء البيروتوني والمساريقي

رسالة مقدمة من

داليا عبد رب النبي محمد عبد الهادي

للحصول على درجة

الماجستير

فى

الأشعة التشخيصية والتدخلية

التوقيع

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د/ صلاح الدين دسوقي أبو العينين
أستاذ الأشعة التشخيصية والتدخلية
قسم الأشعة التشخيصية والتدخلية
كلية الطب
جامعة الإسكندرية

أ.د/ خالد محمد مغازي
أستاذ الأشعة التشخيصية والتدخلية
قسم الأشعة التشخيصية والتدخلية
كلية الطب
جامعة الإسكندرية

أ.د/ أحمد سعد أحمد
أستاذ الجراحة العامة
قسم الجراحة العامة
معهد البحوث الطبية
جامعة الإسكندرية

التاريخ / /

موافقون

لجنة الإشراف

.....

أ.د/ خالد محمد مغازي

أستاذ الأشعة التشخيصية والتدخلية
قسم الأشعة التشخيصية والتدخلية
كلية الطب
جامعة الإسكندرية

.....

د/ نيفين محمد فريد الديب

أستاذ مساعد الباثولوجي
قسم الباثولوجي
كلية الطب
جامعة الإسكندرية

المشرف المشارك

.....

د/ عمرو محمد العبد

مدرس الأشعة التشخيصية والتدخلية
قسم الأشعة التشخيصية والتدخلية
كلية الطب
جامعة الإسكندرية