

LIST OF TABLES

<u>Title</u>	Page
Table (1): Sugarcane genotypes parentage and time of release used in the present study.	39
Table (2): Murashige and Skoog medium composition.	40
Table (3): The nucleotide sequences of primers used for RAPD analysis.	42
Table (4): Callus induction percentage and day to callus initiation of three sugarcane genotypes	53
Table (5): Percentage of embryogenic callus of three sugarcane genotypes	55
Table (6): Means of relative growth rate (RGR) and Percentage water content with standard error between sugarcane genotypes.	60
Table (7): Means of callus relative growth rate (RGR) and water content (% WC) derived from leaf explants of sugarcane after treated with different concentrations mannitol.	60
Table (8): K ⁺ and Na ⁺ ions content between the three sugarcane genotypes.	69
Table (9): Means of K ⁺ and Na ⁺ in callus in sugarcane after treated with different concentrations mannitol.	69
Table (10): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in GT 54-9 callus.	74
Table (11): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in G 84-47 callus.	75
Table (12): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in ph 8013 callus.	76
Table (13): Shoot, root and number of shoot formation for the three genotypes under the study.	79
Table (14): Interaction between shoot, root formation and number day for root formation for the three genotypes under the study.	79
Table (15): DNA polymorphism using randomly amplified polymorphic DNA with primer OPD-02 (54: GT 54-9, 80: ph 8013, 84:G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	87
Table (16): DNA polymorphism using randomly amplified polymorphic DNA with primer OPD-03(54: GT 54-9 80:ph 8013 84:G 84-47) (C: 0 I: 100mM II: 200mM III: 300mM mannitol).	89
Table (17): DNA polymorphism using randomly amplified polymorphic DNA with primer OPH-03(54: GT 54-9, 80: ph 8013, 84:G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	91
Table (18): DNA polymorphism using randomly amplified polymorphic	93

DNA with primer OPO-01(54: GT 54-9, 80: ph 8013, 84:G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).

Table (19): DNA polymorphism using randomly amplified polymorphic DNA with primer OPO-02 (54: GT 54-9, 80: ph 8013, 84:G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	95
Table (20): Similarity indices (%) among the three cultivars using five random amplified (RAPD) primers.	97

LIST OF FIGURES

<u>Title</u>	Page
Figure (1): Material of sugarcane used in the current study (a) Sugarcane leaf roll and (b) leaf sections on solidified medium.	45
Figure (2): Different callus induction in three sugarcane genotypes GT 54-9, G 84-47 and ph 8013	54
Figure (3): Callus induction and day of callus initiation in three sugarcane genotypes.	54
Figure (4): Embryogenic and non-embryogenic callus induced from leaf explant of sugarcane genotypes.	56
Figure (5): percentage of embryogenic callus in three sugarcane genotypes.	56
Figure (6): Difference in relative growth rate for all genotypes.	61
Figure (7): Means of callus relative growth rate derived from leaf explants of sugarcane after treated with different concentrations of mannitol.	61
Figure (8): Effect of different concentrations of mannitol (a) 0, (b) 100, (c) 200 and (d) 300 mM on callus relative growth rate of GT 54-9 genotype.	62
Figure (9): Effect of different concentrations of mannitol (a) 0, (b) 100, (c) 200 and (d) 300 mM on callus relative growth rate of G84-47 genotype.	63
Figure (10): Effect of different concentrations of mannitol (a) 0, (b) 100, (c) 200 and (d) 300 mM on callus relative growth rate of ph8013 genotype	64
Figure (11): Percentage of water content for sugarcane genotypes.	65
Figure (12): Percentage of callus water content of sugarcane after treated with different concentrations of mannitol.	65
Figure (13): Effect of different concentrations of mannitol (a) 0, (b) 100, (c) 200 and (d) 300 mM on percentage of callus water content .	66
Figure (14): K ⁺ and Na ⁺ ions determination for sugarcane genotypes.	70
Figure (15): Means of K ⁺ in callus derived from leaf explants of sugarcane after treated with different concentrations of mannitol.	70
Figure (16): Means of Na ⁺ in callus derived from leaf explants of sugarcane after treated with different concentrations of mannitol.	71
Figure (17): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in GT 54-9 callus.	74

Figure (18): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in G 84-47 callus.	75
Figure (19): Regression between mannitol concentrations (X-factor) and the proline (Y-factor) in ph 8013 callus.	76
Figure (20): Morphological performance of sugarcane genotypes (No. SF: number of shoot formation; No. DRF: day numbers for root formation; NO. R: Root number).	80
Figure (21): Interaction of shoot number formation of sugarcane genotypes (No. SF: number of shoot formation)	80
Figure (22): Regeneration of GT54-9 obtained from tolerant callus under different mannitol concentrations.	81
Figure (23): Regeneration of ph 80-13 obtained from tolerant callus under different mannitol concentrations.	82
Figure (24): Day number of root formation for three sugarcane genotypes.	83
Figure (25): Root number formation for three sugarcane genotypes.	83
Figure (26): Difference in number of root formation for GT-54-9 under different mannitol concentration.	84
Figure (27): Difference in number of root formation for G84-47 under different mannitol concentration.	85
Figure (28): Dendrogram of the three cultivars and using five random amplified (RAPD) primers (54: GT 54-9, 80: ph 8013, 84:G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	97

LIST OF PLEATS

	<u>Title</u>	page
Plate (1):	DNA polymorphism of the Sugarcane (<i>Saccharum officinarum</i> L.) treated with mannitol using randomly amplified polymorphic DNA with primer OPD-02. (54: GT 54-9, 80: ph 8013, 84: G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol)	86
Plate (2):	DNA polymorphism of the Sugarcane (<i>Saccharum officinarum</i> L.) treated with mannitol using randomly amplified polymorphic DNA with primer OPD-03. (54: GT 54-9, 80: ph 8013, 84: G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	88
Plate (3):	DNA polymorphism of the Sugarcane (<i>Saccharum officinarum</i> L.) treated with mannitol using randomly amplified polymorphic DNA with primer OPH-03 (54: GT 54-9, 80: ph 8013, 84: G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	90
Plate (4):	DNA polymorphism of the Sugarcane (<i>Saccharum officinarum</i> L.) treated with mannitol using randomly amplified polymorphic DNA with primer OPO-01. (54: GT 54-9, 80: ph 8013, 84: G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	92
Plate (5):	DNA polymorphism of the Sugarcane (<i>Saccharum officinarum</i> L.) treated with mannitol using randomly amplified polymorphic DNA with primer OPO-02 (54: GT 54-9, 80: ph 8013, 84: G 84-47, M: marker, C: control, I: 100mM, II: 200mM and III: 300mM mannitol).	94

Acknowledgement

First and foremost, I feel always indebted to Allah, the most beneficent and merciful, god provided me with strength to accomplish this investigation.

I would like to express my deep thanks and gratitude to **Prof. Dr. Ahmed Elsayed Khaled**, Professor of Genetics, Department of Botany, Faculty of Agriculture, Saba-Basha, Alex., Univ., for his real help and for his valuable advice. I wish for him more success and all happiness.

Faithful thanks to **Prof. Dr. Nabawya Saleh Abdo Ghura**, Head of Research, Agricultural Research Centre, Sugar Crops Research Institute, for her valuable advices and constant encouragement to complete the present work under his supervision.

I am highly obliged to **Dr. Nader Ragab Abdelsalam**, Assistant professor of plant genetic, Agricultural Botany Dept., Faculty of Agriculture, Saba-Basha, Alex., Univ., for his constructive criticism during the revision of the thesis, for his real help and for his valuable advice. he offered me throughout this work, I wish acceptance of his prayers for anything he wishes for.

I honestly lack words to express my appreciation **Prof. Dr. Mona massoud**, Head of Research, Agricultural Research Centre, Sugar Crops Research Institute, for her patience, limitless help and the time and effort she dedicated. I wish her more success, happiness and acceptance of her prayers for anything she wishes for.

I would like to record my almost appreciation to **Dr. Mohamed Abdel-Monem Ghonima**, Senior researcher, Agricultural Research Center, Sugar Crops Research Institute, Sabbaheya Research Station, Alex., for his supervision and sincere guidance.

My appreciation to my dear friend **Dr. Sara Emad El-Din Gomma**, Researcher, Agricultural Research Center, Horticultural Research Institute, Sabbaheya Research Station, Alex., for her support in statistical regression analysis and encouragement.

Special thanks to **Dr. Amr Mohamed El-Sheikh**, Researcher and all my colleagues in El-Sabahya Research Branch, for their assistance and valuable cooperation.

Finally special thanks to my **mother, father, sisters, my husband Mohamed, sons Youssef and Yahia and my sweetly daughter Mariam.**

ABBREVIATIONS

%	: Percentage
μl	: Micro liter
μM	: Micro molar
2,4-D	: 2,4- Dichlorophenoxy Acetic Acid
AC	: Activated Charcoal
bp	: Base Pair
Cv(s)	: Cultivar(s)
CW	: Coconut Water
DW	: Dry Weight
EC	: Embryogenic Callus
FW	: Fresh Weight
hrs	: Hours
K ⁺	: Potassium
Kin	: Kinetin
M	: Molar
Mg l ⁻¹	: Milligram per liter
ml	: Milliliter
mM	: Milli molar
MS	: Murashige and Skoog
Na ⁺	: Sodium
NAA	: 1-Nphthylacetic acid
NaOH	: Sodium Hydroxide
NO.DRF	: Number of Day for Root Formation
NO.R	: Number of Root
NO.SF	: Number of Shoot Formation
°C	: Degree Celsius
PCR	: Polymerase Chain Reaction
PEG	: Poly Ethylene Glycol
pH	: - Log [H ⁺]
PVP	: Poly Vinyl Pyrrolidone
RAPD	: Random Amplified Polymorphic DNA
RGR	: Relative Growth Rate
SE	: Somatic Embryogenesis
WC	: Water Content