# LIST OF FIGURES


| Figure | | Page |
|---|---|---|
| **(1)** | Abdominal wall contusion in a handlebar injury in 10 years child | **1** |
| **(2)** | The intra-abdominal organs | **3** |
| **(3)** | Hepatic injuries | **4** |
| **(4)** | Coronal reformation of contrast-enhanced CT scan through upper abdomen shows complex hepatic laceration | **4** |
| **(5)** | Contrast-enhanced CT scan through upper abdomen shows laceration extending to periphery of liver with associated subcapsular hematoma | **4** |
| **(6)** | Grade 2 splenic injury | **5** |
| **(7)** | Contrast-enhanced CT scans through upper abdomen showing shattered spleen | **5** |
| **(8)** | The types of kidney injuries | **6** |
| **(9)** | Contrast enhanced CT scan through kidneys shows show a right upper pole renal laceration | **6** |
| **(10)** | Contrast-enhanced CT scan through upper abdomen shows bowel rupture with extra-luminal air | **7** |
| **(11)** | Jaw thrust maneuver, (right) Chin lift maneuver | **8** |
| **(12)** | Electrical pulses from the ultrasound machine make the transducer create sound waves at the desired frequency | **15** |
| **(13)** | Types of scanners which are used in abdominal sonography | **15** |
| **(14)** | Linear array probe | **17** |
| **(15)** | Curvilinear array probe | **17** |
| **(16)** | FAST examination views | **17** |
| **(17)** | FAST-oriented algorithm | **19** |
| **(18)** | Positive FAST scan showing blood anterior to the right kidney | **20** |
| **(19)** | Positive FAST scan showing blood around the spleen | **20** |
| **(20)** | Coronal section of CT abdomen with I.V contrast showing grade V splenic injury. These arrows demonstrate the possible pathway of blood | **22** |
| **(21)** | Sagittal section of CT abdomen with I.V contrast showing haemoperitoneum from liver injury The 'sentinel clot' sign is seen as a high-attenuation collection adjacent to the liver surface | **22** |
| **(22)** | Plain erect x-ray abdomen showing pneumoptonium | **23** |
| **(23)** | Toshiba Nemio 20 Ultrasound device | **28** |
| **(24)** | Ultrasound probes | **28** |

| Figure | | Page |
|---|---|---|
| **(25)** | Phillips MX 16 slice computerized tomographic scanner device | **29** |
| **(26)** | Distribution of cases according to age group | **35** |
| **(27)** | Area chart showing distribution of age frequency | **35** |
| **(28)** | Bar graph showing distribution of age groups compared to gender | **36** |
| **(29)** | Distribution of cases according to mode of trauma | **37** |
| **(30)** | Relation between age and MOT | **38** |
| **(31)** | Distribution of cases according to grading of hypovolemic shock | **39** |
| **(32)** | Distribution of cases according to abdominal examination | **40** |
| **(33)** | Distribution of the studied cases according to patient outcome | **41** |
| **(34)** | Relation between abdominal examination and patient outcome | **42** |
| **(35)** | Distribution of the studied cases according to radiological investigations | **44** |
| **(36)** | Percentage distribution of FAST finding | **45** |
| **(37)** | Distribution of the studied cases according to FAST site of collection | **46** |
| **(38)** | Relation between Grade of shock and FAST fluid grade | **47** |
| **(39)** | Relation between FAST fluid grade and patient outcome | **48** |
| **(40)** | Relation between CT and patient outcome | **50** |
| **(41)** | FAST Vs CT regarding timing | **51** |
| **(42)** | Comparison between FAST and CT according to grade of collection | **52** |
| **(43)** | Comparison between FAST and CT according to organ injury | **54** |

# LIST OF ABBREVIATIONS

| | |
|---|---|
| **AA** | Alleged Assault |
| **ABCDE** | Airway, Breathing, Circulation, Disability and Environment &Exposure |
| **A&E** | Accident and Emergency department |
| **ALARA** | As Low As Reasonably Achievable |
| **ALT** | Alanine AminoTransferase |
| **APLS** | Advanced Pediatric Life Support |
| **AST** | Aspartate AminoTransferase |
| **ATLS** | Advanced Trauma Life Support |
| **AVPU** | Alert, respond to Verbal, respond to Pain, Unresponsive |
| **BAT** | Blunt Abdominal Trauma |
| **CBC** | Complete Blood Count |
| **CI** | Confidence Intervals |
| **CT** | Computed Tomography |
| **CVC** | Central Venous Catheter |
| **CVP** | Central Venous Pressure |
| **DAMA** | Discharged Against Medical Advice |
| **DPL** | Diagnostic Peritoneal Lavage |
| **DL** | Diagnostic Laparoscopy |
| **ED** | Emergency Department |
| **EFAST** | Extended Focused Assessment with Sonography for Trauma |
| **FAST** | Focused Assessment with Sonography for Trauma |
| **FE** | Fisher Exact test |
| **FFH** | Falling From Height |
| **GCS** | Glasgow Coma Scale/Score |
| **HU** | Housnfields Units |
| **IAI** | Intra-Abdominal Injuries |
| **ICU** | Intensive Care Unit |
| **IV** | Intra- Venous |

| | |
|---|---|
| **KHz** | Kilo Hertz |
| **LUQ** | Left Upper Quadrant |
| **MC** | Monte Carlo test |
| **MHz** | Mega Hertz |
| **ml/k.g** | milliliter per kilogram |
| **mm Hg** | millimeter mercury |
| **MOT** | Mode Of Trauma |
| **No.** | Number |
| **NPV** | Negative Predictive Value |
| **OR** | Operating Room |
| **PPV** | Positive Predictive Value |
| **RTA** | Road Traffic Accident |
| **RUQ** | Right Upper Quadrant |
| **SAMPLE** | Signs and symptoms, Allergy, Medications, Past medical history, Last meal and Events related and environment |
| **SD** | Standard Deviation |
| **t** | student t-test |
| **TBI** | Traumatic Brain Injury |
| **U/S** | Ultrasound |
| **$\chi 2$** | value for Chi square test |