

LIST OF FIGURES

Figure	Page
1. The cardiac diamond	2
2. Suggested algorithm for traumatic cardiac arrest	3
3. Traumatic cardiac arrest treatment algorism	4
4. Chest radiographs. A: Initial supine chest radiograph a trauma patient. B: Repeat chest X-ray study 22 h later demonstrating significant increased density involving the right hemithorax, believed to represent mediastinal blood.	6
5. Heamothorax at the left side with mediastinal shift to the right side	7
6. FAST scan showing echo-free space around the heart consistent with haemopericardium	8
7. Transthoracic four-chamber echocardiographic view showing the large pericardial fluid (marked with arrow).	9
8. Large pericardial effusion at CT scan.	9
9. J wave on the ECG (J-wave indicated by the arrow).	10
10. Indications for thoracotomy	11
11. Antero-lateral thoracotomy, opening of the pericardium to relieve tamponade by incising the pericardium anterior to the phrenic nerve.	12
12. A: An instrument passed through the aortic root emerges in the right ventricular outflow tract, confirming the presence of a defect in the fibrous outlet ventricular septum, B: associated with a perforation of the right coronary leaflet of the aortic valve and C: a perforation of the anterior mitral leaflet. This lesion pattern reflects the trajectory of a single anterolateral to posteromedial cardiac stab injury.	14
13. A: Percentages of the patients according to systolic blood pressure and the mechanism of injury B: Percentages of the patients according to pulse and the mechanism of injury C: Percentages of the patients according to respiratory rate and the mechanism of injury D: Percentages of the patients according to Glasgow coma scale and the mechanism of injury E: Percentages of the patients according to central venous pressure and the mechanism of injury	22-24

Figure	Page
<p>14. A: Percentages of the patients according to systolic blood pressure and the age 27-29 B: Percentages of the patients according to pulse and the age C: Percentages of the patients according to respiratory rate and the age D: Percentages of the patients according to Glasgow Coma Scale and the age E: Percentages of the patients according to central venous pressure and the age</p>	
<p>15. A: Percentages of the patients according to clinical changes on neck veins and the mechanism of injury 30-31 B: Percentages of the patients according to chest auscultation findings and the mechanism of injury C: Percentages of the patients according to other associated injuries and the mechanism of injury</p>	
<p>16. A: Percentages of the patients according to clinical changes on neck veins and the age 33-34 B: Percentages of the patient according to chest auscultation findings and the age C: Percentages of the patients according to other associated injuries and the age</p>	
<p>17. A: Percentages of the patients according to pre-operative focused assessment with sonography for trauma and the mechanism of injury 36-38 B: Percentages of the patients according to pre-operative electrocardiogram and the mechanism of injury C: Percentages of the patients according to pre-operative CT-Chest and the mechanism of injury D: CT-Chest of patients with penetrating cardiac injury by gunshot and stab E: Percentages of the patients according to pre-operative Chest X-ray and the mechanism of injury F: Chest X-ray of patients with penetrating cardiac injury by gunshot and stab</p>	
18. Arterial blood gases interpretation	39
<p>19. A: Percentages of the patients according to pre-operative focused assessment with sonography for trauma and the age 41-42 B: Percentages of the patients according to pre-operative CT-Chest and the age C: Percentages of the patients according to pre-operative Chest X-ray and the age</p>	

Figure	Page
	D: Percentages of the patients according to pre-operative electrocardiogram and the age
20.	44
	Percentages of the patients according to types of management and the mechanism of injury
21.	45
	Percentages of the patients according to types of management and the age
22.	46
	Percentages of the patients according to the time spent before going to operating theater and the mechanism of injury
23.	47
	Percentage of the patients according to the time spent before going to operating theater and the age
24.	48
	Percentage of internal cardiac injuries of the patients according to the mechanism of injury
25.	49
	Percentage of internal cardiac injuries of the patients according to the age
26.	51
	Percentage of intra-abdominal injuries of the patients according to the age
27.	53
	Amount of the blood transfusion and blood loss according to the age
28.	57-60
	A: Effect of age on mortality
	B: Effect of Mechanism of injury on mortality
	C: Effect of Systolic blood pressure on mortality
	D: Effect of Glasgow coma scale on mortality
	E: Effect of Other associated injuries on mortality
	F: Effect of Time spent before going to operating theater on mortality
	G: Effect of Tamponade on mortality
	H: Effect of Intra-abdominal injuries on mortality
29.	61-62
	A: ECG shows sinus tachycardia
	B: ECG shows normal sinus rhythm
30.	62
	Echo show minimal pericardial effusion

LIST OF ABBREVIATIONS

ABG	Arterial Blood Gases
ALS	Advanced Life Support
AMUH	Alexandria Main University Hospital
CPA	Cardio-Pulmonary Arrest
CPR	Cardio-Pulmonary Resuscitation
CT	Multi slice Computed Tomography
CVC	Central Venous Catheter
CVP	Central Venous Pressure
ECG	Electrocardiogram
Echo	Echocardiogram
ED	Emergency Department
ET	Emergency Thoracotomy
FAST	Focused Assessment Sonography for Trauma
GCS	Glasgow Coma Scale
ICU	Intensive Care Unit
ISS	Injury Severity Score
PHC	Pre Hospital Care
RR	Respiratory Rate
RTA	Road Traffic Accidents
RTS	Revised Trauma Score
SBP	Systolic Blood Pressure
SPW	Sub-xiphoid Pericardial Window
TCA	Trauma Cardiac Arrest
TEE	Trans-esophageal Echocardiogram
TRISS	Trauma Revised Injury Severity Score
VSD	Ventricular Septal Defect