

## **CHAPTER 6**

## **REFERENCES**

## 6. REFERENCES

- Abbott, W.S. (1925).** A method of computing the effectiveness of an insecticide. J. Econ. Entomol., 18: 265-267.
- Abdel-Mallck, A. Y.; M. A. A. Abdel-Rahman and G. R. A. Hamam (2003).** Survey of entomopathogenic fungi naturally infecting cereal aphid (Homoptera Aphididae) in Southern Egypt. (Abst.). 9<sup>th</sup> European meeting of the IOBC/WPRS working Group, Insect pathogens and Entomoparasitic nematodes in cooperation with cost actions 842 and 850 "Entomophothorales & Biocontrol Symbiosis" "Growing research and development, Schlos Salzau, Germany, 23-29 May, 2003 .
- Abd-Rabou, S.; N. Ahmed and H. Badary (2012).** Scale insects (Hemiptera: Coccoidea) infesting apple, apricot, pear trees and their abundant parasitoids in Egypt. Acad. J. Biol. Sci., 5(1): 25-29.
- Abid, H.; M. Yi Tina; Yu. He; L. Ruan and S. Ahmed (2010).** *In vitro* and *in vivo* culturing impacts on the virulence characteristics of serially passed entomopathogenic fungi. J. Food, Agric. Environ., 8 (3& 4): 481 - 487.
- Adames, M.; M. Fernández; G. Peña and V. Hernández (2011).** Effects of passages through a suitable host of the fungus, *Metarhizium anisopliae*, on the virulence of acaricide-susceptible and resistant strains of the tick, *Rhipicephalus microplus*. J. Insect Sci., 21: 1-13.
- Ajaykumar, P. and K. R. Kanaujia (2005).** Effect of different grain media on sporulation, germination and virulence of *Beauveria bassiana* against *Spodoptera litura* larvae. J. Biol. Control, 19(2): 129-133.
- Ali, A.; C. G. Rojas and M. Torre (2005).** Isolation of dipicolinic acid as an insecticidal toxin from *Paecilomyces fumosoroseus*. J. Appl. Microbiol. Biotechnol., 68: 542– 547.
- Altre, J. A.; J. D. Vandenberg and F. A. Cantone (1999).** Pathogenicity of *Paecilomyces fumosoroseus* isolates to diamondback moth, *Plutella xylostella*: correlation with spore size, germination speed, and attachment to cuticle. J. Invert. Pathol., 73: 332-338.
- Alves, R. T.; P. R. Bateman; J. Gunn; C. Prior and S. R. Leather (2002).** Effects of different formulation of viability and medium term storage of *Metarhizium anisopliae* conidia. Neotropical Entomol., 31 (1): 91-99.

- Amala, U.; T. Jiji and A. Naseema (2012).** Mass multiplication of entomopathogenic fungus, *Paecilomyces lilacinus* (Thom) Samson with solid substrates. J. Biopest., 5(2): 168-170.
- Anitha, K. R. ( 2007).** Seasonal incidence and management of sucking pest of okra. M. Sc.(Agric) Thesis, Univ. Agric. Sci., Dharwad, Karnataka, India.
- Anonymous (2005).** Efficient input management. The Hindu survey, Indian Agriculture. pp 47-49.
- Ansari, M. A. and T. M. Butt (2011).** Effects of successive subculturing on stability, virulence, conidial yield, germination and shelf-life of entomopathogenic fungi. J. Appl. Microbiol., 110: 1460-1469.
- Asghar, M. (2013).** Virulence of *Beauveria bassiana* and *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) passaged through artificial media and an insect host *Uvarovistia zebra* (Orthoptera: Tettigoniidae), Int. J. Agric Crop Sci., 16: 1147-1152.
- Ayyasamy, R. and P. Baskaran (2006).** Evaluation of various synthetic and grain media for radial growth, sporulation and infectivity of the entomopathogenic fungi *Paecilomyces farinosus*. J. Ecobiol., 19 (5) : 229-233.
- Banu, J. G. and N. Gopalakrishanan (2012).** Development of formulations of a native entomopathogenic fungus, *Lecanicillium lecanii* and testing virulence against mealybug, *Paracoccus marginatus* infesting cotton. Indian J. Plant Prote. 40 (3): 182- 186
- Barranco-Florido, J. E.; R. Alatorre-Rosas; M. Gutierrez-Rojas; G. Viniegra-Gonzalez and G. Saucedo-Castaneda (2002).** Criteria for the selection of strains of entomopathogenic fungi *Verticillium lecanii* for solid state cultivation. Enzyme Microb. Technol., 30:910-915.
- Bekheit, H. K. M. (2005).** Biological control tools and their use in insect mangement in Egypt. Egypt. J. Agric. Res., 83(3):1249-1288.
- Bhagat, R. M.; R. B. L. Gupta and C. P. S. Yadava (2003).** Field efficacy of two entomopathogenic fungal formulations against whitegrubs in Himachal Pradesh. Indian J. Entomol., 65 (1): 76-81.
- Bhanu, P. B.; S. Puri and P. K. Singh (2012).** Mass production of entomopathogenic fungi using agricultural products. J. Life Sci., 7(2): 229-232.
- Bhattacharyya, A.; A. C. Samal and S. Kar (2004).** Entomophagous fungus in pest management . News Letter, 5: 2-9

- Boucias, D. G. and J. C. Pendland (1998).** Principles of insect pathology. Kluwer Academic Publishers, 537 p.
- Brownbridge, M.; A. Adamowicz; M. Skinner and B. L. Parker (1999).** Prevalence of fungal entomopathogens in the life cycle of pear thrips, *Taeniothrips inconsequens* (Thysanoptera : Thripidae) in Vermont sugar maple forests. Biol. Control, 16(1): 54-59.
- Brownbridge, M.; S. Costa and S. T. Jaronski (2001).** Effects of *in vitro* passage of *Beauveria bassiana* on virulence of *Bemisia argentifolii*. J. Invert. Pathol., 77: 280- 283.
- Burges, H. D. (1981).** Microbial Control of Pests and Plant Diseases :1970-1980. Academic Press, London. 949 pp.
- Burgess, H. D. and N.W. Hussey (1971).** Microbial control of pests and plant Diseases, 1970-1980, Academic press, London.
- Burges H. D. and N.W. Hussey (1981).** Microbial Control of Insect Pests and Mite, Academic Press, London, pp. 161-167.
- Butt, T. M; M. Goettel (2000).** Bioassays of entomopathogenic fungi. In: Bioassays of entomopathogenic microbes and nematodes. Navon, A. and K. R. S. Ascher (editors). pp. 141-195. CAB International.
- Butt, T. M.; C. Wang; F. A. Shah and R. Hall (2006).** Degeneration of entomogenous fungi. In: An ecological and societal approach to biological control. Eilenberg, J. H. M. T. Hokkanen (editors). pp. 213–226. Springer
- Carruthers, R. I. and R. S. Soper (1987).** Fungal diseases. In: Epiizootiology of Insect Diseases. Fuxa, J. R. and Tanaday (eds) Wiley. Interscience, New York, pp. 357-416.
- Chandrashekharaiyah, G.; B. Gowda; A. Kumar and A. K. Chakravarthy (2013).** Bio-efficacy of *Verticillium lecanii* against white fly on greengram Insect Environ., 19(2):83-85.
- Cooper, R. and A.W. Sweeney (1986).** Laboratory studies on the recycling potential of the mosquito pathogenic fungus *Culicinomyces clavisporus*. J. Invert. Pathol., 48: 152–158.
- Dai, Y.; H. Hiromori and M. Hatsukade (2006).** Virulence of the entomopathogenic fungus *Beauveria brongniartii* to several life stages of the yellowish elongate chafer *Heptophylla picea* Motschulsky (Coleoptera: Scarabaeidae). Appl. Entomol. Zool., 41 (2): 287–293 .
- Daust, A. R.; G. M. Ward and W. D. Roberts ( 1983).** Effect of formulation on the viability of *Metarhizium anisopliae* conidia. J. Invert. Pathol., 41:151-160.

- Dent, D. (1991).** Insect Pest Management C.A.B International. ISBB No-85198-666-8) 604 pages.
- Derakhshan, A.; R. J. Rabindra; B. Ramanujam and M. Rahimi (2008).** Evaluation of different media and methods of cultivation on the production and viability of entomopathogenic fungi, *Verticillium lecanii* (Zimm.) Viegas. Pakistan J. Biol. Sci., 11: 1506-1509.
- Derakhshan, A.; R. J. Rabindra and B. Ramanujam (2008).** Effect of storage conditions of formulations on viability of *Verticillium lecanii* (Zimm.) Viegas and its virulence to *Brevicoryne brassicae* (L.). Pakistan J. Biol.Sci., 8 (2): 498-501.
- De-Souza, L.; N. C. De-Oliveira; C. F.Wilcken and A. B. Filho (2004).** Pathogenicity of *Verticillium lecanii* to pine aphid. Rev . Arvore, 28 (5): 765-770.
- Dreistadt, S. H.; J. K. Clark and M. L. Flint (1994).** Pests of landscape trees and shrubs: An integrated pest management guide. Oakland: Univ. Calif. Div. Agric. Nat. Res., 3359.
- Duncan, D. B. (1955).** Multiple range and multiple F tests. Biometrics, 11:1-44.
- Easwaramoorthy, S and S. Jayaraj (1978).** Effectiveness of the white halo fungus, *Cephalosporium lecanii*, against field populations of coffee green bug, *Coccus viridis*. J. Invert. Pathol., 32:88-96.
- Easwaramoorthy, S.; J. Srikanth; G. Santhalakshmi and N. Geetha (2003).** Mass culture and formulation of three entomogenous fungi, with special reference to *Beauveria brongniartii* (Sacc.) petch, against *Holotrichia serrata* F. (Coleptera : Scarabaeidae). Proceedings of the 64<sup>th</sup> Annual Convention of the Sugar Technologists Association of India, Cochin, Kerala, 17<sup>th</sup>-19<sup>th</sup> August, 2002.
- Ekesi, S.; N. K. Maniania; I. Onu and B. Lohr (1998).** Pathogenicity of entomopathogenic fungi (Hyphomycetes) to the legume flower thrips, *Megalurothrips sjostedtoi* (Trybom) (Thysanoptera : Thripidae). J. Appl. Entomol., 122 (9-10): 629-634.
- Eliana, T. I.; G. V. Pereira; D. T. Miyagui; M. H. P. Pinotti and P. M. O. J. Neves (2007).** Production of extracellular protease by a Brazilian strain of *Beauveria bassiana* reactivated on coffee berry borer *Hypothenemus hampei*. Brazilian Arch. Biol. Technol. J., 50(2):217-223.
- El-Serafi, H. A.; A. A. Ghanim; A. H. El-Heneidy and M. K. El-Sherbenie (2004).** Ecological studies on certain insects infesting guava orchards and their predatory insects at Mansoura District. Egypt. J. Biological Pest Control, 14: (1) 77-85.

- El-Sinary, N. H. and S. A. Rizk (2007).** Entomopathogenic fungus, *Beauveria bassiana* (Bals.) and gamma irradiation efficiency against the greater wax moth, *Galleria melonella* (L.). American-Eurasian J. Sci. Res., 2 (1): 13-18.
- Esfandiari, M.; M. S. Mossadegh and R. Eslamizadeh (2007).** Biology of the cottony cushion scale, *Icerya purchasi* Mask. (Hom. : Margarodidae) in laboratory and its seasonal fluctuations in citrus orchards of Northern Khuzestan. J. Sci. Tech. Agric. Nat. Res., 10 (4B): 393-403.
- Evans, H. C. (1974).** Natural control of arthropods, with special reference to ants (Formicidae), by fungi in The Tropical High Forest of Ghana. J. Appl. Eco. , 11:37-49.
- Evans, H. C. (1999).** Biological control of weed and insect pests using fugal pathogen, with particular reference to Srilanka. Bio control News and Information, 20 (2): 63 -68.
- Ezzat, Y. M. and S. M. A. Nada (1986).** List of super family Coccoidea as known to exist in Egypt. Ball. Lab. Entomol. Agric. 43: 85-90.
- Ezzati-tabrizi, R.; T. Reza, P. Hamid-reza (2009).** Effect of formulating of *Beauveria bassiana* conidia on their viability and pathogenicity to the onion thrips, *Thrips tabaci* Lind. (Thysanoptera: Thripidae) J. Plant Protec. Res., 49 (1):97-104.
- Fadayivata, S.; F.Moravvej and J. Karimi (2014).** Pathogenicity of the fungus *Lecanicillium longisporum* against *Sipha maydis* and *Metopolophium dirhodum* in laboratory conditions, J. of Plant Protec. Res., 54: (1): 67-73.
- Fargues, J. F. and P. H. Robert (1983).** Effect of passaging through scarabaeid hosts on the virulence and host specificity of two strains of the entomopathogenic hyphomycete *Metarhizium anisopliae*. Can. J. Microbiol., 29: 575 – 583.
- Farooq A. S.; C. S. Wang and T. M. Butt (2005).** Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. FEMS Microbiol. Letters, 251: 259–266.
- Farsi, M. J.; H. Askari; K. T. Jahromi and A. K. Pakdel, (2005).** Effect of important ecological factors on blastospore production of *Verticillium lecanii* DAOM 198499 in liquid medium. Iranian J. Agric. Sci., 36: 109-119.
- Ferron, P. (1977).** Influence of relative humidity on the development of fungal infection caused by *Beauveria bassiana* (Fungi imperfecti, Moniliales), in imagines of *Acanthoscelides obtectus* (Coleoptera: Bruchidae). Entomophega, 22: 393-396.

- Ferron, P. (1985).** Fungal control. *In: Comprehensive Insect Physiology, Biochemistry and Pharmacology*, G.A. Kerkut and L.I. Gilbert [eds.], pp 313-316., Academic, New York.
- Finney, D. N. (1971)** Probit analysis. 3<sup>rd</sup> Ed, Cambridge Univ. Press London pp: 318.
- Forschler, B. T. and G. L. Nordin (1989).** Impact of *Beauveria bassiana* on the cotton wood borer *Plectrodera sclator* (Coleoptera : Cerambycidae), in a commercial cotton wood nursery. J. Entomol. Sci., 24: 186-190.
- Fransen, J. J.; K. Winkelman and J. C. Van Lenteren (1987).** The differential mortality at various life stages of the greenhouse whitefly, *Trialeurodes vaporariorum* ( Homoptera: Aleyrodidae), by infection with the fungus *Aschersonia aleyrodis*( Deuteromycotina: Coelomycetes). J. Invert. Pathol., 50: 158-165.
- Fumio, I.; K. Yaginuma; N. Kobayashi; K. Mishiro and T. Sato (2001).** Screening of entomopathogenic fungi against the brown-winged green bug, *Plautia stali* Scott (Hemiptera: Pentatomidae) Appl. Entomol. Zool., 36 (4): 495–500 .
- Fumio, I.; M. Toyama and T. Sato (2003).** Pathogenicity of *Metarhizium anisopliae* to the chestnut weevil larvae under laboratory and field conditions. Appl. Entomol. Zool., 38 (4): 461–465.
- Fuxa, J. R. (1987).** Ecological considerations for the use of entomopathogens in IPM. Ann. Rev. Entomol., 32: 225-251.
- Ganga Visalakshy, P. N.; A. Manoj kumar and A. Krishnamoorthy (2004).** Evaluation of entomopathogenic fungi *Verticillium lecanii* against *Scirtothrips dorsalis*. Project Directorate of Biological Control, Bangalore, Annu. Progr Repo : 2004-2005, pp. 125-126.
- Garcia, M. F. (2006).** Phytosanitary status of citrus in Spain: Insects and mites. Informatore Fitopatologico, 56(1): 28-31.
- Gerlach, J. (2010).** Margarodidae (Hemiptera: Insecta) of the Seychelles islands. *Phelsuma* 18: 70-73.
- Gindin, G.; N.U.Geschtovt; B. Raccah and I. Barash (2000).** Pathogenicity of *Verticillium lecanii* to different developmentl stages of the silverleaf whitefly, *Bemisia argentifolii*. Phytoparasitica, 28 (3): 1-11.
- Godonou, I.; K. R. Green; K. A. Oduro; C. J. Lomer and K. Afreh-Nuaman (2000) .** Field evaluation of selected formulations of *Beauveria bassiana* for the management of the banana weevil

- (*Cosmopolites sordidus*) on plantain (*Musa* spp.). International Symposium on Biological Control Agents in Crop and Animal Protection, Swansea, U. K. 24-28 August 1999. Bio control Sci. & Technol., 10 (6): 779-788.
- Goettel, M. S., G. D. Inglis and S. P. Wraight (2000).** Fungi *In: field manual technique in invertebrate pathology*, L.A. lacey and H.K. kaya (eds.) Kluwer Academic publisher , NetherLands ., pp.255-282 .
- Gopalkrishnan, C. and K. S. Mohan ( 2000).** A simple and cost effective *in vitro* method for mass production of conidia of *Nomuraea rileyi*. Pest Mgt. Hort. Ecosyst., 6: 36-39.
- Green, E. E. (1907).** Notes on the Coccidae collected by the Percy Sladen Trust expedition to the Indian Ocean: supplemented by a collection received from Mr R. Dupont, Director of Agriculture, Seychelles. Trans. Linn. Soc. London, 2 (12): 197-207.
- Guedes-Frazzon, A. P.; I. Vaz Junior; A. Masuda; A. Schrank and M. Henning (2000).** *In vitro* assessment of *Metarhizium anisopliae* isolates to control the cattle tick *Boophilus microplus*. Vet. Parasitol., 94: 117-125.
- Guillon, M. (1997).** Mass-production of biopesticides: Scale up and quality assurance. British Crop Production Council Symposium, 68. 1997, Farnham. Insecticides: novelty or necessity: proceedings. Farnham: The British Crop Protection Council., p.151 - 162.
- Gullan, P.J. and M. Kosztarab (1997).** Adaptations in scale insects. Ann. Rev. Entomol., 42: 23–50.
- Hajek, A. E.; R. A. Humber and M. H. Griggs (1990).** Decline in virulence of *Entomophaga maimaiga* (Zygomycetes: Entomophthorales) with repeated *in vitro* Subculture. J. Invert. Pathol. 56: 91-97.
- Hall, R. A. and B. Papierok (1982).** Fungi as biological control agent of arthropods of agricultural and medical importance. Parasitol., 84: 205-240.
- Hall, R. A. (1982).** Deuteromycetes : Virulence and bioassay design. *In: Invertebrate pathology and microbial control*. Proceedings III International Colloquium Invertebrate Pathology, XV Annual Medium Soc. of Invertebrate Pathology, Brighton, U.K. pp.191-196
- Hamam, G. H. (2003).** Studies of entomopathogenic fungi of cereal aphids infecting wheat plants in Assiut, Egypt. M.Sc. Thesis, Assiut Univ. Egypt..
- Hassan, N. A. and S. G. Radwan (2008).** Population dynamics of *Icerya seychellarum* (Homoptera: Margaroididae) and *Rodalia cardinalis*

(Coleoptera : Coccinellidae) on persimmon (*Diospyros kaki* ) at Qalubiya Governorate. Egypt. J. Agric. Res., 86(3):1015-1027.

**Hatting, J. L., R. A. Humber, T. J. Poprawski and R. M. Miller (1999).** A survey of fungal pathogens of aphids from South Africa, with special reference to cereal aphids. Biol. Control, 16 (1): 1–12.

**Henderson, C. F. and E. W. Tilton (1955).** Tests with acaricides against the brown wheat mite, J. Econ. Entomol., 48:157-161.

**Hedgecock, D.; P. M. Moore; P. M. Higgins and C. Prior (1995).** Influence of moisture content on on temperature tolerance and storage of *Metarhizium flavoviride* conidia in oil formulation. Biocontrol Sci.Technol., 5(3):371-377.

**Hidalgo, E.; D. Moore and G. N. Lepatourel ( 1998).** The effect of different formulations of *Beauveria bassiana* on *Sitophilus zeamais* in stored maize. J. Stored Prod. Res., 34 (2-3) : 171-179.

**Hincapie, R. H.; A. Ospina; Y. Bustillo and A. Saldarriaga ( 1990).** Evaluacion del entomopatogeno *Verticillium lecanii* en el control del afido *Mysus persicae* encrisantemos. Revista Colombiana de Entomologia, 16(2): 21-27.

**Hsia, I. C. C.; M.T. Islam; Y. Ibrahim; T. Y. Howand and D. Omar (2014).** Evaluation of conidial viability of entomopathogenic fungi as influenced by temperature and additive. Int. J. Agric. Biol., 16: 146–152.

**Ibrahim, L.; T. M. Butt and P. Jenkinson (2002).** Effect of artificial culture media on germination, growth, virulence and surface properties of the entomopathogenic hyphomycete *Metarhizium anisopiae*. Mycol. Res., 106: 705-715.

**Ignoffo, C. M.; D. L. Hostetter; P. P. Sikorowski; G. Sutter and W. M. Brooks (1977).** Inactivation of representative species on entomopathogenic virus, a bacterium, fungus and protozoa by the ultraviolet light source. Environ Entomol., 6: 411-415.

**Ignoffo, C. M.; A. H. McIntosh; C. Garcia; M. Kroha and J. M. Johnson (1982).** Effective of successive *in vitro* and *in vivo* passages on the virulence of the entomopathogenic fungus, *Nomuraea rileyi*. Entomophaga, 27: 371-378.

**Inglis, G. D.; T. M. Goettel and B. Strasser (2001).** Use of hyphomycetous fungi for managing insect pest. In: Fungi as biocontrol agentes. Butt, T.M.; Jackson, C.; Magan, N. (Ed.). Wallingford: CAB International, pp 23-69.

**Jana, S. (2009).** Influence of different storage conditions on vitality and virulence of *Beauveria bassiana* spores, J.Agrobiol., 26 (2): 75-81.

- Jaronski, S.T. (1986).** Commercial development of deuteromycetous fungi of arthropods: a criterial appraisal. *In: Fundamental and applied aspects of invertebrate pathology*. Samson, R.A., J.M. Vlask, R. Peters, (Ed.). Wageningen: Society of Invertebrate Pathology: 653-656.
- Jayaraj, S. (1989).** Integrated management of coffee green scale *Coccus viridis* (Homoptera: Coccidae). *J. Plantation Crops*, 16: 195-201.
- Jenkins, N. E.; G. Heviefo; J. Langewald; A. J .Cherry and C. J. Lomer (1998).** Development of mass production technology for aerial conidia for use as mycopesticides. *Biocontrol News and Information*, 19: 21-31.
- Jeong, J. K. and K. C. Kim (2008).** Selection of a highly virulent isolate of *Lecanicillium attenuatum* against cotton aphid. *J. Asia-Pacific Entomol.*, 11(1):1- 4.
- Kadir, A. A. and H. S. Barlow (1992).** Pest Management and the Environment in 2000. C.A.B. International, Malaysia, pp. 401.
- Karthikeyan, A. and V. Selvanarayanan (2011).** Studies on the suitability of certain culture media for *Beauveria bassiana* (Bals.) Vuill., and *Verticillium lecanii* (Zimm.) Viegas *J. Res. Biol.*, 7: 508-512.
- Kawakami, K. ( 1960).** On the change of characteristics of the silkworm muscardines through successive cultures. *Bull. Seric. Exp. Stn. Japan*. (16): 83-99.
- Khalil, S. K.; M. Kahn and M. Naeem (1990).** Studies on entomopathogenic fungi *Verticillium lecanii* (Zimm.) for the control of green pearl aphid, *Myzus persicae* Sulz. *Sarhad J. Agric.*, 6: 597-600.
- Khalil, S. K.; V. Taborsky and J. Bartos (1983).** Studies on *Verticillium lecanii* for the biological control of aphids. In 10<sup>th</sup> International Congress of Plant Protection at Brighton, England, 2: 20-25.
- Knudsen, G. R.; J. B. Johnson and D. J. Eschen (1990).** Alginate pellet formulation of a *Beauveria bassiana* (Fungi: Hyphomycetes) isolate pathogenic to cereal aphids. *J. Economic Entomol.*, 83 (6): 2225-2228.
- Krishnamurthy, P. N. M.; S. G. Eswara- Reddy ( 2004).** Survey on potential microbial agent against thrips. *Pest Management in Hortil. Ecosystem*, 7(2) : 117-123.

- Kulat, S. S.; N. N. Zade; L. N. Peshkar and S. S. Varhade (2002).** Influence of different culture media on the growth and sporulation of *Metarhizium anisopliae* (Metsch.) Sorokin. J. Biolo. Control. 16(2): 177-179.
- Kulkarni, N. S. (1999).** Utilisation of fungal pathogen *Nomuraea rileyi* (Farlow) Samson in the management of lepidopterous pests. Ph.D Thesis, Univ. Agric. Sci., Dharwad, India.
- Landa, Z. (1984).** Protection against glasshouse whitefly, *Trialeurodicus vaporariorum* in integrated protection programmes for glasshouse chambers. Sbornik Urtiz Zahoraduichini, 11: 215-218.
- Latgé, J. L.; R. A. Hall; R. I. Cabrera and J. C. Kerwin (1986).** Liquid fermentation of entomopathogenic fungi. In: Fundamental and applied aspects of invertebrate pathology. Samson, R.A., J.M. Vlcek, R. Peters (Ed.). Wageningen: Society of Invertebrate Pathology, p.603-606.
- Lawrence, A. L. and D. I. Shapiro (2003).** The potential role for microbial control of orchard insect pests in sustainable agriculture Food, Agric. Environ., 1(2): 326-331.
- Lee, J. C. and D. A. Landis (2001).** Natural enemies in field crops: A guide to biological control. Michigan State Univ. Ext. Bull., 2721.65p.
- Leena, M. D.; S. Easwaramoorthy and R. Nirmala (2003).** In vitro production of entomopathogenic fungi, *Paecilomyces farinosus* (Hottnskiold) and *Paecilomyces lilacinus* (Thom.) Samson using by products of sugar industry and other agro industrial by-products and wastes. Sugar Technol., 5 (4): 231-236.
- Lopes, E.; M. I. Lopes and N. M. Barros (1995).** Virulence of stored conidia of *Nomuraea rileyi* against soybean caterpillar, *Anticarsia gemmatilis*. Ciencia Rural, 25: 197-200.
- Machado, A.; A. C. Monteiro; A. M. B. de Almeida and M. E. Martins (2010).** Production technology for entomopathogenic fungus using a biphasic culture system Pesq. Agropec. Bras., Brasília, 45 (10):1157-1163.
- Makoto, A. and T. Ikegami (2005).** Susceptibility of five species of thrips to different strains of the entomopathogenic fungus, *Beauveria bassiana* .Appl. Entomol. Zool., 40 (4): 667-674.
- Mamta, T.; M. Thakur; N. Malik and S. Ganger (2011).** Mass scale cultivation of entomopathogenic fungus *Nomuraea rileyi* using agricultural products and agro wastes J. Biopestic., 4 (2): 176-179.
- Mangoud, A. A. H. (2000).** Integrated pest management of apple trees. Ph.D. Thesis, Fac. Agric. Cairo Univ., Cairo, Egypt
- Mangoud, A. A. H. and H. A. S. Abd El-Gawad (2003).** Evaluation of different integrated pest management concepts for controlling the

- Egyptian fluted mealybug, *Icerya aegyptiaca* on the ornamental plants. Bull. Ent. Soc. Egypt, Econ. Ser., 29 : 137-149.
- Mangoud, A. A. H.; M. A. Salem and M. A. Abd El-Aziz (2012).** Effect of different compounds on *Icerya seychellarum* (Hemiptera: Monophlebidae) and *Rodalia cardinalis* (Coleoptera: Coccinellidae) on mango leaves under laboratory conditions. Egypt. Acad. J. Biolog. Sci., 5(3): 113 -119.
- Mangoud, A. A. H.; M. S. Abd El-Wahid and M. A. Abd El-Aziz (2012).** Screening test of different compounds against *Icerya seychellarum* and harmful effect on the beneficial predator *Rodalia cardinalis* under laboratory conditions. Egypt, J. Agric. Res., 86 (5): 1861-1875.
- Marcos, R. F. and S. P. Wraight (2007).** Mycoinsecticides and Mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. Biological Control, 43: 237–256.
- Masuda, T. and O. Kikuchi (1992).** Pathogenicity of *Verticillium lecanii* isolates to whitefly and aphids. Japanese J. Appl. Entomol. Zool., 36: 239-245.
- McCoy, C. W. (1990).** Entomogenous fungi as microbial pesticides, *In: New directions in biological control*. R. R. Baker and P. E. Dunn (eds.). Alan R. Liss, New York, NY, 139-159.
- McCoy, C. W.; R. A. Samson and D. G. Boucias (1988).** Entomogenous fungi, *In: Handbook of Natural Pesticides*, Vol.5 microbial insecticides, part A, C. M. Ignoffo and N. B. Mandava (eds.). CRC Press, Inc., Boca Raton, FL, pp. 151-236.
- Meade, D. L. and D. N. Bruce (1991).** The use of *Verticillium lecanii* against submarginal instars of *Bemisia tabaci*. J. Invert. Pathol., 57: 296-298.
- Mehta, J.; D. J. Kiran; K. Ambika; S. Priya; K. Neha and D. Sakshi (2012).** Biomass Production of entomopathogenic fungi using various agro products in Kota Region, India. Res. J. Biol. Sci., 1(4):12-16.
- Mesbah, H. A.; K. S. Moursi ; A. K. Mourad; E. A. Zakzouk and R. S. Abdel-Fattah (2012).** Ecological studies on the common white mealybug, *Icerya seychellarum seychellarum* (Hemiptera: Monophlebidae) associated with *Dodonia viscosa* in Alexandria, Egypt. Egypt. Acad. J. Biolog. Sci., 5(3): 27-31.
- Midwest biological control news online (2014).**
- C.F: <http://www.entomology.wisc.edu/mbcn/kyf612.html>.

- Milner, R. J. and G. G. Lutton (1986).** Dependence of *Verticillium lecanii* on high humidities for infection and sporulation using *Myzus persicae* as host. *Environmental Entomology*, (15): 380-382.
- Miranpuri, G. S. and G. G. Khachatourians (1995).** Application of *Beauveria bassiana* and *Verticillium lecanii* against Saskatoon berry leaf aphid *Acyrtosiphon macrosiphum*. *J. Insect Sci.*, 8: 93-95.
- Mitsuaki, S. (2004).** A novel technique to inoculate conidia of entomopathogenic fungi and its application for investigation of susceptibility of the Japanese pine sawyer, *Monochamus alternatus*, to *Beauveria bassiana*. *Appl. Entomol. Zool.*, 39 (3): 485–490.
- Mitsuaki, S. (2001).** *Paecilomyces cateniannulatus* Liang, a commonly found, but an unrecorded entomogenous fungus in Japan. *Appl. Entomol. Zool.*, 36 (3): 283-288.
- Mitsuaki, S.; H. Sato and N. Maehara (2002).** Density of the entomopathogenic fungus, *Beauveria bassiana* Vuillemin (Deuteromycotina: Hyphomycetes) in forest air and soil. *Appl. Entomol. Zool.*, 37 (1): 19–26.
- Mohammad, A. H.; S. F. Moussa; A. H. Abo-Ghalia and S. A. Ahmed (2010).** Efficiency of certain insecticides on the population of the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Green) and their natural enemies under the field condition in Ismailia governorate. *Egypt. Acad. J. Biolog. Sci.*, 2 (2): 11- 17.
- Mondal, P. and A. K. Bhattacharya (2004).** Assessment of different media for mass multiplication of entomopathogenic fungus, *Beauveria bassiana* (Balsamo) Vuillemin. *Proc. Nat. Acad. Sci., India Section B, Biol. Sci.*, 74 (2): 161-169.
- Moore, D.; O. K. Douro-kpindou; N. E. Jenkins and C. J. Lomer (1996).** Effects of moisture content and temperature on storage of *Metarhizium flavoviride* conidia. *Biocontrol Sci. Technol.*, 6(1):51-61.
- Mote, U. N.; P. R. Mahajan and J. R. Kadam (2003).** Bioefficacy of *Verticillium lecanii* (Zimm.) Viegas against sucking pests of gerbera in polyhouse. *In: Green Pesticides for Insect Pest Management*. Chennai, pp. 179-183.
- Murphy, B. C.; T. Morisawa and M. P. Parrella (1998).** Insect-killing fungi. *Floriculture's IPM future Grower Talks*, 61: 60 - 68.
- Nada, M. (1990).** The scale insects (Homoptera : Coccoidea ) on citrus in Egypt . *J. Pest Cont. Environ. Sci.*, 2:201-209.
- Nada, M.; S. Abd-Rabou and G. E. Hussien (1990).** Scale insects infesting mango trees in Egypt (Homoptera: Coccoidea). *Proc. ISSIS*, VI, Part II: 133-134.

- Nagaich, B. B. (1973).** *Verticillium* species pathogenic on aphids. Indian J. Pathol. Microbiol., 26: 163-165.
- Nagaraja, S. D. (2005).** Effect of formulations of *Nomuraea rileyi* (Farlow) Samson and spray equipments in the management of tobacco caterpillar in groundnut and podborer in chickpea ecosystem. M.Sc. (Agric.) Thesis, University of Agric. Sci., Dharwad, India.
- Nahar, P.; P. Yadav; M. Kulye; A. Hadapad; M. Hassani; U. Tuor; S. Keller; A. G. Chandele; B. Thomas and M. V. Deshpande (2004).** Evaluation of indigenous fungal isolates, *Metarhizium anisopliae* M 34412, *Beauveria bassiana* B3301 and *Nomuraea rileyi* N 812 for the control of *Helicoverpa armigera* (Hub.) in pigeonpea field. J. Biol. Control, 18 (1): 1-8.
- Nankinga, C. M. and D. Moore (2000).** Reduction of banana weevil populations using different formulations of entomopathogenic fungus *Beuveria bassiana*. Biocontrol Sci. Technol., 10(5): 645-657.
- Nelson T. L.; A. Low and T. R. Glare (1996).** Large scale production of New Zealand strains of *Beauveria* and *Metarhizium*. Proceedings of the 49<sup>th</sup> New Zealand Plant Protection Conf., 257-261.
- Nier, T.; F. Rive and J. C. Bermudez (1993).** First report for Mexico on the isolation of *Verticillium lecanii* from whitefly and *in vitro* pathogenicity tests on this insect. Revista Mexicana De Micologia, 7: 149-156.
- Nirmala, R.; B. Ramanujam; R. J. Rabindra and N. S. Rao (2005).** Growth parameters of some isolates of entomofungal pathogens and production of dust free spores in rice medium. J. Biol. Control, 19 (2): 121-128.
- Nirmala, R.; B. Ramanujam; R. J. Rabindra and N. S. Rao (2006).** Effect of entomofungal pathogen on mortality of three aphid species. J. Biol. Control, 20(1): 89-94.
- Oetting, R. ( 2004).** Insect and arthropod pest identification and management. A hand-out for Southeast Greenhouse Conference workshops 2000-2004. UGA/CAES/Griffin campus. Available on-line at [http://mrec.ifas.ufl.edu/lso/Manual/insect-textspecies only.htm](http://mrec.ifas.ufl.edu/lso/Manual/insect-textspecies%20only.htm) accessed Aug. 2008 (verified 27 May 2009).
- Olga, M.; M. Kilian; E. C. Oerke and H. Wilhelmdehne (2002).** Oils for increased efficacy of *Metarhizium anisopliae* to control whiteflies, Biocontrol Sci. Technol., 12: 337- 348.
- Orozco Santo, S. M.; J. Farias-Larios; J. Lopez-Perez and N. R. Ramirezvazquez (2000).** Use of *Beauveria bassiana* for the control

of *Bemisia argentifolli* in melon. Maneho-Integrado-de-plagas, 56: 45-51.

- Osborne, L. S.; F. L. Pettitt; Z. Landa and K. A. Hoelmer (1994).** Biological control of pests attacking crops grown in protected culture: The Florida experience. *In: Pest management in the subtropics: biological control: a Florida perspective*, D. Rosen, F.D. Bennett and J.L. Capinera (eds.). Vol. 1. pp. 327-342. Intercept Limited, Andover, UK.
- Ota, M.; A. Ozawa and H. Kobayashi (1999).** Efficacy of *Beauveria bassiana* preparation against whiteflies on tomato. Annual Report of the Kanto-Tosan- Plant Protection Soc., 46: 103-112.
- Pena, J. E.; R. M. Baranowski and R. E. Litz (1987).** Life history, behavior and natural enemies of *Philephedra tuberculosa* (Homoptera: Coccidae). Florida Entomologist, 70(4):421-427.
- Pereira, R. M. and D. W. Roberts (1991).** Alginate and corn starch mycelial formulations of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae*. J.Econ. Entomol. 84(6): 1657-1661.
- Peter, W. I. and M. S. Tigano (2006).** Identification and taxonomy of some entomopathogenic *Paecilomyces* spp. (Ascomycota) isolates using rDNA-ITS Sequences Genetics and Molecular Biology, 29(1): 132-136.
- Prior, C.; P. Jollnds and G. Le Patourel (1988).** Infectivity of oil and water formulation of *B. bassiana* to the cocoa weevil pest J. Invert. Pathol., 52:62-72.
- Quesada-Moraga, E. and A. Vey (2003).** Intra-specific variation in virulence and *in vitro* production of macromolecular toxins active against locust among *Beauveria bassiana* strains and effects of *in vivo* and *in vitro* Passage on these factors .Biocontrol, Sci. Technol., 13: 323-340.
- Quinden, R. J. (1984).** The use of *Verticillium lecanii* an entomopathogenic fungus to control of glasshouse whitefly. Proceedings Brighton, England, : 9-12
- Rachappa, V. (2003).** Occurrence of entomopathogenic fungi and utilization of *Metarhizium anisopliae* in the management of selected crop pests in Northern Karnataka, Ph.D Thesis. Univ. Agric. Sci., Dharwad, India.
- Ramarethinam, S.; S. Marimuthu; W. V. Murugesan and S. Lognathan (2000).** Evaluation of *Paecilomyces fumosoroseus*, on entomopathogenic fungus for controlling red spider mite,

*Oligonychus coffeae* (Nietner) (Acarina: Tetranychidae), infesting tea in India. Pestology, 24 (9): 1-5.

**Ramarethinam, S.; S. Marimuthu and V. Murugesan (2005).** Entomopathogenic fungi *Verticillium lecanii* – an overview. Pestology, 29 (12): 9-28.

**Ramarethinam, S.; S. Marimuthu; S. Loganathan and N. V. Murgesan (2002a).** Potentials of entomopathogenic fungal based commercial formulations on some important pests of selected vegetable crops in India. Pestology, 26(7): 17-21.

**Ramarethinam, S.; S. Marimuthu; W. V. Muragesan and S. Loganathan (2002b).** Evaluation of *B. bassiana* against diamond back moth, *P. xylostella* L. (Lepidoptera: Xponomenidae). J. Entomol. Res., 25(4) : 267-271.

**Ramarethinam, S.; S. Marimuthu; N. V. Murgesan and S. R. Vishwanathan (2001).** Studies on the shelf life and container content compatibility of a commercial *B. bassiana* formulation – Bio-Power®. Pestology, 25 (91): 5-12.

**Ramegowda, G. K. (2005).** Aerobiology, Epizootiology and Utilization of *Nomuraea rileyi* (Farlow) Samson. Ph. D. Thesis, University of Agric. Sci., Dharwad, India.

**Ravensberg, W. J. (2010).** The development of microbial pest control products for control of arthropods: a critical evaluation and a roadmap to success. Ph.D Thesis Wageningen Univ., Netherlands.

**Raymond, J. S. L. S. and B. Shams-Pirzadehi (2000).** Lack of host specialization in *Aspergillus flavus* Appl. Environ. Microbial., 66(1): 320 –324.

**Reda F. A. B.; S. F. Mousa; L. S. Hamouda; R. M. Badawy and S. A. Atteia (2012).** Scale insects infesting guava trees and control measure of *Pulvinaria psidii* (Hemiptera: Coccidae) by using the alternative insecticides. Egypt. Acad. J. Biolog. Sci., 5(3): 89 -106

**Rezk, M. A. A. (2009).** Biological control of certain scale insects and mealybugs infesting certain fruit trees. M.SC. Thesis, Alexandria, Univ., Egypt.

**Roberts, D.W. (1981).** Toxins of entomopathogenic Fungi. In: Microbial control of pests and plant Diseases 1970-1980. Burgess, H.D. (ed.). Academic press, London pp 441- 84.

**Romback, M. C. (1989).** Production of *Beauveria bassinaa* conidia in submerged culture. Entomophaga, 5: 45-52.

**Rousson, S.; M. Rainbault and B.K. Lonsane (1983).** Zymotics a large scale fermenter design and evaluation. Appl. Biochem. Biotechnol., 42: 161-167 .

- Sahayaraj, K. and S. Karthick (2008).** Mass production of entomopathogenic fungi using agricultural products and by products, Afr. J. Biotechnol., 7 (12): 1907-1910.
- Salman, A. M. A. and M. M. S. Bakry (2012).** Relationship between the rate of infestation with the mealybug, *Icerya Seychellarum* (Westwood) (Margarodidae: Homoptera) and the yield loss of seedy balady mango trees at Luxor governorate. World Rural Observations, 4(4): 50-56.
- Santoro, P. H.; P. M. Neves; R. Z. Silvada; S. Akimi and J. Zorzetti (2005).** Produção de esporos de *Beauveria bassiana* (Bals.) Vuill. num processo bifásico utilizando diferentes meios líquidos. Semina Ciências Agrárias, 26: 313-320.
- Schaerffenberg, B. (1964).** Biological and environmental conditions for the development of mycoses caused by *Beauveria* and *Metarhizium*. Insect. Pathol., 6: 8-20.
- Scully, L. R. and M. J. Bidochka (2005).** Serial passage of the opportunistic pathogen *Aspergillus flavus* through an insect host yields decreased saprobic capacity. Can. J. Microbiol., 51:185-189.
- Shah, F. A.; C. S. Wang and T. M. Butt (2005).** Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. FEMS Microbiol. Lett., 51: 259 -266.
- Sharma, S.; R. B. L. Gupta and C. P. S. Yadav (1999).** Mass multiplication and formulation of entomopathogenic fungi and their efficacy against white grub. J.Mycol. Plant Pathol., 29 (3): 299-305.
- Sood, A. K.; P. K. Mehta; N. P. Kashyap and R. Lal (2001).** Bioefficacy of *Beauveria bassiana* against diamond back moth, *Plutella xylostella* L. (Lepidopera : Yponomenidae). J.Entomol. Res., 25 (4): 267-271.
- Sprenkel, R. K. and W. M. Brooks (1977).** Winter survival of the entomogenous fungus *Nomuraea. rileyi* in North Carolina. J.Invert. Pathol., 29: 262-266.
- Sreeramkumar, P.; L. Singh and H. Tabassum (2005).** Potential use of polyethylene glycol in the mass production of non-synnematous and synnematus strains of *Hirsutella thompsonii* Fisher in submerged culture. J. Biol. Control, 19 (2) : 105-113.
- Starnes, R. L; C. L. Liu and P. G. Marrone ( 1993).** History, use and future of microbial insecticides. American Entomol., 39 (2): 83-91.
- Stathers, T. E.; D. Moore and C. Prior (1993).** The effect of different temperatures on the viability of *Metarhizium flavoviride* conidia stored in vegetable and mineral oils. J. Invert. pathol., 62(2):111-115.

- Steel, G. D. and J. H. Torrie (1981).** Principles and Procedures of Statistics (2<sup>nd</sup> edition). McGraw-Hill Book Company. Inc. N. Y. 633pp.
- Steenberg, T. and R. A. Humber (1999).** Entomopathogenic potential of *Verticillium* and *Acremonium* species (Deuteromycotina: Hyphomycetes). J. Invert. Pathol., 73: 309-314.
- Steinhaus, E. A. (1949).** Principles of Insect Pathology- New york ; McGraw – Hill.
- Steinkraus, D. C.; C. J. Geden and D. A. Rutz (1991).** Susceptibility of lesser meal worm (Coleoptera: Tenebrionidae) to *Beauveria bassiana* (Moniliales: Moniliaceae): Effects of host stage, substrate, formulation and host passage. J. Med. Entomol., 28: 314-321.
- Stimac, L. J.; M. R. Pereira; B. S. Alves and A. L. Wood (1993).** Mortality in laboratory colonies of *Solenopsis invicta* (Hymenoptera : Formicidae) treated with *Beaveria bassiana* (Deutromycetes). J. Econ. Entomol., 86 (1):1083-1087.
- Suklova, T. I. (1989).** The biological method in the greenhouse. Rashichita Rastenii, 2: 37-38.
- Susumu, S. and M. Yamaji (2003).** Effect of density of the termite, *Reticulitermes speratus* Kolbe (Isoptera: Rhinotermitidae), on the susceptibilities to *Metarhizium anisopliae*. Appl. Entomol. Zool., 38 (1): 125–130.
- Tamizharasi, V.; J. Srikanth and G. Santhalakshmi (2005).** Molasses based medium requiresno nitrogen suppliment for culturing three entomopathogenic fungi. J. Biol. Control, 19(2): 135-140.
- Tawfik, M. H. and Z. K. Mohammad (2001).** Ecological studies of two scale insects (Hemiptera, Coccoidea) on *Morus alba* in Egypt. Bollettino di Zoologia Agraria e di Bachicoltura, 33 (3): 267-273.
- Tounou, K.; H. M. Agboka; K. J. poehling; G. Z. Langewald and C. Borgemeister (2003).** Evaluation of the entomopathogenic fungi *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* (Deuteromycotina:Hyphomycetes) for control of the green leafhopper *Empoasca decipiens* (Homoptera: Cicadellidae) and potential side effects on the egg parasitoid *Anagrus atomus* (Hymenoptera: Mymaridae). J. Biocont. Sci. Technol., 13(8): 715-728.
- Tsutomu, S. (2005).** Preliminary experiments to control the silverleaf whitefly with electrostatic spraying of a mycoinsecticide. Appl. Entomol. Zool., 40 (2): 289–292.

- Tsutomu, S. and K. Sugiyama (2005).** Pathogenicity of three Japanese strains of entomopathogenic fungi against the silverleaf whitefly, *Bemisia argentifolii*. Appl. Entomol. Zool., 40 (1): 169 –172.
- Unruh C. M. and P. J. Gullan (2008).** Identification guide to species in the scale insect tribe *Iceryini* (Coccoidea: Monophlebidae). Zootaxa, 1803: 1-106.
- Van Emden, H. F. (1996).** Pest Control. Combridge press 117 pp.
- Van- Hanh, V.; S. Hong and K. Kim (2008).** Production of aerial conidia of *Lecanicillium lecanii* 41185 by Solid-State. Mycobiol., 36 (3): 183-189.
- Vandenberg, J. D.; A. M. Shelton; W. T. Wilsey and M. Rmos (1998).** Assessment to *B. bassiana* sprays for control of diamond back moth (Lepidoptera: Plutellidae) on crucifers. J.Econ.Entomol., 91(3): 624-630.
- Vandenberg, J. D. and F. A. Cantone (2004).** Effect of serial transfer of three strains of *Paecilomyces fumosoroseus* on growth *in vitro*, virulence and host specificity. J. Invert. Pathol., 85: 40–45.
- Vimaladevi, P. S. (1994).** Conidia production of the entomopathogenic fungus *N. rileyi* and its evaluation for control of *Spodoptera litura* on *Ricinus communis*. J. Invert. Pathol., 63: 145-150.
- Vyas, R. V.; D. N. Yadav and R. J. Patel (1991).** Mass production of entomopathogenic fungus, *Beauveria brongniartii* on solid substrates. Indian J. Expl. Biol., 29: 795-797.
- Wasti, S. S. and G. C. Hartman (1975).** Experimental parasitization of larvae of the gypsy moth, *Porthetria dispar* (L) with the entomogenous fungus, *Beauveria bassiana* (Balsamo) Vuill. Parasitol., 70: 341-346.
- Watson, G.W. and J. Kubiriba (2005).** Identification of mealybugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. African Entomol., 13:35-47.
- Westwood, J. O. (1855).** The Seychelles Dorthesia. Gardeners' Chronicle and Agricultural Gazette 51: 836.
- Williams, D. J. and G. W. Watson (1988).** The scale insects of the tropical south pacific Region . part 2. The mealy bugs ( Pseudococcidae). CAB International Walling, Oxford, UK., 260pp
- Williams, D. J. (2004).** Mealybugs of southern Asia. The Natural History Museum, UK.
- Wiwat, C. (2004).** Development of *Nomuraea rileyi* based biopesticide for controlling Lepidopteran larvae. Ph. D. Thesis, Mahidol University, Malaysia.

- Wraight, S. P.; R. I. Carruthers; S. T. Jaronski; C. J. Bradley; C. A. Garza and S. Galani-Wraight (2000).** Evaluation of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* for microbial control of the silverleaf whitefly, *Bemisia argentifolii*. Biol. Control, 17:203-217.
- Zare, R. and W.Gams (2001).** A revision of *Verticillium* sect. *Prostrata*. III. Generic classification. Nova Hedwigia, 72: 329- 337.
- Zhang, A. W.; W. Z. Liu; X. Q. Nong; C. S. Deng; W. L. Guo and B. Jang (1992).** A trial production of wettable powder of *Beauveria bassiana*. Chinese J.Biol. Control, 8 (3): 118-120.