

CONCLUSIONS

CONCLUSIONS

- Digital radiography and Computed Tomography are valuable for evaluating pelvic fractures.
- MDCT is the first imaging study today in most poly traumatic patients.
- MDCT decrease the uncomfortable positions in order to obtain diagnostic images. This is high value in patients who are already in pain.
- MDCT Multi-planar (MPR) and the 3-D capabilities improving, detection, characterization, staging of pelvic fractures and the pre-surgical planning.
- MDCT provides detailed information about the soft tissues in and around the pelvis.
- MDCT very important component in the pre-operative assessment of pelvic fractures.
- MDCT is the most suitable technique for assessment of fractures of pelvic and their classification.
- MDCT is more sensitive than plain radiography in detection and diagnosis of the pelvic fractures and can change the stability diagnosis of the fractures.

REFERENCES

REFERENCES

- 1- Ooi CK, Goh HK, Tay SY, Phua DH. Patients with pelvic fracture: what factors are associated with mortality. *Int J Emerg Med* 2010; 3: 299-304.
- 2- Shlamovitz GZ, Mower WR, Bergman J, Crisp J, DeVore HK, Hardy D, et al. Poor test characteristics for the digital rectal examination in trauma patients. *Ann Emerg Med* 2007; 50: 25-33, 33.
- 3- Giannoudis PV, Pape HC. Damage control orthopaedics in unstable pelvic ring injuries. *Injury* 2004; 35: 671-7.
- 4- Grotz MR, Allami MK, Harwood P. Open pelvic fractures: epidemiology, current concepts of management and outcome. *Injury* 2005; 36: 1-13.
- 5- Mohanty K, Musso D, Powell JN. Emergent management of pelvic ring injuries: an update. *Can J Surg* 2005; 48: 49-56.
- 6- Starr AJ, Griffin DR, Reinert CM. Pelvic ring disruptions: prediction of associated injuries, transfusion requirements, pelvic arteriography, complications, and mortality. *J Orthop Trauma* 2002; 16: 553-61.
- 7- Blackmore CC, Jurkovich GJ, Linnau KF. Assessment of volume of hemorrhage and outcome from pelvic fracture. *Arch Surg* 2003; 138: 504-9.
- 8- Senohradski K, Kavoric B, Miric D. Computer tomography in the diagnosis and therapy of acetabular fractures. *Srp Arh Celok Lek* 2001; 129: 194-8.
- 9- Guillamondegui OD, Pryor JP, Gracias VH, Gupta R, Reilly PM, Schwab CW. Pelvic radiography in blunt trauma resuscitation: a diminishing role. *Trauma* 2002; 53: 1043-7.
- 10- Chen TW, Yang ZG, Dong ZH, Chu ZG, Yao J, Wang QL. Pelvic crush fractures in survivors of the Sichuan earthquake evaluated by digital radiography and multidetector computed tomography. *Skeletal Radiol* 2010; 39: 1117-22.
- 11- Ohashi K, El-Khoury GY, Abu-Zahra K, Berbaum K. Interobserver agreement for Letournel acetabular fracture classification using MD-CT: are the standard Judet radiographic views necessary. *Radiology* 2006; 241: 386-91.
- 12- Falchi M, Rollandi GA. CT of pelvic fractures. *Eur J Radiol* 2004; 50: 96-105.
- 13- Butler P, Mithchel AM, Ellis H. Applied radiological anatomy for medical students. Vol. 13. Cambridge, UK, New York: Cambridge University Press, 2007. 129-45.
- 14- Agur A, Dally A. Grant's atlas of anatomy. 12th ed. Vol. 3. Philadelphia, PA: Lippincott Williams & Wilkins, 2009. 194-285.
- 15- Martinez CR, Di Pasquale TG, Helfet DL, Graham AW, Sanders RW, Ray LD. Evaluation of acetabular fractures with two -and three-dimensional CT. *Radiographics* 1992; 12: 227-41.
- 16- Hunter JC, Brandser EA, Tran KA. Pelvic and acetabular trauma. *Radiol Clin North Am* 1997; 35: 559-90.

- 17- Harley JD, Mack LA, Winquist RA. CT of acetabular fractures: Comparison with conventional radiography. *AJR Am J Roentgenol* 1982; 138: 413-7.
- 18- Goulet J, Bray T. Complex acetabular fractures. *Clin Orthop Relat Res* 1989; 240: 9-20.
- 19- Standrings S. *Gray's anatomy: the anatomical basis of clinical practice*. 39th ed. Edinburgh, New York: Elsevier Churchill Livingstone, 2005. 66-116.
- 20- Heare MM, Heare TC, Gillespy T. Diagnostic imaging of pelvic and chest wall trauma. *Radiol Clin North Am* 1989; 27: 873-89.
- 21- Butler P, Mitchel A, Ellis H. *Applied radiological anatomy for medical students*. 1st ed. Cambridge, UK, New York: Cambridge University Press, 2007. 47-64.
- 22- Snell R. *Clinical anatomy by regions*. 9th ed. Philadelphia: Lippincott Williams & Wilkins, 2012. 240-301.
- 23- Wegener OH. *Whole body computerized tomography*. 2nd ed. Edinburgh: Churchill Livingstone, 2000. 553-67.
- 24- Tile M. *Fractures of the pelvis and acetabulum*. 2nd ed. Philadelphia, Williams & Wilkins, 1995. 240-320.
- 25- Hansen J, Henary F. *Netter's clinical anatomy*. 2nd ed. Philadelphia: Saunders/Elsevier, 2010. 181-226.
- 26- Reinert CM, Starr AJ, Griffin DR. Pelvic ring disruptions: prediction of associated injuries, transfusion requirements, pelvic arteriography, complications, and mortality. *J Orthop Trauma* 2002; 16: 553-61.
- 27- Wegener OH. *Whole body computerized tomography*. 2nd ed. Vol. 27. Edinburgh: Churchill Livingstone, 2000. 553-67.
- 28- Kricum ME. Fractures of the pelvis. *Orthop Clin North Am* 1990; 21: 573-90.
- 29- Laasonen EM. Missed sacral fractures. *Ann Clin Res* 1997; 9: 84-7.
- 30- Smith TK, Jhonstong RM, Bucholz RW, Browner BD. Symposium: external fixation of pelvic fractures. *Contemp Orthop* 1999; 7: 81-101.
- 31- Ashman CJ, Yu J, Chung CB. Chance fractures; antero posterior radiographic signs. *Emerg Radiol* 1997; 14: 320-5.
- 32- Furey A, Otool R, Nascone J. Classification of pelvic fractures: analysis of inter- and intraobserver variability using the Young-Burgess and Tile classification systems. *Orthopedics* 2009; 32: 401.
- 33- Stambaugh L, Blackmore CC. Pelvic ring disruptions in emergency radiology. *Eur J Radiol* 2003; 48: 71-87.
- 34- Marsh JL, Slongo TF, Agel J, Broderick JS, Creevey W, DeCoster TA, et al. Fracture and dislocation classification compendium-2007: Pelvis. *J Orthop Trauma* 2007; 21(Suppl 10): S59-S67.
- 35- Young JWR. Skeletal trauma: general consideration. In: Sutton D (ed). *Text Book of radiology and imaging*. 7th ed. Vol. 1. Churchill and Livingstone, 2004. 1400-4.

- 36- Rogers LF. Radiology of skeletal trauma. 3rd ed. New York: Churchill Livingstone, 2000. 991-1105.
- 37- Harris JH. The Radiology of emergency medicine. 4th ed. Philadelphia, Baltimore: Lippincott Williams & Wilkins Company, 2000. 725-814.
- 38- Young JW, Burgess AR, Brumback RJ, Poka A. Lateral compression fractures of the pelvis: the importance of plain radiographs in the diagnosis and surgical management. *Skeletal Radiol* 1986; 15: 103-4.
- 39- Kane WJ. Fractures of the pelvis. In: Rockwood CCA, Green DP (eds). *Fractures in adults*. 2nd ed. Philadelphia: J. B. Lippincott, 2000. 1093-209.
- 40- Watura R, Cobby M, Taylor J. Multislice CT in imaging of trauma of the spine, pelvis and complex foot injuries. *Br J Radiol* 2004; 77: S46-S63.
- 41- Koo H, Leveridge M, Thompson C, Zdero R, Bhandari M, Hans JK, et al. Interobserver reliability of the young-burgess and tile classification systems for fractures of the pelvic ring. *J Orthop Trauma* 2010; 22: 379-84.
- 42- Greenspan A, Michael W. *Orthopedic radiology*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins, 2004. 179-207.
- 43- Rensik CS, Stackhouse DJ, Shanmuganathan K, Young J. Diagnosis of pelvic fractures in patients with acute pelvic trauma: efficacy of plain radiographs. *AJR Am J Roentgenol* 1992; 158: 109-12.
- 44- Montana MA, Richardson ML, Kilcoyne RF, Harley JD, Shuman WP, Mack LA. CT of sacral injury. *Radiology* 1986; 161: 499-503.
- 45- Dunn EL, Berry PH, Connally JD. Computed tomography of the pelvis in patients with multiple injuries. *J Trauma* 1983; 23: 378-83.
- 46- Jackson H, Kam J, Harris JH, Harle TS. The sacral arcuate lines in upper sacral fractures. *Radiology* 1998; 145: 35-9.
- 47- Gill K, Bucholz RW. The role of computerized tomographic scanning in the evaluation of major pelvic fractures. *J Bone Joint Surg Am* 1984; 66: 34-9.
- 48- Buckely SL, Burkus JK. Computerized axial tomography of pelvic ring fractures. *J Trauma* 1987; 27: 496-502.
- 49- Fishman EK, Drebin B, Magid D, Scott WW Jr, Ney DR, Brooker AF Jr, et al. Volumetric rendering techniques: application for three dimensional imaging of the hip. *Radiology* 1987; 163: 737-8.
- 50- Raffi M, Firooznia H, Golimbu C, Waugh T Jr, Naidich D. The impact of CT in clinical management of pelvic and acetabular fractures. *Clin Orthop* 1983; 178: 228-35.
- 51- Marincek B. *Emergency radiology imaging and intervention; with 49 tables*. Berlin, Heidelberg, New York: Springer, 2007. 225-49.
- 52- Geusens E, Breuseghem V, Pans S, Brys P. Imaging in trauma of the pelvis and hip region. *JBR-BTR* 2004; 87: 190-202.

- 53- Falchi M, Rollandi GA. CT of pelvic fractures. *Eur J Radiol* 2004; 50: 96-105.
- 54- Joseph N. Trauma imaging of the pelvis. *CE Essentials: Online Radiography Continuing Education for Radiologic X ray Technologist*. 2006. Available at: <https://www.ceessentials.net/article34.html>.
- 55- Butler P, Mitchel A, Ellis H. *Applied radiological anatomy for medical students*. Cambridge, UK, New York: Cambridge University Press, 2007. 47-64.
- 56- Graf KW Jr, Bal BS, Talavera F, McCarthy JJ. Unstable pelvic fractures. *Medscape*. 2014. Available at: <http://emedicine.medscape.com/article/1247426-overview>.
- 57- Dalal SA, Burgess AR, Siegel JH, Young JW, Brumback RJ, Poka A, et al. Pelvic fracture in multiple trauma: classification by mechanism is key to pattern of organ injury, resuscitative requirements, and outcome. *J Trauma* 1989; 29: 981-1000.
- 58- Adam A, Dixon AK, Grainger RG, Allison DJ. *Grainger & Allison's diagnostic radiology: A textbook of medical imaging*. 5th ed. Philadelphia, PA: Churchill Livingstone, 2007. 46.
- 59- Hilty MP, Behrendt I, Benneker LM, Martinolli L, Stoupis C, Buggy DJ, et al. Pelvic radiology in ATLS algorithms: A diminishing role. *World J Emerg Surg* 2008; 3: 11.
- 60- Horton KM, Sheth S, Corl F, Fishman EK. Multidetector row CT: principles and clinical applications. *Crit Rev Comput Tomogr* 2002; 43: 143-81.
- 61- Yitta S, Hetch E, Slywotzky C, Bennet G. Added value of multiplanar reformation in the multidetector CT evaluation of the female pelvis: a pictorial review. *Radiographics* 2009; 29: 1987-2003.
- 62- Watura R, Cobby M, Taylor J. Multislice CT in imaging of trauma of the spine, pelvis and complex foot injuries. *Br J Radiol* 2004; 1: S46-63.
- 63- Falchi M, Rollandi GA. CT of pelvic fractures. *Eur J Radiol* 2004; 50: 96-105.
- 64- Flohr T, Stierstorfer K, Bruder H, Simon J, Polacin A, Schaller S. Image reconstruction and image quality evaluation for a 16-slice CT scanner. *Med Phys* 2003; 30: 832-45.
- 65- Linsenmaier U, Krötz M, Häuser H, Rock C, Rieger J, Bohndorf K, et al. Whole-body computed tomography in polytrauma: techniques and management. *Eur Radiol* 2002; 12: 1728-40.
- 66- Wintermark M, Mouhsine E, Theumann N, Mordasini P, van Melle G, Leyvraz PF, et al. Thoracolumbar spine fractures in patients who have sustained severe trauma: depiction with multi-detector row CT. *Radiology* 2003; 227: 681-9.
- 67- Herzog C, Ahle H, Mack MG, Maier B, Schwarz W, Zangos S, et al. Traumatic injuries of the pelvis and thoracic and lumbar spine: does thin-slice multidetector-row CT increase diagnostic accuracy? *Eur Radiol* 2004; 14: 1751-60.
- 68- Geijer M, El-Khoury G. MDCT in the evaluation of skeletal trauma: principles, protocols, and clinical applications. *Emerg Radiol* 2006; 13: 7-18.

- 69- Smith W, Williams A, Agudelo J, Shannon M, Morgan S, Stahel P, et al. Early predictors of mortality in hemodynamically unstable pelvis fractures. *J Orthop Trauma* 2007; 21: 31-7.
- 70- Gurevitz S, Bender B, Tytiun Y, Velkes S, Salai M, Stein M. The role of pelvic fractures in the course of treatment and outcome of trauma patients. *Isr Med Assoc J* 2005; 7: 623-6.
- 71- Dente CJ, Feliciano DV, Rozycki GS, Wyrzykowski AD, Nicholas JM, Salomone JP, et al. The outcome of open pelvic fractures in the modern era. *Am J Surg* 2005; 190: 830-5.
- 72- Gylling SF, Ward R, Holcroft J, Bray T, Chapman M. Immediate external fixation of unstable pelvic fractures. *Am J Surg* 1985; 150: 721-4.
- 73- Maier B, Schwarz W, Herzog C, Ahle H, Mack MG, Zangos S, et al. Traumatic injuries of the pelvis and thoracic and lumbar spine: does thin-slice multidetector-row CT increase diagnostic accuracy? *Eur Radiol* 2004; 14: 1751-60
- 74- Gänsölen A, Pohlemann T, Paul C, Lobenhoffer P, Tschernke H. Epidemiology of pelvic ring injuries. *Injury* 1996; 27(Suppl 1): S-A13-20.
- 75- Dong ZH, Yao J, Wang QL, Chen TW, Yang ZG, Chu ZG. Pelvic crush fractures in survivors of the Sichuan earthquake evaluated by digital radiography and multidetector computed tomography. *Skeletal Radiol* 2010; 39: 1117-22.
- 76- Li FP, Li M, Hua Q, Zhao LJ, Luo JN. Study on the spiral CT reconstruction in the diagnosis and treatment of pelvic ring fractures. *Zhongguo Gu Shang* 2010; 23: 204-7.
- 77- Wedegärtner U, Gatzka C, Rueger JM, Adam G. Multislice CT (MSCT) in the detection and classification of pelvic and acetabular fractures. *Rofo* 2003; 175: 105-11.
- 78- Li M, Xu RM, Zheng Q, Xiao BP, Wang GP. Imaging diagnoses and clinical meaning for easy neglect occult posterior pelvic ring injury. *Zhongguo Gu Shang* 2008; 21: 810-3.
- 79- Chmelová J, Mrázková D, Dzupa V, Báca V, Grill R, Pleva L. The role of plain radiography in pelvic trauma in the era of advanced computed tomography. *Acta Chir Orthop Traumatol Cech* 2006; 73: 394-9.
- 80- Koo H, Leveridge M, Thompson C, Zdero R, Bhandari M, Kreder HJ, et al. Interobserver reliability of the young-burgess and tile classification systems for fractures of the pelvic ring. *J Orthop Trauma* 2008; 22: 379-84.
- 81- Manson T, O'Toole RV, Whitney A, Duggan B, Sciadini M, Nascone J. Young-Burgess classification of pelvic ring fractures: does it predict mortality, transfusion requirements, and non-orthopaedic injuries. *J Orthop Trauma* 2010; 24: 603-9.
- 82- Furey AJ, O'Toole RV, Nascone JW, Sciadini MF, Copeland CE, Turen C. Classification of pelvic fractures: analysis of inter- and intraobserver variability using the Young-Burgess and Tile classification systems. *Orthopedics* 2009; 32: 401.

- 83- Dente CJ, Feliciano DV, Rozycki GS, Wyrzykowski AD, Nicholas JM, Salomone JP, et al. The outcome of open pelvic fractures in the modern era. *Am J Surg* 2005; 190: 830-5.
- 84- Chu ZG, Yao J, Wang QL, Chen TW, Yang ZG, Dong ZH. Pelvic crush fractures in survivors of the Sichuan earthquake evaluated by digital radiography and multidetector computed tomography. *Skeletal Radiol* 2010; 39: 1117-22.
- 85- Gabbe BJ, de Steiger R, Esser M, Bucknill A, Russ MK, Cameron PA. Predictors of mortality following severe pelvic ring fracture: results of a population-based study. *Injury* 2011; 42: 985-91.
- 86- Lefaiivre K, Padalecki J, Starr A. What constitutes a Young and Burgess lateral compression-I (OTA 61-B2) pelvic ring disruption? A description of computed tomography-based fracture anatomy and associated injuries. *J Orthop Trauma* 2009; 23: 16-21.

PROTOCOL

**ROLE OF MULTI-DETECTOR COMPUTED TOMOGRAPHY
IN EVALUATION OF PELVIC FRACTURES**

دور الأشعة المقطعية متعددة المقاطع في تقييم كسور الحوض

Protocol of a thesis submitted
to the Faculty of Medicine
University of Alexandria
In partial fulfillment of the
requirements of the degree of
**Master of Radiodiagnosis and
Intervention**

خطة بحث مقدمة
لكلية الطب
جامعة الإسكندرية
إيفاء جزئياً
لشروط الحصول على درجة
الماجستير في الأشعة التشخيصية
والتدخلية

by

من

Feras Ahmed Khalaf
MBBCh, Yemen.
Visiting Resident
Alexandria University Hospitals
Department of Radiodiagnosis
Faculty of Medicine
University of Alexandria
2012

فراس احمد خلف
بكالوريوس الطب والجراحة ، اليمن
طبيب مقيم زائر
مستشفيات جامعة الإسكندرية
قسم الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية
٢٠١٢

2.

SUPERVISORS**المشرفون****Prof. Dr. Hesham Taha Kotb**

Professor of Radiodiagnosis,
Faculty of Medicine,
University of Alexandria.

أ.د/ هشام طه قطب

أستاذ الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية

Prof. Dr. Yosry Emad Eldin Eid

Professor of Orthopedic Surgery
and Traumatology,
Faculty of Medicine,
University of Alexandria.

أ.د/ يسري عماد الدين عيد

استاذ جراحة العظام والكسور
كلية الطب
جامعة الإسكندرية

CO-SUPERVISOR**المشرف المشارك****Dr. Sherif Abdelmonem Shama**

Lecturer in Radiodiagnosis,
Faculty of Medicine,
University of Alexandria.
For his experience in MDCT Imaging

د/ شريف عبد المنعم شامة

مدرس الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية
وذلك لخبرته في فحوصات
الأشعة المقطعية متعددة المقاطع



3.

ASSISTANT-RESEARCHER

الباحث المساعد

Ammar Ali Alani

عمار علي العاني

3rd grade student

طالب بالفرقة الثالثة

Faculty of Medicine,



كلية الطب

University of Alexandria.

جامعة الإسكندرية

Mobile phone: 01113399928

E-mail: dr.ammar07@hotmail.com



مركز البحوث والبحوث

4.

INTRODUCTION

The pelvis is a complex bony ring comprised of multiple components. The sacrum and ilia make up the posterior portion of the ring, with the ischia and pubic bones anteriorly providing a connection between the lower limbs and trunk. However, the bony articulations, sacroiliac and pubic symphysis, are not inherently stable.⁽¹⁾

Pelvic fracture is a serious injury that is associated with significant morbidity and mortality. This is a result of the high energy transfer to the pelvic region. Road traffic accidents and falls from height are by far the most common mechanisms of injury.⁽²⁾

Fractures of the pelvic ring have been reported to comprise 2% to 8% of all skeletal injuries.⁽³⁻⁵⁾ Patients are most often young males in their early 30s, and motor vehicle collisions are the most common inciting cause.^(6,7) Because of the high energy involved, patients with pelvic ring disruptions usually have multiple injuries.⁽⁶⁾ Morbidity and mortality is also high, with overall mortality rates of 17% for patients with pelvic fractures.⁽⁷⁾

Pelvic fractures were classified according to pelvic instability; if a pelvic fracture is stable, it is Type A; a fracture that is unstable to rotation but vertically stable is termed Type B; severe pelvic ring disruption that is unstable to both rotation and vertical displacement is Type C.⁽⁸⁾


Detection and assessment of pelvic fractures is a critical component of the initial evaluation of any major trauma patient.⁽⁹⁾ pelvic fractures are frequently complex and the precise pathological anatomy is not easily demonstrated by routine radiographs.⁽¹⁰⁾ Plain radiographs alone have limited sensitivity for detection of pelvic fractures compared with CT.⁽¹¹⁾ Fractures of the sacrum and acetabulum are most easily overlooked.^(10, 11)

Digital radiography and Computed Tomography are valuable for evaluating pelvic fractures.⁽¹²⁾ In most polytrauma patients, the first imaging study today is MDCT. Although axial CT images have a higher diagnostic accuracy in detecting and characterizing pelvic injuries than radiography, evaluation with MPR images has been shown to increase diagnostic accuracy.⁽¹³⁾ Pelvic trauma is where 3-D imaging first made a significant clinical impact, and it is now a standard requisite for surgical planning, being a valuable complement to axial and MPR images.⁽¹⁴⁾


CT is useful for detecting intra-articular fragments, fragment interposition, marginal impaction, occult pelvic ring fractures and evaluating the sacroiliac complex to determine if vertical instability is present.^(15, 16)

MSCT detected additional fractures in 17% of patients not seen on conventional radiographs. In some patients, MSCT changed the fracture classification by delineating the full extent of the fracture. MPR visualize sacral fractures better than axial source images. MSCT of the pelvis is required for correct fracture detection.⁽¹⁷⁾ MSCT has increased the speed, efficiency and accuracy of computed tomographic imaging in musculoskeletal trauma. Added advantages such as MPRs and 3D imaging have greatly enhanced the ability to evaluate complex anatomical areas such as the spine, pelvis and foot.^(18, 19)

ترقی عوامی



MPR technique proved to be superior in the detection of fractures of the pelvis, leading to significantly better results than conventional radiographs and markedly better results than transverse slices alone.⁽²⁰⁾ In severe trauma patients MDCT proved to be far superior in the detection and classification of traumatic fractures of the spine and the pelvis. Particularly thin-slice technique combined with multiplanar reformation leads to significantly better results than conventional radiographs or transverse images alone.⁽²¹⁾



Ju

7.

AIM OF THE WORK

The aim of this work will be directed to study the role of Multi-Detector Computed Tomography (MDCT) over Conventional Radiography in evaluation and clearing the diagnosis of osseous injuries of the pelvis.

Hes

STX

الهدف من هذا العمل

[Handwritten mark]

PATIENTS

A prospective study will be conducted on 30 patients presenting with pelvic trauma referred to the radiology department at Alexandria university Hospitals.

[Handwritten signature]


[Handwritten signature]

[Handwritten signature]

METHODS

Selected patients will be subjected to:

- Full history taking.
- Thorough clinical examination.
- Plain X-ray of the pelvis:
 - AP view.
 - Oblique views (external , internal).
 - Inlet / Outlet views.
- Non enhanced MDCT of the pelvis:
 - Multi-planar volume reformation (MPR).
 - Three-dimensional VR.
- The medical ethics will be considered. The patient has to get benefit from the examination [An informed consent will taken from patient in case of incompetent patient the informed consent will be from the guardians].



حکومت سندھ

X

Ethics Of Research

Research on human or human products:

- Prospective study:** informed consent will taken from patient. In case of incompetent patient the informed consent will be from the guardians.

- Retrospective study:** Confidentiality of record will be considered.

- DNA/ genomic material:** informed consent for DNA/ genomic test and for research will be taken from patients. No further tests will be carried out except with further approval of committee and patients. If the samples will travel outside Egypt the researcher will be responsible for transportation and security approval.

- All drugs used in the research is approved by the Egyptian ministry of health.

Research on animal:

- The animal species is appropriate to the test.

- After test, if the animal will suffer it will be euthanized and properly disposed.

- After operation, it will have a proper postoperative care.



HS

Handwritten signature in Arabic

Handwritten signature

RESULTS


The results of this study will be calculated, tabulated and statistically analyzed according to the appropriate methods.



سید ذکریا کمالی

DISCUSSION

The results will be discussed in view of achievement of the aim, their significance and their comparison with other available previous related researches.





REFERENCES

1. Attias N, Arzani S, Duncan G, Taber KH, Hayman LA. Sectional imaging anatomy: pelvic ring ligaments. *J Comput Assist Tomogr* 2001; 25:975-9.
2. Ooi CK, Goh HK, Tay SY, Phua DH. Patients with pelvic fracture: what factors are associated with mortality. *Int J Emerg Med* 2010; 3:299-304.
3. Giannoudis PV, Pape HC. Damage control orthopaedics in unstable pelvic ring injuries. *Injury* 2004; 35(7):671-7.
4. Grotz MR, Allami MK, Harwood P. Open pelvic fractures: epidemiology, current concepts of management and outcome. *Injury* 2005; 36(1):1-13.
5. Mohanty K, Musso D, Powell JN. Emergent management of pelvic ring injuries: an update. *Can J Surg* 2005; 48(1):49-56.
6. Starr AJ, Griffin DR, Reinert CM. Pelvic ring disruptions: prediction of associated injuries, transfusion requirements, pelvic arteriography, complications, and mortality. *J Orthop Trauma* 2002; 16: 553-61.
7. Blackmore CC, Jurkovich GJ, Linnau KF. Assessment of volume of hemorrhage and outcome from pelvic fracture. *Arch Surg* 2003; 138:504-9.
8. Tile M. *Fractures of the Pelvis and Acetabulum*. 3rd ed. Baltimore, MD: Williams & Wilkins; 2003.




9. Stambaugh LE, Blackmore CC. Pelvic ring disruptions in emergency radiology. *European Journal of Radiology* 2003; 48 :71-87.
10. Senohradski K, Kavoric B, Miric D. Computer tomography in the diagnosis and therapy of acetabular fractures. *Srp Arh Celok Lek* 2001; 129:194–8.
11. Guillaumondegui OD, Pryor JP, Gracias VH, Gupta R, Reilly PM, Schwab CW. Pelvic radiography in blunt trauma resuscitation: a diminishing role. *Trauma* 2002; 53: 1043–7.
12. Tian-wu C, Zhi-gang Y, Zhi-hui D, Zhi-gang C, Jin Y, Qi-ling W. Pelvic crush fractures in survivors of the Sichuan earthquake evaluated by digital radiography and multidetector computed tomography. *Skeletal Radiol* 2010; 39:1117–22.
13. Ohashi K, El-Khoury GY, Abu-Zahra K, Berbaum K. Interobserver agreement for Letournel acetabular fracture classification using MD-CT: are the standard Judet radiographic views necessary? *Radiology* 2006; 241:386-91.
14. Falchi M, Rollandi GA. CT of pelvic fractures. *Eur J Radiol* 2004; 50:96–105.
15. Otto W. Acetabulum fractures. Diagnosis, classification, evaluation. *Zentralbl Chir* 2000; 125:725–9.
16. McCormack R, Strauss EJ, Alwattar BJ, Tejwani NC. Diagnosis and Management of Pelvic Fractures *Bulletin of the NYU Hospital for Joint Diseases* 2010;68(4):281-91.

17. Wedegartner U, Gatzka C, Rueger JM, Adam G. Multislice CT (MSCT) in the detection and classification of pelvic and acetabular fractures. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 2003; 175:105–11.
18. Watura R, Cobby M, Taylor J. Multislice CT in imaging of trauma of the spine, pelvis and complex foot injuries. *The British Journal of Radiology* 2004; 77:S46–S63.
19. Koo H, Leveridge M, Thompson C, Zdero R, Bhandari M, Hans JK, et al. Interobserver Reliability of the Young-Burgess and Tile Classification Systems for Fractures of the Pelvic Ring. *J Orthop Trauma* 2010; 22:379-84.
20. Wedegartner U, Lorenzen M, Lorenzen J, Nolte-Ernsting C, Weber C, Dieckmann C, et al. Multislice CT of the pelvis: dose reduction with regard to image quality. *Fortschr Rontgenstr* 2004; 176:106–12.
21. Herzog C, Ahle H, Mack MG, Maier B, Schwarz W, Zangos S, et al. Traumatic injuries of the pelvis and thoracic and lumbar spine: does thin-slice multidetector-row CT increase diagnostic accuracy? *Eur Radiol* 2004; 14:1751–60.





ARABIC SUMMARY

الملخص العربي

شملت هذه الدراسة ٣٠ مريضا باشتباه سريري بكسر في الحوض والمحولين من قسم الأشعة التشخيصية في مستشفى جامعة الإسكندرية، حيث انه تم فحص جميع المرضى بواسطة الأشعة المقطعية . فكان العدد ٢١ من الذكور و ٩ من الإناث ٣٠٪ مع نسبة الذكور إلى الإناث من ٢,٣:١. وتراوح أعمارهم بين ١٣ و ٨٥ عاما مع متوسط عمر ٣٦ عاما.

كان الهدف من هذا العمل هو دراسة دور التصوير المقطعي متعدد المقاطع على الأشعة التقليدية في التقييم وتأكيد تشخيص الإصابات في عظم الحوض.

في هذه الدراسة ، تم أخذ التاريخ الكامل والفحص السريري الدقيق للمرضى، تصوير الحوض بالأشعة السينية كلما أمكن ذلك (امامي خلفي ، "مدخل / مخرج"، مائل ،"داخلي / خارجي") وتصوير بالأشعة المقطعية متعدد المقاطع بدون صبغة للحوض (الفحص الثلاثي الأبعاد والصور ثلاثية الأبعاد).

وقد أجريت الفحوصات على ضوء سرعة جي اي رباعي المواسح وسيمنس سداسي المواسح. كانت عوامل الفحص ١٤٠ كيلو فلت و ١٤٥ مللي أمبير لكل مقطع ودوران الأنبوبة ٠,٥ ثانية وسمك المقطع ١ مم .

كسر الحوض هو إصابة خطيرة ذو معدلات الاعتلال والوفيات كبيرة. هذا هو نتيجة لنقل الطاقة العالية إلى منطقة الحوض. وقد تم الإبلاغ عن كسور في الحوض لتشمل ٢٪ إلى ٨٪ من جميع إصابات الهيكل العظمي. غالبا ما تكون الحالات من الشباب الذكور في العقد الثالث ، السبب الأكثر شيوعا هو حوادث السير . بسبب الطاقة الهائلة، المرضى الذين يعانون من اضطرابات في حلقة الحوض عادة ما يكونو ذو إصابات متعددة. معدل المرض والوفيات مرتفع أيضا، مع معدلات الوفيات الإجمالية بنسبة ١٧٪ للمرضى الذين يعانون من كسور الحوض.

التصوير الشعاعي الرقمية والتصوير المقطعي ذات قيمة لتقييم كسور الحوض. في معظم مرضى متعددي الإصابات تعتبر الأشعة المقطعية متعددة المقاطع هي الخيار الاول في التصوير. بالإضافة الي أن الصور المقطعية المحورية لديها أعلى دقة التشخيص في الكشف وتوصيف إصابات الحوض من التصوير الشعاعي، قد تبين ان زيادة التقييم ودقة التشخيص تأتي مع الصور في الأبعاد الثلاثة وثلاثية الأبعاد. في إصابات الحوض التصوير ثلاثي الأبعاد يعتبر ذو تأثير سريري كبير، وتعتبر الان المعيار الاول لتخطيط العمليات الجراحية.

الملخص العربي

لجنة الإشراف

.....

أ.د. / هشام طه قطب
أستاذ الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية

.....

أ.د. / يسري عماد الدين عيد
أستاذ جراحة العظام والكسور
كلية الطب
جامعة الإسكندرية

.....

د. / شريف عبد المنعم شامة
أستاذ مساعد الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية

دور الأشعة المقطعية متعددة المقاطع في تقييم كسور الحوض

مقدمة من

فراس احمد خلف

بكالوريوس الطب والجراحة - جامعة العلوم التطبيقية - اليمن

للحصول على درجة

الماجستير

في

الأشعة التشخيصية

موافقون

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د. / هشام طه قطب
أستاذ الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية

أ.د. / عادل علي رمضان
أستاذ الأشعة التشخيصية
كلية الطب
جامعة الإسكندرية

أ.د. / هشام علي بدوي
أستاذ الأشعة التشخيصية
معهد البحوث الطبية
جامعة الإسكندرية



دور الأشعة المقطعية متعددة المقاطع في تقييم كسور الحوض

رسالة علمية

مقدمة إلى كلية الطب - جامعة الإسكندرية
إستيفاء للدراسات المقررة للحصول على درجة

الماجستير

في

الأشعة التشخيصية

مقدمة من

فراس احمد خلف

بكالوريوس الطب والجراحة - جامعة العلوم التطبيقية - اليمن