

REFERENCES

1. Sternberg K, Greenfield SP, Williot P, Wan J. Pediatric stone disease: an evolving experience. *J Urol* 2005;174:1711-16.
2. Rizvi SA, Naqvi SA, Hussain Z, Hashmi A, Hussain M, Zafar MN, et al. Pediatric urolithiasis: Developing nation perspectives. *J Urol* 2008;168:1522-5.
3. Hesse A, Brøndle E, Wilbert D, Køhrmann KU, Alken P. Study on the prevalence and incidence of urolithiasis in Germany comparing the years 1979 vs. 2006. *Eur Urol* 2003; 44:709-13.
4. Tanaka ST, Pope JC., 4th Pediatric stone disease. *Curr Urol Rep.* 2009;10:138, 43.
5. Pearle MS, Calhoun E, Curhan GC. Urolithiasis. In: Litwin MS, Saigal CS, editors. *Urologic Diseases in America*. US Government Printing Office, NIH Publication No. 07-5512; 2007. pp. 283–319.
6. Taylor EN, Stampfer MJ, Curhan GC. Dietary factors and the risk of incident kidney stones in men: new insights after 14 years of follow-up. *J Am Soc Nephrol* 2004; 15: 3225-31.
7. Goldfarb DS. Increasing prevalence of kidney stones in the United States. *Kidney Int* 2003; 63: 1951-5.
8. Ayoob R, Wang W, Schwaderer A. Body fat composition and occurrence of kidney stones in hypercalciuric children. *Pediatr. Nephrol.* 2011; vol. 26 (12) pp. 2173-8.
9. Kieran K, Giel DW, Morris BJ. Pediatric urolithiasis, does body mass index influence stone presentation and treatment? *J Urol* 2010; 184:1810-5.
10. Saxena A, Sharma RK. Nutritional aspect of nephrolithiasis. *Indian J Urol.* 2010;26:523-30.
11. Wu, HY, Docimo, SG Surgical management of children with urolithiasis. *Urol Clin North Am* 2004;31:589-94.
12. Kruse K, Kracht U, Kruse U: Reference values for urinary calcium excretion and screening for hypercalciuria in children and adolescents. *Eur J Pediatr* 1984, 143:23–31.
13. Sargent JD, Stukel TA, Kresel J, Klein RZ: Normal values for random urinary calcium to creatinine ratios in infancy. *J Pediatr* 1993, 123:393–7.
14. Stapleton FB, Noe HN, Jerkins GR, Roy S III: Urinary excretion of calcium following an oral calcium-loading test in healthy children. *Pediatrics* 1982, 69:594–7.
15. Bartosh SM: Medical management of pediatric stone disease. *Urol Clin North Am* 2004, 31:575–87.

Reference

16. Polinsky MS, Kaiser KA, Baluarte HJ, et al.: Renal stones and hypercalciuria. *Adv Pediatr* 1993, 40:353–84.
17. Watts RW, Morgan SH, Danpure CJ, et al.: Combined hepatic and renal transplantation in primary hyperoxaluria type I: Clinical report of nine cases. *Am J Med* 1991, 90:179–88.
18. Tekin A, Tekgül S, Atsu N, et al.: A study of the etiology of idiopathic calcium urolithiasis in children: hypocitruria is the most important risk factor. *J Urol* 2000, 164:162–5.
19. Tekin A, Tekgül S, Atsu N, et al.: Oral potassium citrate treatment for idiopathic hypocitruria in children with calcium urolithiasis. *J Urol* 2002, 168:2572–4.
20. Milliner DS, Murphy ME: Urolithiasis in pediatric patients. *Mayo Clin Proc* 1993, 68:241–8.
21. Hagar BH, Magath TB: The etiology of incrustrated cystitis with alkaline urine. *JAMA* 1925, 85:1352–5.
22. Nemoy NJ, Staney TA: Surgical, bacteriological, and biochemical management of “infection stones.” *JAMA* 1971, 215:1470–6.
23. Straub M, Strohmaier WL, Berg W, et al.: Diagnosis and metaphylaxis of stone disease. Consensus concept of the National Working Committee on Stone Disease for the Upcoming German Urolithiasis Guideline. *World J Urol* 2005, 23:309–23.
24. Tekgül S, Riedmiller H, Beurton D, et al.: Guidelines on Pediatric Urology. *EAU Update March* 2006, 42–8.
25. Takazawa R, Kitayama S, Tsujii T. Successful outcome of flexible ureteroscopy with holmium laser lithtripsy for renal stones 2 cm or greater. *Int J Urol* 2012;19:264-7.
26. Honeck P, Wendt-Nordahi G, Krombach P, Bach T, Hacker A, Alken P, et al. Does open stone surgery still play a role in the treatment of urolithiasis. Data of a primary urolithiasis center? *J Endourol.* 2009;23:1209–12.
27. Alivizatos G, Skolarikos A. Is there still a role for open surgery in the management of renal stones? *Curr Opin Urol.* 2006;16:106–11.
28. Paik ML, Resnick MI. Is there a role for open stone surgery? *Urol Clin North Am.* 2000;27:323–31.
29. Rizvi SAH, Naqvi SAA, Hussain Z, Hashmi A, Hussain M, Zafar MN, et al. Management of pediatric urolithiasis in Pakistan: Experience with 1,440 children. *J Urol.* 2003;169:634–7.
30. Buchholz NNP, Hitchings A, Albanis S. The (soon forgotten) art of open surgery: to train or not to train? *Ann R Coll Surg Engl.* 2006;88:214–7.
31. Zargooshi J. Open stone surgery in children. Is it justified in the era of minimally invasive therapies? *BJU Int.* 2001;88:928–31.

Reference

32. Rizvi SAH, Sultan S, Zafar MN, Ahmed B, Faiq SM, Hossain KZ, et al. Evaluation of children with urolithiasis. In *J Urol*. 2007;23:420–7.
33. Farhat WA, Kropp BP: Surgical Treatment of Pediatric Urinary Stones: Lesson 3 AUA Update 2007: 26.
34. Muslumanoglu AY, Tefekli A, Sarilar O, Binbay M, Altunrende F, Ozkuvanci U. Extracorporeal shock wave lithotripsy as first line treatment alternative for urinary tract stones in children: a large scale retrospective analysis. *J Urol*. 2003 Dec;170:2405-8.
35. Shouman AM, Ziada AM, Ghoneim IA, Morsi HA. Extracorporeal shock wave lithotripsy monotherapy for renal stones >25 mm in children. *Urology*. 2009 Jul;74(1):109-11.
36. Wu HY, Docimo SG: Surgical management of children with urolithiasis. *Urol Clin North Am*. 2004; 31 (3):589-594.
37. Muslumanoglu AY, Tefekli A, Sarilar O, et al.: Extracorporeal shock wave lithotripsy as first line treatment alternative for urinary tract stones in children: a large scale retrospective analysis. *J Urol* 2003, 170:2405–8.
38. Raza A, Turna B, Smith G, et al.: Pediatric urolithiasis: 15 years of local experience with minimally invasive endourological management of pediatric calculi. *J Urol* 2005, 174:682–5.
39. Ather MH, Noor MA: Does size and site matter for renal stones up to 30-mm in size in children treated by extracorporeal lithotripsy? *Urology* 2003, 61:212–5.
40. Rodrigues Netto N Jr, Longo JA, Ikonomidis JA, Rodrigues Netto M: Extracorporeal shock wave lithotripsy in children. *J Urol* 2002, 167:2164–6.
41. Afshar K, McLorie G, Papanikolaou F, et al.: Outcome of small residual stone fragments following shock wave lithotripsy in children. *J Urol* 2004, 172:1600–3.
42. Tan AH, Al-Omar M, Watterson JD, et al.: Results of shockwave lithotripsy for pediatric urolithiasis. *J Endourol* 2004, 18:527–30.
43. Wu HY, Docimo SG: Surgical management of children with urolithiasis. *Urol Clin North Am* 2004, 31:589–94.
44. Lee RS, Passerotti CC, Cendron M, et al. Early results of robot assisted laparoscopic lithotomy in adolescents. *J Urol* 2007; 177:2306–9.
45. Young HH, McKay RW. Congenital valvular obstruction of the prostatic urethra. *Surg Gynecol Obstet* 1929;48:509.
46. Kim SS, Kolon TF, Canter D, White M, Casale P. Pediatric flexible ureteroscopic lithotripsy: the children’s hospital of Philadelphia experience. *J Urol* 2008;180:2616-9.

Reference

47. Ritchey M, Patterson DE, Kelalis PP, Segura JW: A case of pediatric ureteroscopic lasertripsy. *J Urol* 1988, 139:1272–4.
48. Hill DE, Segura JW, Patterson DE, Kramer SA: Ureteroscopy in children. *J Urol* 1990, 144:481–3.
49. Caione P, De Gennaro M, Capozza N, et al.: Endoscopic manipulation of ureteral calculi in children by rigid operative ureterorenoscopy. *J Urol* 1990, 144:484–5.
50. Smaldone MC, Cannon GM Jr, Wu HY, et al.: Is ureteroscopy first line treatment for pediatric stone disease? *J Urol* 2007; 178:2128–31.
51. Traxer O, Lechevallier E, Saussine C: Urétroscopie souple-laser Holmium-YAG : le materiel. *Prog Urol*, 2008; 18, 12: 917-928.
52. Soygur T, Zumrutbas AE, Gulpinar O, et al.: Hydrodilation of the ureteral orifice in children renders ureteroscopic access possible without any further active dilation. *J Urol* 2006, 176:285–7.
53. Bassiri A, Ahmadnia H, Darabi MR, Yonessi M. Transureteral lithotripsy in pediatric practice. *J Endourol* 2002;16: 257–60.
54. Al Busaidy SS, Prem AR, Medhat M. Paediatric ureteroscopy for ureteric calculi: a 4-year experience. *Br J Urol* 1997; 80:797–801.
55. Adanur S, Aydin HR, Ozkaya F, Ziypak T, Polat O. Holmium Laser Lithotripsy with Semi-Rigid Ureteroscopy: A First-Choice Treatment for Impacted Ureteral Stones in Children? *Med Sci Monit.* 2014; 20: 2373–2379
56. Delakas D, Daskalopoulos G, Metaxari M, et al. Management of ureteral stones in pediatric patients. *J Endourol* 2001; 15:675–80.
57. Thomas R, Ortenberg J, Lee BR, Harmon EP. Safety and efficacy of pediatric ureteroscopy for management of calculous disease. *J Urol* 1993;149:1082–4.
58. Scarpa RM, De Lisa A, Porru D, et al. Ureterolithotripsy in children. *Urology* 1995;46:859–62.
59. Reddy PP, Barrieras DJ, Bagli DJ, et al. Initial experience with endoscopic holmium laser lithotripsy for pediatric urolithiasis. *J Urol* 1999;162:1714–6.
60. Al-Busaidy SS, Prem AR, Medhat M: Paediatric ureteroscopy for ureteric calculi: a 4-year experience. *Br J Urol* 1997, 80:797–801.
61. Schuster TG, Russell KY, Bloom DA, et al.: Ureteroscopy for the treatment of urolithiasis in children. *J Urol* 2002, 167:1813.
62. Richter S, Shalev M, Lobik L, et al.: Early postoperative vesicoureteral reflux-a temporary and infrequent complication: prospective study. *J Endourol* 1999, 13:365–6.
63. Barcellos Sampaio FJ, Mandarim-de-Lacerda CA. 3-Dimensional and radiological pelvicaliceal anatomy for endourology. *J Urol.* 1988;140:1352–5.

Reference

64. Gupta M, Ost MC, Shah JB, Mcdougall EM, Smith AD. Percutaneous management of the upper urinary tract. In: Wein, Kavoussi, Novick, Partin, Peter, editors. Campbell-Walsh Urology. 9th ed. Philadelphia: Elsevier Saunders Publication; 2007. p. 1528.
65. Jackman SV, Hedican SP, Peters CA, Docimo SG. Percutaneous nephrolithotomy in infants and preschool age children: experience with a new technique. *Urology* 1998; 52:697–701.
66. Kurzrock EA, Huffman JL, Hardy BE, Fugelso P. Endoscopic treatment of pediatric urolithiasis. *J Pediatr Surg* 1996; 31:1413–6.
67. Mor Y, Elmasry YE, Kellett MJ, Duffy PG. The role of percutaneous nephrolithotomy in the management of pediatric renal calculi. *J Urol* 1997;158(3 Pt 2):1319–21.
68. Zeren S, Satar N, Bayazit Y, et al.: Percutaneous nephrolithotomy in the management of pediatric renal calculi. *J Endourol* 2002, 16:75–78.
69. Desai MR, Kukreja RA, Patel SH, Bapat SD: Percutaneous nephrolithotomy for complex pediatric renal calculus disease. *J Endourol* 2004, 18:23–27.
70. Aron M, Yadav R, Goel R, et al.: Percutaneous nephrolithotomy for complete staghorn calculi in preschool children. *J Endourol* 2005, 19: 968–72.
71. Jackman SV, Hedican SP, Peters CA, Docimo SG: Percutaneous nephrolithotomy in infants and preschool age children: experience with a new technique. *Urology* 1998, 52:697–701.
72. Lojanapiwat B, Prasopsuk S. Upper-pole access for percutaneous nephrolithotomy: Comparison of supracostal and infracostal approaches. *J Endourol*. 2006;20:491–4.
73. Kawahara T, Ito H, Terao H, Yoshida M, Ogawa T, Uemura H, et al. Ureterscopy assisted retrograde nephrostomy: A new technique for percutaneous nephrolithotomy (PCNL) *BJU Int*. 2012;110:588–90.
74. Honey RJ, Wiesenthal JD, Ghiculete D, Pace S, Ray AA, Pace KT. Comparison of supracostal versus infracostal percutaneous nephrolithotomy using the novel prone-flexed patients position. *J Endourol*. 2011;25:947–54.
75. Moraitis K, Philippou P, El-Husseiny T, Wazait H, Masood J, Buchholz N. Simultaneous antegrade/retrograde upper urinary tract access: Bart's modified lateral position for complex upper tract endourologic pathologic features. *Urology*. 2012;79:287–92.
76. Lezrek M, Ammani A, Bazine K, Assebane M, Kasmaoui H, Qarro A, et al. The split-leg modified lateral position for percutaneous renal surgery and optimal retrograde access to the upper urinary tract. *Urology*. 2011;78:217–20.
77. El-Nahas AR, Shokeir AA, El-Assmy AM, Mohsen T, Shoma AM, Eraky I, et al. Post-percutaneous nephrolithotomy extensive hemorrhage: A study of risk factors. *J Urol*. 2007;177:576–9.

Reference

78. Valdivia Uria JG, Valle Gerhold J, Lopez Lopez JA, Villarroja Rodriguez S, Ambroj Navarro C, Ramirez Fabian M, et al. Technique and complications of percutaneous nephroscopy: Experience with 557 patients in the supine position. *J Urol*. 1998;160:1975–8.
79. Juul N, Nielsen V, Torp-Peterson S. Percutaneous balloon catheter nephrostomy guided by ultrasound. Results of a new technique. *Scand J Urol nephrol*. 1985;19:291–4.
80. Tepeler A, Armağan A, Akman T, Polat EC, Ersöz C, Topaktap R, et al. Impact of percutaneous renal access technique on outcomes of percutaneous nephrolithotomy. *J Endourol*. 2012;26:828–33.
81. Chen ML, Shukla G, Jackman SV, Tsao AK, Smaldone MC, Ost MC, et al. Real-time tomographic reflection in facilitating percutaneous access to the renal collecting system. *J Endourol*. 2011;25:743–5.
82. Desai M. Ultrasonography-guided punctures with and without puncture guide. *J Endourol*. 2009;23:1641–3.
83. Hosseini MM, Hassanpour A, Farzan R, Yousefi A, Afrasiabi MA. Ultrasonography-guided percutaneous nephrolithotomy. *J Endourol*. 2009;23:603–7.
84. Basiri A, Ziaee SA, Nasseh H, Kamranmanesh M, Masoudy P, Heidary F, et al. Totally ultrasonography-guided percutaneous nephrolithotomy in the flank position. *J Endourol*. 2008;22:1453–7.
85. Safak M, Olgar T, Bor D, Berkmen G, Gogus C, et al. Radiation doses of patients and urologists during percutaneous nephrolithotomy. *J Radiol Prol*. 2009;29:409–15.
86. Majidpour HS. Risk of radiation exposure during PCNL. *Urol J*. 2010;7:87–9.
87. Tepeler A, Binbay M, Yuruk E, Sari E, Kaba M, Muslumanoglu AY, et al. Factors affecting the fluoroscopic screening time during percutaneous nephrolithotomy. *J Endourol*. 2009;23:1825–9.
88. Tepeler A, Armağan A, Akman T, Polat EC, Ersöz C, Topaktap R, et al. Impact of percutaneous renal access technique on outcomes of percutaneous nephrolithotomy. *J Endourol*. 2012;26:828–33.
89. Soria F, Delgado MI, Sánchez FM, Allona A, Jiménez Cruz JF, Morell E, et al. Effectiveness of three-dimensional fluoroscopy in percutaneous nephrostomy: An animal model study. *Urology*. 2009;73:649–52.
90. Andonian S, Scoffone CM, Louie MK, Gross AJ, Grabe M, Daels FP, et al. Does Imaging Modality Used for Percutaneous Renal Access Make a Difference. A Matched Case Analysis? *J Endourol*. 2013;27:24–8.
91. Spirnak JP, Resnick MI. Retrograde percutaneous stone removal using modified Lawson technique. *Urology*. 1987;30:551–3.
92. Grasso M, Lang G, Taylor FC. Flexible ureteroscopically assisted percutaneous renal access. *Tech Urol*. 1995;1:39–43.

Reference

93. Bader MJ, Gratzke C, Seitz M, Sharma R, Stief CG, Desai M. The “all-seeing needle”: Initial results of an optical puncture system confirming access in percutaneous nephrolithotomy. *Eur Urol.* 2011;59:1054–9.
94. Tefekli A, Altunrende F, Tepeler K, Tas A, Aydin S, Muslumanoğlu AY. Tubeless percutaneous nephrolithotomy in selected patients: A prospective randomized comparison. *Int Urol Nephrol.* 2007;39:57–63.
95. Dwaba MS, Shokeir AA, Hafez AT, et al.: Percutaneous nephrolithotomy in children: early and late anatomical and functional results. *J Urol* 2004, 172:1078–81.
96. Salah MA, Tallai B, Holman E, et al.: Simultaneous bilateral percutaneous nephrolithotomy in children. *BJU Int* 2005, 95:137–9.
97. Lingeman J, Matlaga B, Evan A. Surgical management of upper urinary tract calculi. In: Wein AJ, Kavoussi LR, Novick AC, Partin AW, Peters CA (eds). *Cambell-Walsh Urology*, 9th edn. Saunders Elsevier, Philadelphia, PA, 2007; 1437–8.
98. European Association of Urology (EAU) Guidelines. Guidelines on urolithiasis. EAU 2011.
99. Unsal A, Resorlu B. Retrograde intrarenal surgery in infants and preschool-age children. *J Pediatr Surg.* 2011;46:2195-9.
100. Abu Ghazaleh LA et al. Retrograde intrarenal lithotripsy for small renal stones in prepubertal children. *Saudi J Kidney Dis Transpl.* 2011;22(3):492-6.
101. Salerno A et al. Treatment of pediatric renal stones in a Western country: a changing pattern. *J Pediatr Surg.* 2013;48:835-9.
102. Tekgöl S et al. Guideline on pediatric urology. 2013:1-126. Available at: http://www.uroweb.org/gls/pdf/22%20Paediatric%20Urology_LR.pdf.
103. Unsal A et al. Bilateral böbrek ve üreterinde sistin taşları olan 9 aylık infantta anegrad perkütan yaklaşım. *Türk Üroloji Dergisi.* 2009;35:56-60.
104. Wadhwa P et al. Critical prospective appraisal of renal morphology and function in children undergoing shockwave lithotripsy and percutaneous nephrolithotomy. *J Endourol.* 2007;21: 961-6.
105. Özden E et al. Percutaneous renal surgery in children with complex stone. *J Pediatr Urol.* 2008;4:295-8.
106. Grasso M, Conlin M, Bagley D. Retrograde ureteropyeloscopic treatment of 2 cm or greater upper urinary tract and minor staghorn calculi. *J. Urol.* 1998; 160: 346–51.
107. Breda A, Ogunyemi O, Leppert JT, Lam JS, Schulam PG. Flexible ureteroscopy and laser lithotripsy for intrarenal stones 2 cm or greater – Is this the new frontier? *J. Urol.* 2008; 179: 981–4.
108. Riley JM, Stearman L, Troxel S. Retrograde ureteroscopy for renal stones larger than 2.5 cm. *J. Endourol.* 2009; 23: 1395–8.

Reference

109. Ricchiuti DJ, Smaldone MC, Jacobs BL, Smaldone AM, Jackman SV, Averch TD. Staged retrograde endoscopic lithotripsy as alternative to PCNL in select patients with large renal calculi. *J. Endourol.* 2007; 21: 1421–4.
110. Hyams ES, Munver R, Bird VG, Uberoi J, Shah O. Flexible ureteroscopy and holmium laser lithotripsy for the management of renal stone burdens that measure 2 to 3 cm: a multiinstitutional experience. *J. Endourol.* 2010; 24:1583–8.
111. Takazawa R, Kitayama S, Tsujii T. Successful outcome of flexible ureteroscopy with holmium laser lithotripsy for renal stones 2 cm or greater. *International journal of Urology.* 2012 March;19(3):264-267.
112. Yeow WC, Pemberton R, Barker A. Flexible ureterorenoscopy and laser lithotripsy in children. *J Indian Assoc Pediatr Surg.* 2009 Apr;14(2):63-5.
113. Kim SS, Kolon TF, Canter D, White M, Casale P. Pediatric flexible ureteroscopic lithotripsy: the children's hospital of philadelphia experience. *J Urol.* 2008 Dec;180(6):2616-9.
114. Kotz S, Balakrishnan N, Read CB, Vidakovic B. *Encyclopedia of statistical sciences.* 2nd ed. Hoboken, N.J.: Wiley-Interscience; 2006.
115. Kirkpatrick LA, Feeney BC. *A simple guide to IBM SPSS statistics for version 20.0.* Student ed. Belmont, Calif.: Wadsworth, Cengage Learning; 2013.
116. Afshar K, McLorie G, Papanikolaou F, Malek R, Harvey E, Pippi-Salle JL, Bagli DJ, Khoury AE, Farhat W. Outcome of small residual stone fragments following shock wave lithotripsy in children. *J Urol.* 2004 Oct; 172(4 Pt 2):1600-3.
117. Jayanthi VR, Arnold PM, Koff SA. Strategies for managing upper tract calculi in young children. *J Urol.* 1999 Sep; 162(3 Pt 2):1234-7.
118. Muslumanoglu AY, Tefekli A, Sarilar O, Binbay M, Altunrende F, Ozkuvanci U. Extracorporeal shock wave lithotripsy as first line treatment alternative for urinary tract stones in children: a large scale retrospective analysis. *J Urol.* 2003 Dec; 170(6 Pt 1):2405-8.
119. Aksoy Y, Ozbey I, Atmaca AF, Polat O. Extracorporeal shock wave lithotripsy in children: experience using a mpl-9000 lithotripter. *World J Urol.* 2004 Jun; 22(2):115-9.
120. Strauss KJ, Kaste SC. The ALARA (as low as reasonably achievable) concept in pediatric interventional and fluoroscopic imaging: striving to keep radiation doses as low as possible during fluoroscopy of pediatric patients--a white paper executive summary. *Radiology.* 2006 Sep; 240(3):621-2.
121. Krambeck AE, Gettman MT, Rohlinger AL, Lohse CM, Patterson DE, Segura JW. Diabetes mellitus and hypertension associated with shock wave lithotripsy of renal and proximal ureteral stones at 19 years of followup. *J Urol.* 2006 May; 175(5):1742.
122. Straub M, Gschwend J, Zorn C. Pediatric urolithiasis: the current surgical management. *Pediatr Nephrol.* 2010 Jul; 25(7):1239-44.

Reference

123. Desai MR, Kukreja RA, Patel SH, Bapat SD. Percutaneous nephrolithotomy for complex pediatric renal calculus disease. *J Endourol.* 2004;18:23–7.
124. Desai M. Endoscopic management of stones in children. *Curr Opin Urol.* 2005;15:107–12.
125. Jackman SV, Hedican SP, Peters CA, Docimo SG. Percutaneous nephrolithotomy in infants and preschool age children: Experience with anew technique. *Urology.* 1998;52:697–701.
126. Ritchey M, Patterson DE, Kelalis PP, Segura JW. A case of pediatric ureteroscopic lasertripsy. *J Urol.* 1988;139:1272–4.
127. Schuster TG, Russell KY, Bloom DA, Koo HP, Faerber GJ. Ureteroscopy for the treatment of urolithiasis in children. *J Urol.* 2002;167:1813.
128. Lesani OA, Palmer JS. Retrograde proximal rigid ureteroscopy and pyeloscopy in prepubertal children: Safe and effective. *J Urol.* 2006;176:1570–3.
129. De Dominicis M, Matarazzo E, Capozza N, Collura G, Caione P. Retrograde ureteroscopy for distal ureteric stone removal in children. *BJU Int.* 2005;95:1049–52.
130. Raza A, Turna B, Smith G, Moussa S, Tolley DA. Pediatric urolithiasis: 15 years of local experience with minimally invasive endourological management of pediatric calculi. *J Urol.* 2005;174:682–5.
131. El-Hout Y, Elnaeema A, Farahat WA. Current status of retrograde intrarenal surgery for management of nephrolithiasis in children. *Indian J Urol.* 2010 Oct-Dec; 26(4): 568–572.
132. Tan AH, Denstedt JD, Razvi H. Ureteroscopy for pediatric urolithiasis: An evolving first-line therapy. *Pediatric Urol.* 2004;65:153–6. [PubMed]
133. Minevich E, Defoor W, Reddy P, Nishinaka K, Wacksman J, Sheldon C, et al. Ureteroscopy is safe and effective in prepubertal children. *J Urol.* 2005;174:276–9.
134. Thomas JC, DeMarco RT, Donohoe JM, Adams MC, Brock JW, 3rd, Pope JC., 4th Pediatric ureteroscopic stone management. *J Urol.* 2005;174:1072–4.
135. Sofer M, Binyamini J, Ekstein PM, Bar-Yosef Y, Chen J, Matzkin H, et al. Holmium laser ureteroscopic treatment of various pathologic features in pediatrics. *Urology.* 2007;69:566–9.
136. Smaldone MC, Cannon GM GM, Jr, Wu HY, Bassett J, Polsky EG, Bellinger MF, et al. Is ureteroscopy first line treatment for pediatric stone disease? *J Urol.* 2007;178:2128–31.
137. Cannon GM, Smaldone MC, Wu HY, Bassett JC, Bellinger MF, Docimo SG, et al. Ureteroscopic management of lower pole stones in a pediatric population. *J Endourol.* 2007;21:1179–82.
138. Dave S, Khoury AE, Braga L, Farhat WA. Single-institutional study on role of ureteroscopy and retrograde intrarenal surgery in treatment of pediatric renal.

Reference

- Urology. 2008;72:1018–21
139. Kim SS, Kolon TF, Canter D, White M, Casale P. Pediatric flexible ureteroscopic lithotripsy: The children's hospital of Philadelphia experience. *J Urol*. 2008;180:2616–9.
 140. Tanaka ST, Makari JH, Pope JC, 4th, Adams MC, Brock JW, 3rd, Thomas JC. Pediatric ureteroscopic management of intrarenal calculi. *J Urol*. 2008;180:2150–3.
 141. Bassiri A, Ahmadnia H, Darabi MR, Yonessi M. Transureteral Lithotripsy in Pediatric Practice. *J Endourol*. 2002;16:257–60.
 142. Herndon CD, Viamonte L, Joseph DB. Ureteroscopy in children: Is there a need for ureteral dilation and postoperative stenting? *J Pediatr Urol*. 2006;2:290–3.
 143. Smaldone MC, Corcoran AT, Docimo SG, et al. Endourological management of pediatric stone disease: present status. *J Urol*. 2009; 181:17-28.
 144. Ozden E, Sahin A, Tan B, et al. Percutaneous renal surgery in children with complex stones. *J Pediatr Urol*. 2008; 4:295-298.
 145. Traxer O, Smith TG, Pearle MS, et al. Renal parenchymal injury after standard and mini percutaneous nephrostolithotomy. *J Urol*. 2001;165:1693-5.
 146. Bilen CY, Kocak B, Kıtırcı G, et al. Percutaneous nephrolithotomy in children: lessons learned in 5 years at a single institution. *J Urol*. 2007;177:1867-71.
 147. Zeren S, Satar N, Bayazit Y, Bayazit AK, Payasli K, Ozkeçeli R. Percutaneous nephrolithotomy in the management of pediatric renal calculi. *J Endourol*. 2002;16:75–8
 148. Lee KL, Stoller ML. Minimizing and managing bleeding after percutaneous nephrolithotomy. *Curr Opin Urol*. 2007;17:120–4.
 149. Michel MS, Trojan L, Rassweiler JJ. Complications in percutaneous nephrolithotomy. *Eur Urol*. 2007;51:899–906.
 150. Desai M, Ridhorkar V, Patel S, Bapat S, Desai M. Factors affecting blood loss during percutaneous nephrolithotomy: prospective study. *Journal of Endourology*. June 1999, 13(5): 359-364.
 151. Elghoneimy, T.Z. Orban, K. Mursi. A. Hegazy. Safety and efficacy of supracostal access in percutaneous renal surgery. *J of Endourology*; Jan 2008, Vol. 22 Issue 1, p29 .
 152. Mousavi-Bahar S, Mehrabi S, Moslemi M. The safety and efficacy of PCNL with supracostal approach in the treatment of renal stones. *International Urology & Nephrology*; Dec2011, Vol. 43 Issue 4, p983.
 153. El-Nahas AR, Shokeir AA, El-Kenawy MR, Shoma AM, Eraky I, El-Assmy AM, Ghaly AM, El-Kappany HA. Safety and efficacy of supracostal percutaneous nephrolithotomy in pediatric patients. *J Urol*. 2008 Aug;180(2):676-80.

Reference

154. Anand A, Kumar R, Dogra PN, Seth A, Gupta NP. Safety and efficacy of a superior calyceal puncture in pediatric percutaneous nephrolithotomy. *J of Endourol.* 2010;24(11):1725-8.
155. Roberts S, Bolton DM, Stoller ML. Hypothermia associated with percutaneous nephrolithotomy. *Urology.* 1994;44:832-5.
156. El-Nahas AR, Mansour AM, Ellaithy R, Abol-Enein H. Case Report: Conservative Treatment of Liver Injury during Percutaneous Nephrolithotomy *J of Endourol.* August 2008, 22(8): 1649-1652.
157. Thomas AA, Pierce G, Walsh RM, Sands M, Noble M. Splenic injury during percutaneous nephrolithotomy. *JSLs.* 2009 Apr-Jun;13(2):233-6.
158. Desai AC, Jain S, Benway BM, Grubb RL 3rd, Picus D, Figenshau RS. Splenic injury during percutaneous nephrolithotomy: a case report with novel management technique. *J of Endourol.* 2010 Apr;24(4):541-5.
159. AslZare M, Darabi MR, Shakiba B, Gholami-Mahtaj L. Colonic perforation during percutaneous nephrolithotomy: An 18-year experience. *Can Urol Assoc J.* 2014 May-Jun; 8(5-6).
160. Saad KS, Hanno A, El-Nahas AR. Injury of the ileum during percutaneous nephrolithotomy in a pediatric patient. *Can Urol Assoc J.* 2014 Mar-Apr;8(3-4):E204-6.
161. Tuttle DN, Yeh BM, Meng MV, et al. Risk of injury to adjacent organs with lower-pole fluoroscopically guided percutaneous nephrostomy: Evaluation with prone, supine, and multiplanar reformatted CT. *J Vasc Interv Radiol.* 2005;16:1489-92.
162. Culkin DJ, Wheeler JS, Jr, Canning JR. Nephro-duodenal fistula: A complication of percutaneous nephrolithotomy. *J Urol.* 1985;134:528-30.
163. Traxer O. Management of injury to the bowel during percutaneous stone removal. *J Endourol.* 2009;23:1777-80.
164. Shah HN, Kausik VB, Hegde SS, Shah JN, Bansal MB. Tubeless percutaneous nephrolithotomy: A prospective feasibility study and review of previous reports. *BJU Int.* 2005;96:879-83.
165. Desai MR, Kukreja RA, Desai MM, Mhaskar SS, Wani KA, Patel SH, et al. A prospective randomized comparison of type of nephrostomy drainage following percutaneous nephrostolithotomy: Large bore versus small bore versus tubeless. *J Urol.* 2004 Aug;172(2):565-7.

المخلص العربي

منذ أوائل ١٩٨٠، عندما أدخل منظار الكلي و منظار الحالب لتفتيت و استخراج الحصوات، و مع استحداث تقنية تفتيت الحصوات بواسطة الموجات التصادمية من خارج الجسم، حلت هذه الإجراءات محل العمليات الجراحية المفتوحة في المرضى البالغين.

وقد تم هذا التقدم التكنولوجي بصورة سريعة و كبيرة في المرضى البالغين بينما تم تطبيقه متأخرا على الأطفال، و ذلك في المقام الأول بسبب القيود التقنية المرتبطة بعدم توافر المناظير و الادوات التي تتناسب و صغر حجم المريض، وثانيا بسبب ندرة حالات حصوات المسالك البولية في الأطفال.

و تعد حصوات المسالك البولية في مرحلة الطفولة أمر نادر الحدوث في العالم المتقدم، وهو ما يمثل ١٪ الى ٥٪ من جميع حصوات المسالك البولية بينما تمثل حالات حصوات المسالك البولية في الأطفال في البلدان النامية نسبة قد تصل الي ٣٠٪ من جميع حصوات المسالك البولية و على الرغم من أن الإحصاءات الدقيقة غير متوفرة من البلدان النامية، تعد حصوات المسالك البولية في الأطفال أكثر شيوعا مما هي عليه في البلدان الصناعية.

و قد قمنا في هذه الدراسة، بتقييم مستقبلي لتجربتنا في علاج الحصوات باستخدام الطرق الاقل تداخلا في الأطفال الذين تم علاجهم بقسم جراحة المسالك البولية، جامعة الإسكندرية، حيث تم تقسيم المرضى إلى مجموعتان خضعوا لتفتيت و استخراج حصوات المسالك البولية و ذلك باستخدام منظار الكلي او منظار الحالب لتفتيت و استخراج الحصوات، هذه بالإضافة لاستخدام التفتيت بواسطة الموجات التصادمية.

و تم علاج ١٧ طفلا (٢١ وحدة الكلي) بمنظار الحالب المرن، وقد تمت بنجاح تماما في جلسة واحدة في ١٥ حالة (٧١.٤٪). و قد تم إجراء منظار الحالب دون توسيع الحالب في جميع الحالات. وتعرض اثنان من المرضى لارتفاع حرارة خفيفة إلى معتدلة بعد المنظار الجراحي.

وخضع واحد وعشرين طفلا (٢٢ وحدة الكلي) إلى منظار الكلي. وقد تمت بنجاح تام بمعدل ٩٥.٥٪ من الحالات، مع نسبة المضاعفات ٤١٪، بما في ذلك نزيف في ١٣٪، حمى معتدلة في ١٨٪.

وقد أظهرت نتائج هذه الدراسة أن استخدام الطرق الاقل تداخلا في الأطفال هو وسيلة آمنة وفعالة لعلاج حصوات المسالك البولية حيث أنه يقلل من معدلات الاعتلال والبقاء في المستشفى، كما تبين من إجراء هذه الدراسة ان الحصوات في اغلبها كانت مكونة من اوكسالات الكالسيوم في غالبية الأطفال ، وكان ازدياد نسبة الكالسيوم في البول هو عامل الخطر الأكثر شيوعا في المجموعة البحثية.



جامعة الإسكندرية
كلية الطب
قسم جراحة المسالك البولية و التناسلية

دراسة مستقبلية لمقارنة امان و فاعلية استخراج حصوات الكلى بالمنظار عن طريق الجلد مقابل المنظار الصاعد عن طريق الحالب فى حالات الأطفال ذوى الحصوات الكبيرة والمتعددة

رسالة مقدمة

لقسم جراحة المسالك البولية و التناسلية - كلية الطب - جامعة الإسكندرية
ضمن متطلبات درجة

الدكتوراه

فى

جراحة المسالك البولية و التناسلية

من

كريم سعد محمد سعد

بكالوريوس الطب والجراحة، ٢٠٠٢
ماجستير جراحة المسالك البولية، جامعة الإسكندرية - ٢٠٠٩
كلية الطب، جامعة الإسكندرية

[٢٠١٥]



جامعة الإسكندرية
كلية الطب
قسم جراحة المسالك البولية و التناسلية

دراسة مستقبلية لمقارنة امان و فاعلية استخراج حصوات الكلى بالمنظار
عن طريق الجلد مقابل المنظار الصاعد عن طريق الحالب فى حالات الأطفال
ذوى الحصوات الكبيرة والمتعددة

رسالة مقدمة من

كريم سعد محمد سعد

للحصول على درجة

الدكتوراه

فى

جراحة المسالك البولية و التناسلية

التوقيع

لجنة المناقشة والحكم على الرسالة

.....

أ.د/ عصام مرزوق
أستاذ جراحة المسالك البولية والتناسلية
كلية الطب
جامعة الإسكندرية

.....

أ.د/ أحمد جمال الدين هنو
أستاذ جراحة المسالك البولية والتناسلية
كلية الطب
جامعة الإسكندرية

.....

أ.م.د/ أحمد رفعت النحاس
أستاذ مساعد جراحة المسالك البولية والتناسلية
كلية الطب
جامعة المنصورة

التاريخ / /

لجنة الإشراف

موافقون

أ.د/ . احمد جمال الدين هنو

أستاذ جراحة المسالك البولية والتناسلية
كلية الطب
جامعة الإسكندرية

.....

أ.د/ سيف الأسلام محمود نفيس حمدى

أستاذ جراحة المسالك البولية والتناسلية
كلية الطب
جامعة الإسكندرية

.....

أ.م.د/ محمد السيد يوسف

أستاذ مساعد جراحة المسالك البولية والتناسلية
كلية الطب
جامعة الإسكندرية

.....