
LIST OF FIGURES

Figure (1)	Shape of a surfactant	9
Figure (2)	Corrosion arrested by release of corrosion inhibitor	26
Figure (3)	FTIR-spectrum of 6-(3-amino phenoxy) hexane-1-thiol(C6M)	56
Figure (4)	FTIR-spectrum of 8-(3-amino phenoxy) octane-1-thiol (C8M)	56
Figure (5)	FTIR-spectrum of 10-(3-amino phenoxy) decane-1-thiol (C10M)	57
Figure (6)	FTIR-spectrum of 12-(3-amino phenoxy) dodecane-1-thiol (C12M)	57
Figure (7):	¹ HNMR-spectrum of 6-(3-amino phenoxy) hexane-1-thiol (C6M)	59
Figure (8):	¹ HNMR-spectrum of 8-(3-amino phenoxy) octane thiol (C8M)	59
Figure (9):	¹ HNMR-spectrum of 10-(3-amino phenoxy) decane-1-thiol (C10M)	60
Figure (10):	¹ HNMR-spectrum of 12-(3-amino phenoxy) dodecane-1-thiol (C12M)	60
Figure (11)	FTIR-spectrum of poly 6-(3-amino phenoxy) hexane-1-thiol (C6P)	63
Figure (12)	FTIR-spectrum of poly 8-(3-amino phenoxy) octane-1-thiol (C8P)	63
Figure (13)	FTIR-spectrum of poly 10-(3-amino phenoxy) decane-1-thiol (C10P)	64

Figure (14)	FTIR-spectrum of poly 12-(3-amino phenoxy) dodecane-1- thiol (C12P)	64
Figure (15)	TGA &DSC Chart of poly 6-(3-amino phenoxy) hexane-1- thiol (C6P)	66
Figure (16)	TGA &DSC Chart of poly 8-(3-amino phenoxy) octane-1- thiol (C8P)	67
Figure (17)	TGA &DSC Chart of poly 10-(3-amino phenoxy) decane-1- thiol (C10P)	68
Figure (18)	TGA &DSC Chart of poly 12-(3-amino phenoxy) dodecane-1- thiol (C12P)	69
Figure (19)	XRD spectra of poly 6-(3-amino phenoxy) hexane-1- thiol (C6P)	70
Figure (20)	XRD spectra of poly 8-(3-amino phenoxy) octane-1- thiol (C8P)	71
Figure (21)	XRD spectra of poly 10-(3-amino phenoxy) decane-1- thiol (C10P)	71
Figure (22)	XRD spectra of poly 12-(3-amino phenoxy) dodecane-1- thiol (C12P)	72
Figure (23)	FTIR-spectrum of 6-(3-amino phenoxy) hexane-1- thiol (C6M) with AgNPs	74
Figure (24)	FTIR-spectrum of 8-(3-amino phenoxy) octane-1- thiol(C8M) with AgNPs	74
Figure (25)	FTIR-spectrum of 10-(3-amino phenoxy) decane-1- thiol (C10M) With AgNPs	75
Figure (26)	FTIR-spectrum of 12-(3-amino phenoxy) dodecane-1- thiol (C12M) With AgNPs	75
Figure (27)	UV spectra of prepared silver nanoparticles (AgNPs)	76

Figure (28)	UV-spectrum of 6-(3-amino phenoxy) hexane-1-thiol with AgNPs (C6M+AgNPs)	77
Figure (29)	UV -spectrum of 8-(3-amino phenoxy) octane-1-thiol with AgNPs (C8M+AgNPs)	77
Figure (30)	UV -spectrum of 10-(3-amino phenoxy) decane-1-thiol With AgNPs(C10M+AgNPs)	78
Figure (31)	UV -spectrum of 12-(3-amino phenoxy) dodecane-1- thiol With AgNPs (C12M+AgNPs)	78
Figure (32)	TEM imageof silver nanoparticles (AgNPs)	79
Figure (33)	TEM imageof 6-(3-amino phenoxy) hexane-1-thiol (C6-M) with AgNP	80
Figure (34)	TEM image of 8-(3-amino phenoxy). octane-1-thiol with AgNPs (C8M+AgNPs)	80
Figure (35)	TEM imageof 10-(3-amino phenoxy) decane-1-thiol With AgNPs (C10-M+AgNPs)	81
Figure (36)	TEM imageof 12-(3-amino phenoxy) dodecane-1-thiol With AgNPs (C12M+AgNPs)	81
Figure (37)	XRD spectra of prepared Silver nanoparticles (AgNPs)	82
Figure (38)	XRD spectra of 6-(3-amino phenoxy) hexane-1-thiol with AgNPs (C6M+AgNPs)	83
Figure (39)	XRD spectra of 8-(3-amino phenoxy) octane-1-thiol with AgNPs (C8M+ AgNPs)	83
Figure (40)	XRD spectra of 10-(3-amino phenoxy) decane-1-thiol with AgNPs (C10M+AgNPs)	84
Figure (41)	XRD spectra of 12-(3-amino phenoxy) dodecane-1- thiol With AgNPs (C12M+AgNPs)	84

Figure (42)	FTIR-spectrum of poly 6-(3-amino phenoxy) hexane-1- thiol with AgNPs (C6P+AgNPs)	86
Figure (43)	FTIR-spectrum of poly 8-(3-amino phenoxy) octane-1- thiol with AgNPs (C8P+AgNPs)	86
Figure (44)	FTIR-spectrum of poly 10-(3-amino phenoxy) decane-1- thiol With AgNPs (C10P+AgNPs)	87
Figure (45)	FTIR-spectrum of poly 12-(3-amino phenoxy) dodecane-1- thiol with AgNPs (C12P+AgNPs)	87
Figure (46)	UV-spectrum of poly 6-(3-amino phenoxy) hexane-1- thiol with AgNPs (C6P+AgNPs)	88
Figure (47)	UV-spectrum of poly 8-(3-amino phenoxy) octane-1- thiol with AgNPs (C8P+AgNPs)	88
Figure (48)	UV-spectrum of poly 10-(3-amino phenoxy) decane-1- thiol With AgNPs (C10P+AgNPs)	89
Figure (49)	UV-spectrum of poly 12-(3-amino phenoxy) dodecane-1- thiol with AgNPs (C12P+AgNPs)	89
Figure (50)	TEM imageof poly 6-(3-amino phenoxy) hexane-1- thiol with AgNPs (C6P+AgNPs)	90
Figure (51)	TEM imageof poly 8-(3-amino phenoxy) octane-1- thiol with AgNPs (C8P+AgNPs)	91
Figure (52)	TEM imageof poly 10-(3-amino phenoxy) decane-1- thiol With AgNPs (C10P+AgNPs)	91
Figure (53)	TEM imageof poly 12-(3-amino phenoxy) dodecane-1- thiol with AgNPs (C12P+AgNPs)	92
Figure (54)	XRD-spectrum of poly 6-(3-amino phenoxy) hexane-1- thiol with AgNPs (C6P+AgNPs)	93
Figure (55)	XRD-spectrum of poly 8-(3-amino phenoxy) octane-1- thiol with AgNPs (C8P+AgNPs)	94

Figure (56)	XRD-spectrum of poly 10-(3-amino phenoxy) decane-1- thiol With AgNPs (C10P+AgNPs)	94
Figure (57)	XRD-spectrum of poly 12-(3-amino phenoxy) dodecane-1- thiol with AgNPs (C12P+AgNPs)	95
Figure (58)	Relationship between surface tension and – logarithm concentration of the synthesized monomeric thiol surfactants	98
Figure (59)	Relationship between surface tension and – logarithm concentration of the synthesized monomeric thiol surfactants with silver nanoparticles	98
Figure (60)	Relationship between surface tension and –logarithm concentration of the synthesized polymeric thiol surfactants	99
Figure (61)	Relationship between surface tension and – logarithm concentration of the synthesized polymeric thiol surfactants with silver nanoparticles	99
Figure (62)	Relationship between corrosion rate of steel and concentration of the synthesized monomeric thiol surfactants	105
Figure (63)	Relationship between corrosion rate of steel and concentration of the synthesized monomeric thiol surfactants with silver nanoparticles	105
Figure (64)	Relationship between corrosion rate of steel and concentration of the synthesized polymeric thiol surfactants	106

Figure (65)	Relationship between corrosion rate of steel and concentration of the synthesized polymeric thiol surfactants with silver nanoparticles	106
Figure (66)	The Relation between inhibition efficiency and concentration of the synthesized monomeric thiol surfactants in 1 M HCl for the carbon steel sample 25°C.	112
Figure (67)	The Relation between inhibition efficiency and concentration of the synthesized monomeric thiol surfactants with silver nanoparticles in 1 M HCl for the carbon steel sample at 25°C	112
Figure (68)	The Relation between inhibition efficiency and concentration of the synthesized polymeric thiol surfactants in 1 M HCl for the carbon steel sample at 25°C.	113
Figure (69)	The relation between inhibition efficiency and concentration of the synthesized polymeric thiol surfactants with silver nanoparticles in 1 M HCl for the carbon steel sample at 25°C.	113
Figure (70)	Tafel line for the carbon steel sample in 1 M HCl and in presence different concentrations compound C12M	116
Figure (71)	Tafel line for the carbon steel sample in 1 M HCL and in presence different concentrations compound C12M+AgNP	117
Figure (72)	Tafel line for the carbon steel sample in 1 M HCL and in presence different concentrations compound (C12P)	118

- Figure (73)** Tafel line for the carbon steel sample in 1 M HCL and in presence different concentrations compound (C12P+AgNPs). 119
- Figure (74)** Energy dispersive X-rays analysis (EDAX) of carbon steel samples: (a) After immersion in 1 M HCl solution for 24 h. (b) After immersion in 1 M HCl solution for 24 h in the presence of C12M. (C) After immersion in 1 M HCl solution for 24 h in the presence of C12M self-assembled on silvernanoparticles. 121
- Figure (75)** Energy dispersive X-rays analysis (EDAX) of carbon steel samples: (a) after immersion in 1 M HCl solution for 24 h in the presence of C12P. (b) after immersion in 1 M HCl solution for 24 h in the presence of C12P self-assembled on silver nanoparticles. 122
-