

S^eNSITIVITY OF ANABANA ARYZAE AND TOLYPOTHRIX
TENUIS (CYANODACTERIA) TO POLYTRIN INSECTICIDE

-17-

Mehreshan T.El-Mokadem, Gahiza A. Ismail
and Alia A. El Shimy

Bot. Dep. Women's College, Ain Shams Univ.
Heliopolis, Cairo, Egypt

Summary

The effect of increasing concentration of the insecticide Polytrin (0.063 -----1.5 recommended field rate) on the growth of cyanobacteria Anabaena oryzae and Tolypothrix tenuis under aseptic conditions was studied. Percentage dry weight compared to control, total carbohydrates and intracellular nutrient mineral content have been estimated.

addition of Polytrin to growing cultures of A.oryzae and T.Tenuis resulted in a slight reduction of growth at low concentrations, but rapid decrease in growth was observed at higher rates for both strains. The growth of Tolypothrix tenuis was inhibited completely at 1.5 field rate.

Nitogen content of both strains were gradually reduced by increasing the insecticide concentration, while carbohydrate content of algal cells increased.

The intracellular K content of A. oryzae showed slight rise at low insecticide concentrations but a marked reduction at higher concentrations while a marked reduction in K content of T.tenuis was noticed even at the least used concentration.

Slight rise in P and Mg content was observed at low rate followed by decrease at higher applications in both strains. Na did not show wide change by the insecticide. Zn, Cu and Mn content may show some little rise with Polytrin concentration in both organisms.

Introduction

The successful use of algal inoculation as in India and Japan demonstrates that blue-green (BGA) can serve as an alternative or supplemental source of nitrogen for rice cultivation, (Agrwal, 1979; Alyer, et al, 1972; Watanabe, 1967, and Roger & Kuhsooriya, 1980). Blue-green algae may give some advantages not necessarily associated with nitrogen-fixation, such as through the production of growth-promotive substances (Roger and Kulasooriya, 1980).

From the various agronomic practice adopted along the cultivation cycles which may directly influence the growth of blue green algae is the application of

pesticides. The resistance to high levels of pesticides seems to be more characteristic of BGA than of the eukaryotic algae (Watanabe, 1967, Venkataraman & Kajyalakshmi, 1971, 1972; Singh, 1978). But some pesticides like chloropicrin may affect all algae without discrimination (Ishizawa and Musuguchi, 1966). Laboratory experiments showed that, metabolic products of the insecticides Aldrin, Dieldrin and endrin are inhibitory to blue-green algae. (Battino-Viterbo et al., 1973). Also, Wright et al., (1977) showed that, 5 ppm of the herbicide propanil prevented the growth of Anabaena cylindrica, Tolypothrix tenuis and Nostoc endophytum in flask cultures.

Although a number of studies have investigated the growth responses of cyanobacteria to pesticides, available literature shows that, there are only few attempts to define systems for the study of the effect of pesticide on the intracellular mineral constituents of cyanobacteria.

The purpose, of the present study was to assess the changes in growth, intracellular carbohydrates, macro- and microelement of the cyanobacteria Anabaena oryzae and Tolypothrix tenuis when subjected to different concentrations of the foliar pyrethroid insecticide, Polytrin, in batch culture grown photoautotrophically in continuous illumination.

Material and Methods

Cyanobacteria:

Anabaena aryzae, Egyptian strain and Tolypothrix tenuis, Japanese strain were provided from the Agriculture Research Centre Giza, Egypt.

Insecticide:

Plytrin, common name Cypermethrin CGA 55' 186, 200 Emulsifiable concentrate (200 EC) was provided from Ciba-Geigy Limited, Cairo, Egypt. It is a foliar pyrethroid insecticide, its imperical formula is $C_{22}H_{19}Cl_2NO_3$.

Experimental procedure:

Cyanobacteria were grown in batch cultures under sterile conditions. Inocula were grown to mid-exponential phase. Aliquots of (5 ml) cell suspensions were inoculated into 500 ml Erlenmeyer flasks containing 200 ml of (Watanabe et al, 1959) medium. Cultures were grown under conditions of continuous illumination at an incident light intensity of 20 W/m² provided by white fluorescent lamps. The insecticide Polytrin was added at time-zero to cyanobacterial cultures, the concentrations used were 0.063 (1/16), 0.125 (1/8), 0.25 (1/4) 0.5 (1/2), 1.0 and 1.5 recommended field rate (0.5 ml 200 EC/L). Experiments were repeated at least twice.

Cells were harvested by centrifugation at the end of the exponential growth phase, the harvested cells were washed twice with distilled water.

Growth was determined in terms of (mg dry wt/100 ml culture) following drying for 24 hours at 70°C.

Analytical methods:

Total carbohydrates were estimated by the phenol sulphuric acid reagent and calibrated against glucose standard (Dubois et al, 1956).

Intracellular ion concentrations were analyzed from the dried algal cells which were powdered and digested in nitric-perchloric-sulphuric acid. Phosphorus was analyzed using the vanadatomolybdate colorimetric method, (Chapman and Prott, 1978).

Mg, Fe, Mn, Zn, and Cu were determined by atomic absorption spectrophotometry. K and Na were estimated by flame photometry (Jackson, 1967).

Nitrogen was measured by kjeldahl method (Jackson, 1967).

Results

Effect of Polytrin on the growth of algae:

Growth of Anabaena oryzae and Tolypothrix tenuis in the presence of different insecticide concentrations in the medium (Table 1) showed that, growth was slightly

reduced at low insecticide concentrations (0.063 & 0.125 of the recommended field rate) 40.4% - 72.2% of control culture, but rapid decrease in growth was observed at higher concentrations. Minimal growth of 20.3% and 28.5% in dry weight was obtained at 1.5 and 1.0 field rate for A. oryzae, while the corresponding values for T. tenuis amounted only to 0% and 9.5%. (Table 1). The insecticide caused also yellowing of some Anabaena cells after growth at 1.5 field rate.

Effect of Polytrin on cell composition of carbohydrates and nutrient elements:

Polytrin had a detectable effect on the growth of A. oryzae and T. tenuis at concentrations greater than 0.25 field rate. Analysis of organisms exposed to all used concentration of insecticide (Tables 2, 3 & 4) revealed marked differences in the content of carbohydrate, macro and micro-elements as compared to untreated organisms in the exponential phase of growth.

Table 2. shows that Polytrin caused inhibition of nitrogen fixation. The nitrogen content of both strains were gradually reduced by concentration and amounted to 56.7% and 0% of the control for A. oryzae and T. tenuis respectively at 1.5 recommended field rate.

Total carbohydrate content of algal cells increased with increasing the insecticide concentrations (Table 2) reaching about twice the control value at 1.5 field rate in A.oryzae, while growth of T.tenuis was inhibited completely at that concentration.

Analysis of the intracellular macroelements of A.oryzae (Table 3) showed a slight rise in K content at low concentration and a marked reduction at higher concentrations from 0.25 recommended field rate upwards, while the marked reduction in K content with T.tenuis began from the least concentration 0.063 field rate.

Slight rise in P and Mg content was observed at low concentration followed by decrease at higher application in both strains.

Na did not show clear dependence on the insecticide concentration. Zn, Cu and Mn content showed some rise with Polytrin concentrations in both studied organisms.

Discussion

A comparison of the effect of used concentrations of Polytrin on cell growth and adaptation in A.oryzae and T.tenuis reveals that Anabaena oryzae is more tolerant to the used concentrations than Tolypothrix tenuis. Inhibition was observed at concentrations greater than

0.25 field rate. This result is similar to some previous observations on other pesticides. Wright et al, (1977) showed that 5 ppm. of the herbicide propanil prevented the growth of Anabaena cylindrica, Tolypothrix tenuis and Nostoc endophytum in flask cultures. Venkatoraman & Rajyalakshmi (1972) observed that, among 10 Anabaena strains tested for their resistance to Ceresan, 9 could tolerate 100 ppm but 1 was inhibited by concentrations higher than 1 ppm. Cylindrospermum sp. was observed to be less resistant to insecticides than aulosira fertillissima and Plectonema boryanum Singh, (1973).

Under Egyptian conditions (El-Nawawy et al, 1962; Hamdi et al, 1970; Ibrahim, 1972, and Shalan et al, 1984) showed that most Anabaena strains and Tolypothrix tenuis were inhibited by high concentrations of various pesticides not including Polytrin.

The diminution of nitrogen content of A.oryzae and T.tenuis with Polytrin indicates a decrease in nitrogenase activity of these organisms. This loss of activity could reflect a high turn over for some protein (s) involved in N₂ fixation (Romos et al, 1985). Pesticides generally appear to limit N₂-fixing capacities of blue-greenalgae, thereby affecting the overall nitrogen economy of soils (DA Silva et al, 1975; Eid et al, 1962;

El-Nawawy and Hamdi, 1979; Huang, 1978; Ibrahim, 1972 and Inger, 1970).

Total carbohydrate content of A. oryzae and T. tenuis rose with increasing Polytrin concentration in the growth media. This means that, the cells did not probably completely lose the capacity for photosynthesis in the presence of the insecticide. Hamdi et al (1970), stated that, chlorophyll synthesis was stimulated by low levels of some herbicides. Increased levels of carbohydrates were typically found in nitrogen starved cyanobacteria (Allen & Smith, 1969 and Romos, 1985). Accumulation of carbohydrates could somehow counteract the negative effect of the active nitrogenous metabolites, as it seems to be the case for the control of nitrate utilization in the unicellular cyanobacterium Anacysts (Flores et al, 1983; Espin et al, 1982; Merrick, 1982). El-Mokadem et al (1988) in a previous work have found more accumulation of carbohydrate associated with loss of nitrogen content in Anabaena oryzae grown under more saline conditions.

The elemental analysis of the dried cells revealed that, the algae exhibited changes in intracellular elements with increasing the insecticide. Lyzell et al, (1985), had stated that N:P ratio affects the growth rate of cyanobacteria.

It is recommended that for application of Polytrin insecticide in near by cotton or other crop fields, the rate would be adjusted in a manner not to harm the advantageous cyanobacteria added to rice as biofertilizer for nitrogen fixation.

~~_____~~
Acknowledgement: The authors sincerely thank Prof. Dr. M.A. Hammouda, Women's College for his encouragement during this study.

References

- Agarwal, A. (1979). Blue-green algae to fertilize Indian rice paddies. Nature 279:181.
- Allen, M.M. Smith, A.L. (1969). Nitrogen chlorosis in blue-green algae. Arch.Mikrobiol. 69:114-120.
- Alyer, R.S., S. Salahudeen, and G.S.Venkataraman (1972). Long term algalization field trial with high-yielding varieties of rice (*Oryza sativa*). Indian J.Agric. Sci. 42(5): 380-383.
- Battino-Viterbo, A., G. Minervini-Ferrante, and M. Bisiach. (1973). Comparison of activity of some chemicals on Anabaena and Chlorella-Riso 22(4): 327-336.
- Chapman, H.D. and Pratt, P.E. (1978). Methods of analysis for soils, plants and waters. University of California Div. Agric. Sci. Priced Puplication. 4034.

- DA Silva, E. J.L.E. Henricksson, and E.Henricksson (1975).
Effect of pesticides on blue-green algae and
nitrogen fixation. Arch. Environ. Contam.
Toxicol. 3: 193-204.
- Dubois, M.; Gilles, R.A., Hamilton, J.K., Robers, P.A.
and Simth, F. (1956) Colorimetric method for
determindation of sugar and related substances.
Analyt. Chem. 28: 350-356.
- Eid, M.T., M.R. Hamiss, and A.Shoukry (1962). Raddy
fertilization trials in nurseý and field Argi.
Res. Rev. (Cairo) 40: 136.
- El-Mokadem M.T., El-Shimy, A.A. ismail G.A. (1988).
Intracellular mineral content of the
cyanobacterium Anabaena oryzae in response
to saline media. Women's College for Arts,
Science and Education Annual Review (Science
section) (in press).
- El-Nawawy, A.S., M.A. El-Padl, and M.M. Nada (1962).
Economical studies on algae in Egypt. I.Effect
of new isothiouronium derivatives of
arylmercaptoalkane carboxylic acids on the
paday soil flora of algae in Egypt. J. Soc.
Sci. U.A.R. 2(1): 103-113.
- El-Nawawy, A.S. and Y.A. Hamdi (1975). Research on
blue-green algae in Egypt 1958-1972. Pages 219-
228 in W.D.P. Stewart. ed. Nitrogen fixation
by free living microorganisms, Cambridge Univ.
Press. London.
- Espin G., Alvarez-Morales A., Cannon F., Dixon R., Merrick
M. (1982). Cloning of the gln A, ntr B and
ntr C genes of Klebsiella pneumoniae and studies
of their role in regulation of the nitrogen
fixation (nif.) gene cluster. Mol. Gen. Genet.
186: 518-524.

- Flores E., Romero J.M. Guerrero M.G., Losada M. (1983).
Regulatory interaction of photosynthetic nitrate
utilization and carbon dioxide fixation in
the cyanobacterium Anacystis nidulans. Biochim.
Biophys acta. 725: 529-532.
- Hamdi, Y.A.A.S. El-Nawawy, and M.S. Tewfik. (1970).
Effect of herbicides on growth and nitrogen
fixation by alga Tolypothrix tenuis. Acta Microbiol.
Ppl. B2(19)I: 53-56.
- Huang, C.Y. (1978). Effects of nitrogen fixing activity
of blue-green algae on the yield of rice plants
(in Chinese, English summary). Bot. Bull. Acad.
Sin.19(1): 41-52.
- Ibrahim, A.N. (1972). Effect of certain herbicides on
growth of nitrogen-fixing algae and rice plants.
Symp. Biol. Hung. 11: 445-448.
- Inger, L. (1970). Effect of two herbicides on nitrogen
fixation by blue-green. Sven. Bot. Tidskr.
64: 460-461.
- Ishizawa, S., and T. Matsuguchi (1966). Effects of
pesticides and herbicides upon micro-organisms
in soil and water under water logged condition.
Bull. Nat. Inst. Agric. Sci. B. 16: 1-90.
- Jackson, D. (1967). Soil Chemical analysis, Prentice
Hall.
- Layzell, D.B., Turpin D.H., and Elrifid, I.R. (1985).
Effect of N Source on the steady State Growth
and N assimilation of P-Limited Anabaena flos-
aquae. Pl. Phsiol. 78: 739-745.

- Leigh RA. Wyn Jones (1964). A hypothesis relating critical potassium concentrations for growth to the distribution and functions of this ion . plant cell. New Phytol. 97: 1-13.
- Merrick M. (1982). A new model for nitrogen control. Nature 297: 362-363.
- Ramos, J.L., Francisco Madueno, and Miguel G.Guerrero (1985). Regulation of nitrogenase levels in Anabaena sp. ATCC. Arch. Microbiol. 141: 105-111.
- Roger, P.A. and Kulasooriya. S.A. (1980). Blue-green Algae and Rice. The International Rice Research Institute Los Banos, Laguna, Philippines.
- Shalan, S.N. M.N. Alaa El-Din; M.E. Hassan; M.S. Khadr and M. Shakr (1984). Effect of herbicides on the rice yield inoculated with blue-green algae. The second Conf. of the Agric. Res. Centre, Giza, Egypt. 9-11 April.
- Singh, P.K. (1973). Effects of pesticides on blue-green algae. Arch. Mikrob. 89: 317-326.
- Singh, P.K. (1978). Nitrogen economy of rice soils in relation to nitrogen fixation by blue-green algae and azolla. In National symposium on increasing rice yield in Kharif, Cuttack, Central Rice Research institute.
- Venkataraman, G.S., and B. Rajyalakshmi (1971). Tolerance of blue-green algae to pesticides. Curr. Sci. 40: 143-144.

- Venkataraman, G.S. and B. Rajyalakshmi (1972). Relative tolerance of blue-green algae to pesticides. Indian J. Agric. Sci. 42: 119-121.
- Watanabe, A. (1967). The blue-green algae as the nitrogen fixators. Int. Congr. for Microbiol. Moscow 66. Symp. C: 77-86.
- Watanabe, A.A. Hattori, Y. Fu Ta, and T. Kiyohara (1959). Large scale culture of blue green algae Tolypothrix tenuis. Utilizing hot spring and natural gas as heat and carbon dioxide sources. J. Gen. Appl. Microbiol. 5: 51-57.
- Wright, S.J.L., A.F. Stainthorpe, and J.D. Downs. (1977). Interactions of herbicide propanil and a metabolite 3,4. dichloroaniline, with blue-green algae Acta. Phytopathol. Acad. Sci. Hung. 12(1/2): 51:60.

Women's Coll. Ann. Rev.
Vol. 15 (1990).

Table 1: Growth of Anabaena oryzae and Tolypothrix tenuis cultures at different concentrations of Polytrin insecticide.

Polytrin Concentration	Anabaena oryzae		Tolypothrix tenuis	
	mg. dry wt 100/ml. cul.	Growth percent	mg. dry wt/100ml. culture	Growth percent
Control	158 ± 19	100%	138 ± 17	100%
0.063 R.f.r.	127 ± 14	80.4%	109 ± 8	79%
0.0125 " "	114 ± 18	72.2%	78 ± 9	56.6%
0.25 " "	88 ± 10	55.7%	57 ± 4	41.3%
0.5 " "	74 ± 9	41.5%	40 ± 7	29%
1.0 " "	45 ± 7	28.5%	13 ± 3	9.5%
1.5 " "	32 ± 5	20.3%	Dead	0

Growth percent: calculated with reference to control growth as 100%; values based on mg. dry wt./100 ml culture the values are the mean of three independent observations.

R.f.r. : Recommended field rate.

Table 2: Effect of Polytrin insecticide on the intracellular carbohydrate and nitrogen content of Anabaena oryzae and Tolypothrix tenuis. (Cultures grown with and without Polytrin untill rate logarithmic phase).

Polytrin concentration	Anabaena oryzae		Tolypothrix tenuis	
	Carbohydrates	Total nitrogen	Carbohydrates	Total nitrogen
Control	13.4	5.82	21.2	7.38
0.063 R.f.r	17.8	5.31	26.3	6.39
0.125 " "	15.3	5.3	28.3	5.42
0.25 " "	19.5	5.3	28.1	5.28
0.5 " "	22.7	4.91	30.4	5.52
1.0 " "	25.0	4.41	29.8	4.16
1.5 " "	26.3	3.35	---	---

R.f.r Recommended field rate

Table 3: Effect of Polytrin insecticide on the intracellular macro & micro elements of Anabaena oryzae.

Treatment	P mg./g .dry weight	K	Na	Mg	Fe µg./g. dry weight	Mn	Zn	Cu
Control	11.6	14.8	0.14	6.9	192.2	83.95	82.0	24.0
0.063 R.f.r	13.9	17.7	0.14	7.5	128.1	62.1	57.6	25.3
0.125 " "	12.6	16.5	0.28	7.2	138.1	73.0	57.0	20.0
0.25 " "	7.2	0.7	0.14	5.2	73.0	87.9	161.0	32.0
0.5 " "	6.6	0.8	0.14	3.8	64.4	120.5	116.2	34.7
1.0 " "	5.5	0.1	0.14	4.0	61.9	151.5	140.6	61.3
1½ " "	4.9	0.2	0.24	4.2	48.3	93.5	116.0	49.3

R.f.r : Recommended field rate.

Table 4: Effect of Polytrin insecticide on the intracellular
macro and micro elements of Tolypothrix tenuis

Treatment	P mg./g .dry wt.	K	Na	Mg	Fe	Mn	Zn	Cu
						μ g./g. dry wt.		
Control	11.1	3.5	0.21	4.5	222.	58.4	51	50
0.063 R.f.r	14.7	0.4	0.28	5.8	291.	98.1	131	62
0.125 " "	14.2	0.5	0.56	6.4	277	98.1	127	64
0.250 " "	4.7	0.5	0.56	1.9	96.1	65.4	117	50
0.500 " "	5.1	0.3	0.38	2.1	117	62.5	164	80
1.00 " "		N. D.					N. D.	
1.5 " "	--	--	--	--	--	--	--	--

R.F. r: Recommended field rate