

SPECTROSCOPIC BEHAVIOUR OF TRIETHYLENE-  
TETRAAMINEHEXAACETIC ACID WITH TETRA AND  
HEXAVALENT URANIUM IN SOLUTION

By

MOHAMED B. HAFEZ, WAFAA S. HEGAZI<sup>#</sup> and NABIL HAFEZ  
Hot Laboratory Center, Atomic Energy Establishment,  
Cairo- Egypt.

ABSTRACT

The interaction between triethylenetetraaminehexaacetic acid (TTHA) and tetra and hexavalent uranium is investigated. Evidence is given for the formation of a stable and soluble 1:1 tetravalent uranium chelate. An insoluble 2:1 chelate between uranyl ions and TTHA is formed between pH 2 and 4. Also two soluble 1:1 and 2:1 chelates are formed. Stability constants ( $\log K$ ) of the soluble complexes were calculated to be 15 and 11.8 respectively.

INTRODUCTION

Complex formation between tetra and hexavalent uranium and TTHA has been studied by some workers.<sup>1,2</sup> However, the composition and the stability of the formed complexes were not definitely determined. The probable formation of several uranyl-TTHA chelates

---

<sup>#</sup> A University College for Girls, Department of Chemistry, Ain Shams University, Cairo, Egypt.

with different compositions, in the pH range 2-8 was previously reported.<sup>3</sup> Other studies,<sup>4</sup> at low pH, showed that uranyl ions react with TTHA to form only a mononuclear chelate, whose stability varies with the pH and the concentration of the chelating agent. In view of the contradictions in the published data on the uranyl-TTHA chelates, a detailed quantitative study of the equilibria involved in the interaction of tetra and hexavalent uranium ions with TTHA, over a wide range of pH and TTHA concentrations, is undertaken in the present work.

#### EXPERIMENTAL AND CALCULATIONS

All chemicals used were Analar grade. Tetravalent uranium was freshly prepared by reducing uranyl ions with nascent hydrogen gas using orthochloroplatinic acid as a catalyst.<sup>5</sup> The chelate solutions were prepared by mixing solutions of uranyl ions and TTHA in acidic medium, then adjusting the pH of the mixtures by adding carbonate-free sodium hydroxide solution.

pH-measurements were made using a Pye Unicam Model 292 pH-meter. Spectrophotometric studies were carried out with the aid of a Beckman DU-spectrophotometer, using 1 cm quartz cells.

Formation constants of the chelates were evaluated,<sup>6</sup> and accordingly, the stability constants (log K) were then calculated.<sup>7,8</sup>

## RESULTS AND DISCUSSION

### I. Chelation of Tetravalent Uranium (U(IV)):

#### I.1. Absorption spectra of (U(IV)+TTHA) mixture as a function of pH:

The absorption spectrum of  $8 \times 10^{-2} \text{M}$  U(IV) and  $3.2 \times 10^{-1} \text{M}$  TTHA mixture at  $\text{pH} = 1$  is identical with that of free U(IV) ions, with absorption maxima at 458, 532, 618 and 645 nm<sup>4</sup>. In the pH range 2.2 - 9.5 the spectrum exhibits characteristic maxima at 464, 538, 622 and 650 nm (Fig.1). This shows that a U(IV)-TTHA complex is formed momentarily within the pH range 2.2 - 9.5. Between pH 2 and 4 the spectra of the mixture did not change for a long period after preparation. However, at  $\text{pH} \geq 4.5$  the U(IV)-TTHA complex changed gradually with time to U(VI)-TTHA.

#### I.2. Determination of the composition of the formed chelate:

##### I.2.a) The mole ratio method:

The concentration of U(IV) was kept constant at  $8 \times 10^{-2} \text{M}$  and the  $[\text{U(IV)}] / [\text{TTHA}]$  ratio was varied between 1.00 / 0.25 and 1/4. The pH of the mixture was maintained at 2.2, and the absorbance was measured at 650 nm.

##### I.2.b) The continuous variation method:

Different concentrations of U(IV) and TTHA were mixed together such that the total concentration was always  $1.6 \times 10^{-2} \text{M}$ . All the solutions were adjusted at pH 2.2 and the optical density measurements were made at 650 nm.

Both of the above methods point to the formation of 1:1 chelate, as can be seen from Figs. 2 a) and 2 b). The value of the stability constant of the complex ( $\log K$ ) was calculated to be 20.7.

## II. Chelation of Hexavalent Uranium (U(VI)):

### II.1. Absorption spectra of (U(VI)+TTHA) mixture as a function of pH:

The absorption spectra of  $8 \times 10^{-2} M$  uranyl ions in 0.2N HCl and that of (U(VI)-TTHA) mixtures at different pH values are given in Fig. 3. It was found that at pH=1.5 the spectrum of the mixture is similar to that of free uranyl ions. In the pH range 2-3 a yellow precipitate is formed indicating the formation of an insoluble compound. Between pH 3.5 and 8 a soluble complex is formed as indicated by the increase in the absorbance with increasing pH. At pH >8 the absorbance decreases with increasing pH and uranyl hydroxide is precipitated.

### II.2. Determination of the composition of the chelate:

#### II.2.a) The mole ratio method:

The absorption spectra of mixtures composed of  $8 \times 10^{-2} M$  U(VI) and various TTHA concentrations ranging from  $8 \times 10^{-3}$  to  $8 \times 10^{-1} M$  at different pH values are given in Fig.4. An insoluble

2U : 1L complex is formed at a pH range 2-3.5. A soluble 2U : 1L complex is observed at 420 nm in the pH range 4.5 -5.5. Between pH 5.5 and 8 a part of uranyl ions was precipitated indicating that the above mentioned complex does not exist at this pH range. Another soluble 1:1 complex is observed between pH 4.5 and 8. At pH = 8.5, however, the uranyl ions undergo hydrolysis yielding the hydroxide.

#### II.2.b) The continuous variation method:

The total concentration of (U(VI)-TTHA) solution mixtures was kept at  $1.6 \times 10^{-2} M$  and the pH of each solution was adjusted at 4.5. The optical measurements were recorded at 420 nm, from which the formation of two U(VI)-TTHA complexes of mole ratios 1:1 and 2:1 could be concluded. This is in accordance with the results obtained from the mole ratio method.

The values of the stability constants ( $\log K$ ) were found to be 15.0 and 11.8 for the two complexes respectively.

The competition between the oxygen of the uranyl group and the chelate ligand may be a factor which decreases the stability of the formed complex.<sup>9</sup> Similar instability of uranium complexes was also observed with other ligands such as citrates,<sup>10</sup> amines<sup>11</sup> and other polyaminopolycarboxylic acids.<sup>12</sup>

REFERENCES

1. Bhat, T.R. and Krishnamurthy, M.; J. Inorg. Nucl. Chem. 26, 587, 1964.
2. Kozlov, A.G. and Krot, N.N.; J. Inorg. Nucl. Chem. 5, 9, 954, 1960.
3. James, D.B., Poell, J.E. and Spedding, P.H.; J. Inorg. Nucl. Chem. 19, 133, 1961.
4. Hafez, M.B.; "Thèse de Doctorat es science" CEA-R-3521, 1968.
5. Gogliat, G., De Leone, R. and Lanz, R.; RT/CHI(64), 14 Roma Giugno, 1964.
6. Vosburg and Cooper; J. Am. Chem. Soc. 63, 437 (1941); Sherif and Awad; J. Inorg. Nucl. Chem. 24, 197 (1962).
7. Gel'man, A.D. and Maslov'eva, M.P.; Dok. Akad. Nauk, 124, 815, 1959.
8. Grimes, J.N., Huggard, M.J. and Wilford, S.P.; J. Inorg. Nucl. Chem. 25, 1225, 1963.
9. Hafez, M.B. and Mahmoud, K.A.; Isotope and Rad. Res., 6, 2, 1973.
10. Dyatkina, M.E., Markov, V.P., Tsepina, I.V. and Hikhalliev, Yu.M.; Russ. J. Inorg. Chem. 6, 5, 293, 1961.
11. Sacconi, L.; Atti. Accad. Nazl. Lincei. Classe. Sci. Fis. Nat.; 8, 6, 639, 1949.
12. Carell, M.J.; Analyst, 77, 859, 1952.

## المتراكبات الناتجة من تراكب

ثلاثي ايثيلين رباعي امين سداسي حمض الخليك مع كل من

اليورانيم الرباعي واليورانيوم السداسي التكافؤ

محمد بدر الدين حافظ - وفاء صلاح حجازي\* - نبيل حافظ

مؤسسة الطاقة الذرية

تم في هذا البحث دراسة التراكب بين مشتق الحمض الاميني : ثلاثي ايثيلين رباعي امين سداسي حمض الخليك وبين كل من اليورانيم رباعي التكافؤ واليورانيوم سداسي التكافؤ لاحتماليه تكوين مركبات متراكبة . وقد دلت النتائج على تكسب متراكبات ثابت وقابل للذوبان في الماء بنسبة تركيبية 1 : 1 بين اليورانيم الرباعي وبين مشتق الحمض الاميني . كذلك اثبتت النتائج تكوين متراكب غير قابل للذوبان بين اليورانيل وبين مشتق الحمض الاميني بنسبة تركيبية 2 : 1 في المحاليل ذات الأس الهيدروجيني اقل من 2 و اقل من ( اوي ساوي ) . ٤

ايضا تم الاستدلال على تكوين متراكبين قابلين للذوبان في المحلول المائي بنسبة تركيبية 1 : 1 و 1 : 1 بين ثقب اليورانيل ومشتق الحمض الاميني على التوالي . وفي جميع الاحوال تم حساب ثوابت الاتزان للمتراكبات القابلة للذوبان المذكورة اعلاه .

\* كلية البنات - قسم الكيمياء - جامعه عين شمس

1. Effect of pH on the absorption spectra of (U(IV)+TTHA) mixture;  $[U(IV)] = 8 \times 10^{-2} M$ ,  $[TTHA] = 3.2 \times 10^{-1} M$ ;  
1) pH = 2.2, 2) pH = 5.5, 3) pH = 7.5, 4) pH = 9.5 and 5) U(IV) only.
  
- 2.a) Variation of the absorbance at 650 nm with the mole ratio  $[U(IV)] / [TTHA]$  at pH 2.2,  $[U(IV)] = 8 \times 10^{-2} M$ ,
- 2.b) Variation of the absorbance at 650 nm for (U(IV)-TTHA) mixtures with mole fraction at pH 2.2, total concentration of the mixtures  $1.6 \times 10^{-2} M$ .
  
3. Effect of pH on the absorption spectra of (U(VI)+TTHA) mixtures;  $[U(VI)] = 8 \times 10^{-2} M$ ,  $[TTHA] = 3.2 \times 10^{-1} M$ .  
1) pH = 3.5, 2) pH = 4.5, 3) pH = 5.5, 4) pH = 6.0,  
5) pH = 6.5, 6) pH = 7.0, 7) pH = 7.5, 8) pH = 8.0,  
9) pH = 9.5 and 10) U(VI) only in 0.2N HCl.
  
4. Variation of the absorbance at 420 nm with the molar ratio of hexavalent uranium to TTHA;  $[U(VI)] = 8 \times 10^{-2} M$ ,  
A) pH = 3.5, B) pH = 4.5, C) pH = 5.5 and D) pH = 7.5.






Fig. 1  $\lambda$  nm



$$\alpha = \frac{[UUE]}{[UUE] + [THA]}$$

Fig. 26




Fig. 3

