

الفصل الثانى عشر

الأمراض المتسببة عن النباتات العشبية المتطفلة

Diseases Caused by Herbaceous Parasitic Plants on Tomato

سبب النباتات المتطفلة على المحاصيل أضرارًا كبيرة لعوائلها النباتية، ويوجد أكثر من ١٠٠٠ نوع من النباتات الرعمية تتبع ٧ عائلات تتطفل على نباتات زهرية أخرى مهمة اقتصاديًا. تنتج النباتات المتطفلة أزهارًا وبذورًا. ومن أهم النباتات المتطفلة على نبات الطماطم الهالوك (*Orobancha spp*)، والحامول (*Cuscuta spp*) Dodder. ويتشربان فى أقطار غرب آسيا وشمال إفريقيا وجنوب أوروبا وأيضاً فى أمريكا الشمالية. تقوم هذه النباتات بسحب الماء والعادن والمركبات العضوية اللازمة لتغذيتها من النبات العائل مما يؤدي إلى ضعف نمو العائل نتيجة لتأثر العمليات الفسيولوجية المختلفة به.

تتميز نباتات الهالوك والحامول بخلوها من مادة الكلوروفيل ومن ثم لا يستطيعان تكوين غذائهما العضوى دون الاعتماد فى ذلك على النبات العائل. ومع ذلك توجد نباتات متطفلة أخرى لها أوراق خضراء أى تحتوى على كلـيـل لكن لا يوجد لها جذور، وبذلك تخلق المادة العضوية اللازمة لها عن طريق إتمام عملية التمثيل الضوئى وتعصل على الماء والمواد المعدنية اللازمة لها من العائل. ومن الأمثلة على ذلك نبات العدار *Striga hermonthica* ونوع الدبق *Viscum sp*. وبعض أنواع الأوركيد.

تختلف طريقة وطبيعة التطفل تبعاً للنبات المتطفل. الهالوك مثلاً يتطفل تحت سطح التربة ويصيب الأجزاء الأرضية من العائل. أما الحامول فيتطفل من فوق سطح التربة ويصيب السوق أو الفروع فى العائل. وقد يكون التطفل خـلـى حيث يوجد معظم جسم الطفيل خارج العائل، وجزء صغير منه فقط داخل العائل. ويسمى النبات المتطفل فى هذه الحالة نبات ناقص التطفل مثل العدار والدبق، ويوجد أيضاً تطفل داخلى أى أن كل جسم الطفيل يوجد داخل العائل ولا يظهر منه إلا أجزاءه الزهرية فقط ويوجد ذلك فى بعض النباتات كاملة التطفل مثل الرافليزيا.

١- الهالوك Broomrape

لهالوك نبات زهرى متطفل إجبارياً يتبع عائلة Orobanchaceae ويوجد أساساً فى منطقة البحر المتوسط حيث توجد مساحات كبيرة مصابة به. ويوجد أيضاً فى مناطق أخرى ظروفها تشبه ظروف منطقة البحر المتوسط مثل كاليفورنيا بالولايات المتحدة الأمريكية وفى غرب استراليا وكوبا.

يتبع جنس الهالوك *Orobancha spp* حوالى ١٢٠ نوعاً تصيب عديد من العوائل فى جميع أنحاء العالم منها: البطاطس- الطماطم- البقوليات- الكرنب- البطيخ- الدخان ويصيب أيضاً عديد من الحشائش. ويسبب الهالوك أضراراً كبيرة حيث تتطفل بعض أنواعه على مدى واسع من العائلات النباتية منها - Solanaceae - Apiaceae - Cucurbitaceae - Aesteraceae - Fabaceae.

فى مصر يوجد بعض الأنواع من الهالوك تتطفل على الطماطم وقد تتطفل على محاصيل أخرى. ومن هذه الأنواع:



- O. crenata* ويتطفل على الطماطم والبقوليات وعديد من النبات الأخرى.
O. ramosa ويتطفل على الطماطم وهي أهم عائل له.
O. aegyptiaca ويصيب الطماطم والبطاطس أساسًا وتوجد محاصيل أخرى عائلة له.
O. cernua ويتطفل على الطماطم وعباد الشمس.

الشكل الظاهري للهالوك Broomrape morphology

الهالوك نبات زهري كامل التطفل على جذور العائل. يتكون الهالوك من شمراخ زهري حولي قد يكون مدعًا أحيانًا. الأوراق مختزلة خالية من الكلوروفيل وتظهر على هيئة حراشيف بنية اللون. الأزهار خنثى وحيدة العنبر. الثمار علبة تنفتح مصراعيا وتحتوى على مئات من البذور الصغيرة الحجم ذات شكل كمثرى ولون بنى إلى أسود. بذور الهالوك يمكن أن تبقى كامنة فى التربة لعدة سنوات إلى أن تتشجع على الإنبات ببعض مركبات معينة تنتج بواسطة جذور النباتات الحية القريبة منها (Fig 1).

الوضع التقسييمى للهالوك Classification of Broomrape

Kingdom: Plantae

Phylum: Magnoliophyta

Class: Magnoliopsida

Order: Lamiales

Family: Orobanchaceae

Genus: *Orobanche*

Species. *O. aegyptiaca* Pers- *O. crenata* Forsk- *O. cernua* Loefl- *O. ramosa*

أعراض الإصابة بالهالوك Symptoms of broomrape infection

النباتات المصابة بالهالوك ضعيفة ومتقزمة إذا ما قورنت بالنباتات السليمة وتصفّر أوراقها وتظهر نموات الهالوك لسفراء بالقرب من قاعدة نبات الطماطم وعند إزالة التربة من حول النبات يمكن مشاهدة منطقة الاتصال بين الهالوك وجذور الطماطم. ثم بعد فترة قصيرة من نمو نبات الهالوك تنمو الأزهار البيضاء أو الملونة. بعد ذلك تجف نباتات الهالوك ويصبح بنية اللون وتنتثر ثمارها وتنطق منها البذور وتختلط بالتربة بعد سقوطها عليها أو تحمل بواسطة الرياح إلى مسافات بعيدة.

تطفل الهالوك Broomrape pathogenicity

يمكن أن تبقى بذور الهالوك ساكنة فى التربة فى غياب العائل أكثر من ١٠ سنوات. وعند وجود العائل وأى محصول منبه لإنبات بذور الهالوك وتوفر الظروف الملائمة تبدأ البذرة فى الإنبات مكونة أنبوبة إنبات نحو فى اتجاه العائل القريب منها جدًا وتلتصق بجذوره وترسل ممصات إلى داخل الجذر وتعمق فى الداخل إلى أن تص إلى الاسطوانة الوعائية وتقوم بامتصاص العناصر الغذائية اللازمة لها وأيضًا كمية الماء الضرورية لحياتها. وتكود عسما منتفخا نتيجة لامتصاص الغذاء ثم تخرج من هذا الجسم ممصات أخرى تتصل أيضًا بجذور العائل. هذا الجسم ينتفخ يستطيل مكونا شمراخا زهريا - قد يتكون أكثر من شمراخ وتظهر هذه الشمراخ فوق سطح التربة وتنتفخ أزهارها ويتم الاخصاب وتتكون الثمار والبذور وتنضج البذور سريعًا.



يصل عدد البذور الناتجة من نبات واحد من الهالوك إلى أكثر من ٢٠٠ ألف بذرة.

المقاومة Control

هم طرق المقاومة العمل على تخفيض عدد بذور الهالوك في التربة إلى أقل مستوى يمكن الوصول إليه مع الإبقاء عن هذا المستوى للسنوات التالية ويتم ذلك بالآتي :

١ - إزالة الشماريخ الزهرية للهالوك باليد قبل نضجها وإعدامها.
٢ - استعمال المحصول الصائد. وقد استخدم الكتان في مصر كنبات صائد للهالوك حيث تفرز جذوره مواد تنبه إنذار بذور الهالوك.

٣ - تأخير ميعاد زراعة الطماطم.

٤ - زراعة أصناف مقاومة للهالوك.

٥ - تحفيز بذور الهالوك على الإنبات في غياب العائل وبالتالي تموت بادرات الهالوك النابتة قبل زراعة المحصول. وتعرف هذه الطريقة بالإنبات الانتحاري Suicidal germination وتتم باستعمال مركبات خاصة منها مادة Strigol المستخلصة من بذور القطن. وهي ذات فاعلية كبيرة في تحفيز إنبات بذور الهالوك.

المقاومة البيولوجية Biological control

توجد فطريات متطفلة على الهالوك منها *Sclerotium orobanche* و *Fusarium orobanche* وتوجد أيضاً حشرات متطفلة مثل *Phytozoma orobanchia* و *Tropinota squalidis* كذلك جنس *Agrotis sp*.

المقاومة الكيماوية Chemical control

تمسى التربة الملوثة وقبل زراعة البذرة أو الشتلة يستعمل مركب Metam sodium أو أحد المدخنات التجارية الأخرى التي تتبع مجموعة metam sodium وهو مصدر للمركبات العضوية الطيارة Volatile organic compounds - لذلك لا يلجأ إلى المدخنات إلا إذا كانت باقى استراتيجيات المقاومة غير متاحة أو غير ناجحة.

تحلّك يستعمل مبيد الحشائش Sencor DF (metribuzin) قبل زراعة الطماطم بأربعة أشهر حتى لا تضر النباتات. وتحتف جرعة المبيد باختلاف نوع التربة. فى التربة المحتوية على مادة عضوية أقل من ٢٪ تستعمل الجرعة المنخفضة أما جرعة المرتفعة فتستعمل فى التربة التى بها أكثر من ٢٪ مادة عضوية ولا يستعمل فى التربة عالية القلوية أو الرملية الخفيفة.

أما بعد زراعة البذرة أو الشتلات يستعمل مبيد الحشائش Matrix 25DF (nimsulfuron) رشاً على التربة. وإذا كانت هذه الحشائش قد ظهرت فوق سطح التربة يضاف إلى المبيد أحد الناشرات السطحية غير أيونية nonionic surfactant بمعد ٢٥،٠٪ (٧/٧).

كذلك يستعمل مبيد الحشائش Sandea (halosulfuron)، لكن فى حالة معاملة التربة بمبيدات حشرية تتبع Organophosphate insecticides لا يستعمل مبيد الحشائش Sandea ولا يستعمل أى مبيد حشرى يتبع Organophosphate insecticides رشاً على المجموع الخضرى للنبات إلا بعد ٧ أيام من استعمال halosulfuron أو قبل ٢١ يوماً من استعماله.



٢- الحامول Dodder

يتبع الحامول جنس *Cuscuta* الذى ينتشر فى جميع مناطق العالم المعتدلة والحارة ونادرا ما يوجد فى المناطق الباردة ويوجد منه أكثر من ١٠٠ نوع نباتى متطفل تختلف فى ألوانها ما بين الأصفر والبرتقالى والأحمر لكن نادرا ما يوجد اللون الأخضر. يتطفل الحامول على عديد من النباتات تشمل محاصيل حقلية ومحاصيل بستانية منها البرسيم الحجازى- الكتان- البرسيم- البطاطس- الطماطم- الداليا- الكريز انثم- البيتونيا- الهيلينيوم وأنواع أخرى من النباتات المختلفة.

قديمًا كان يطلق على الحامول أسماء عديدة منها: شعر الشيطان Devil's hair - شعر الساحرة Witch's hair - أحشاء أبليس Devil's gut - الخيوط الذهبية Gold thread وغير ذلك من الأسماء.

الشكل الظاهرى للحامول Dodder morphology

الحامول نبات ذات ساق رفيعة لا تحمل أوراق لكن يوجد بها ندب دقيقة مسكان الأوراق المختزلة ويوجد بها نسبة قليلة جداً من الكلوروفيل. لكن بعض أنواع الحامول يمكن أن تقوم بعملية التمثيل الضوئى بدرجة بسيطة جداً مثل *C. reflexa*. بينما أنواع أخرى تعتمد فى تغذيتها اعتماداً كاملاً على النبات العائل مثل *C. europaea*. ويختلف نمو وطول الحامول باختلاف مناطق نموه. فى المناطق الحارة قد يصل ارتفاعه إلى ارتفاع عروش الشجيرات أو يحول الأشجار. أما فى المناطق المعتدلة ينحصر طوله فى النباتات الحولية ذات الطول الخضرى القليل نسبياً. ويمكن أن يتلاصق الحامول مع نبات واحد أو عدة نباتات فى وقت واحد. وتتدرج ألوان أزهار الحامول من الأبيض إلى الوردى الفاتح أو الأصفر أو الكرىمى. هذه الأزهار توجد فى أوائل الصيف فى بعض أنواع الحامول وفى أنواع أخرى من الحامول توجد فى آخر الصيف.

بذور الحامول دقيقة جداً وتنتج بكميات وفيرة ولها غطاء بذرى صلب ويمكن أن تبقى فى التربة من ٥ - ١٠ سنوات أو أكثر.

الوضع التصنيفى للهامولك Classification of Broomrape

Kingdom: Plantae

Phylum: Magnoliophyta

Class: Magnoliopsida

Order: Solanales

Family: Convolvulaceae

Genus: *Cuscuta*

Species: *C. campestris* and *C. pentagona*

صنف جنس *Cuscuta* سابقاً تبع عائلة Cuscutaceae وكان هو الجنس الوحيد بهذه العائلة. والآن تم وضعه فى عائلة Convolvulaceae.



أعراض الإصابة بالحامل Symptoms of dodder infection

يظهر الطفيل فوق سطح التربة بشكل خيط ملتف حول سوق النبات العائل مرصلاً ممصاته داخل أنسجة الساق لتعض الأنسجة الوعائية للطفيل بالأنسجة الوعائية للعائل. يعقب ذلك ذبول ساق الحامل في مكان اتصالها بالتربة ثم يعيش الطفيل بعد ذلك كلية على حساب النبات العائل ويمتص منه الماء والمواد الغذائية المجهزة ويكون الطفيل سيقان عديدة تنمو بسرعة وتلتف بدورها حول العائل ثم يزهر ويكون ثمار (Fig 2) بعد نضج الثمار تنفتح وتتساقط من البذور. تظهر الإصابة في أماكن متفرقة من الحقل وتزداد مساحتها أثناء نمو نباتات الطماطم. ونتيجة الإصابة بالحامل يضعف نبات الطماطم وتصغر أوراقه وتقل مقاومته للأمراض الفيروسية.

تطفل الحامل Dodder pathogenicity

تساقط بذور الحامل بعد نضجها في التربة وعند وجود رطوبة أرضية كافية تنبت البذور - يمكن أن تحتفظ هذه البذور بحيويتها من ١٠ إلى ٢٠ سنة- وتعطي أنابيب نبات صفراء خيطية تظل أطرافها السفلى في التربة بينما تظهِر الأطراف الأخرى فوق سطح التربة. وتتحرك إلى أن تصل إلى ساق النبات العائل. (إذا لم تقابل أنابيب نبات بذور الحامل عائلاً مناسباً فإنها تموت). ثم ترسل ممصاتها داخل الساق للتغذية. يعقب ذلك ذبول ساق الحامل في مكان اتصالها بالتربة ويبقى كامل التطفل على العائل فوق سطح التربة.

المقاومة Control

- ١ - في الحقل الموبوء بالحامل يمنع زراعة محاصيل عائلة له لعدة سنوات.
- ٢ - منع استيراد أو زراعة بذور ملوثة ببذور الحامل.
- ٣ - الدورة الزراعية غير مؤثرة عامة في المقاومة وذلك لكثرة عوائله وأيضاً لأن حيوية بذوره يمكن أن تستمر فترة طويلة ومع ذلك عمل دورة زراعية مع محاصيل غير عائلة مثل القطن والأذرة والقمح والشعير والثوم يمكن أن تقلل تجمع بذور الحامل.
- ٤ - معظم إنبات بذور الحامل يتم في الربيع (من ١ مارس إلى منتصف مايو) لذلك تأخير الزراعة أو الشتل يمكن أن يساعد على تقليل المشكلة مع الحامل.
- ٥ - الطريقة المثلى لمقاومة الحامل هي القضاء على نبات الطماطم العائل مباشرة عند ملاحظة إصابته بالطفيل.
- ٦ - إذا لوحظ الحامل بعد تكوينه للأزهار يزال هو والنبات العائل من الحقل ويحرق لقتل البذور.
- ٧ - بعد العمل في الحقل المصاب بالحامل يجب التخلص من الملابس والأدوات العالق بها بذور الحامل قبل الدخول إلى حقول جديدة خالية من الإصابة.
- ٨ - توجد طماطم مقاومة للإصابة بالحامل وتمنع إنتاج بذوره منها PX 665 - CDS 233 - Heinz 9492.

المقاومة الحيوية Biological control

تتص حشرة *Melanagromyza cascutea* على الحامل ويمكن استخدامها في المقاومة الحيوية له.



المقاومة الكيماوية Chemical control

المبيد الكيماوى Prowl H20 (pendimethalin) ويستعمل قبل زراعة الشتلات فقط . يمكن أن يقلل إنبات ويزع الحامول بنسبة ٨٠٪.
استعمال المبيد Matrix 25 DF (rimsulfuron) عندما تكون نباتات الطماطم فى طور البادرة يقلل نمو الحامول.
يستعمل مبيد الحشائش Dacthal (DCPA) فى مقاومة الحامول أثناء الربيع عندما تكون الطماطم المنزرعة بالبحر طولها من ٤ - ٦ بوصات والطماطم التى تم شتلها بعد ٤ - ٦ أسابيع من الشتل.

مشاكل الحشائش الخاصة بنباتات الطماطم Tomato Weed Problems

الحشائش مصدر من أخطر المصادر فى إصابة حقول الطماطم بالحشرات ومسببات الأمراض المختلفة حيث تعول وتأوى هذه الآفات . كذلك تنافس الحشائش نباتات الطماطم الموجودة فى نفس الحقل على الغذاء والماء وكسوة مما يؤدي إلى ضعف نباتات الطماطم وقلة المحصول . ومن الحشائش التى تسبب مشاكل فى حقول الطماطم : Field bindweed وهو حشيشة معمرة عميقة الجذور فى التربة ومن الصعب مقاومتها عندما تثبت هذه الجنفر . حشائش Nightshade family : ويتبعها مجموعة من الحشائش المختلفة مثل Black nightshade و Heart nightshade و Cutleaf nightshade و Groundcherry وغيرها من الحشائش . هذه الحشائش حولية وتنمو فى طبقة السطحية للتربة على عمق من ١ - ٢ بوصة (٢.٥ - ٥ سم) .
Nutsedge : وهى حشيشة معمرة يمكن أن تكون بذور ودرنات . يمكن للدرنات أن تبقى حيه فى التربة لعدة سنوات إلى أن تتحسن الظروف الملائمة لنموها . الدرنة الواحدة بها ٤ - ٧ براعم وكل برعم له القدرة على تكوين نبات . عادة ينمو برعم واحد فقط من الدرنة . لكن إذا أضر هذا البرعم وانتهى إما عن طريق العمليات الزراعية أو مبيدات الحشائش ينمو نبات جديد من برعم آخر موجود على نفس الدرنة .
ونتيجة للضرر الذى تحدثه الحشائش فى حقول الطماطم يجب التخلص منها ويتم ذلك بالآتى :

أولاً: العمليات الزراعية Practical cultures

١ - التشميس قبل الزراعة : فى المناطق المعتدلة يمكن أن ترتفع درجة حرارة التربة المغطاة بشرايح البولي إيثيلين الشفاف حوالى ٦ - ١٣ م (٤٣ - ٥٥,٥ ف) عن التربة غير المغطاة وقد أدى هذا إلى تقليل كثافة تجمعات الفطر والبكتيريا من ٦٢ - ١٠٠٪ إذا تمت المعاملة قبل الزراعة . وقللت أيضا إنبات وظهور الحشائش بنسبة ٩٧ - ١٠٪ (Stapleton, 1986) . هذه العملية يمكن أن تتم فى مصر فى شهور الصيف الحارة (يولية وأغسطس) بعد أن يُم رى الأرض ربا غزيراً ثم تغطى بشرايح البلاستيك الشفاف لمدة ٦ - ٨ أسابيع .
٢ - العزيق : يتم العزيق اليدوى ٣ مرات : الأولى بعد الشتل بحوالى ٢ - ٣ أسابيع لإزالة الحشائش الصغيرة . والثانية بعد ٢ - ٣ أسابيع من الأولى وتتم الثالثة بعد ٢ - ٣ أسابيع أخرى من الثانية وبذلك يتم التخلص من الحشائش وتغطية الأسمدة المضافة التى توضع فى باطن الخط .
وبالعزيق تنتقل التربة من الريشة البطالة إلى الريشة العمالة والترديم على النباتات وتعديل وضعها .



٣ - الحرث العميق: كثير من أنواع الحشائش الحولية توجد فى الطبقة السطحية للتربة أى على عمق ٣-١ بوصة (٥،٢- ٥ سم) كذلك بعض الحشائش المعمرة توجد درناتها على عمق ٤-٦ بوصات (١٠- ١٥ سم). فإذا تم عرث الأرض بمحراث قلاب فإنه يتم دفن بذور أو درنات هذه الحشائش إلى عمق حوالى ١٢ بوصة (٣٠ سم) على الأقل وهذا العمق يؤدي إلى دفنها وتقليل ظهورها إلى أقصى درجة.

٤ - تغطية خطوط الزراعة بالبلاستيك الأسود: يتم ذلك فى العروة الشتوية أو الصيفية المبكرة مع وجود ثقب على مسافات منتظمة لزراعة الشتلات. وتتبع هذه الطريقة أيضاً فى الطماطم التى تنمو تحت الأنفاق البلاستيكية المنخفضة. هذه الطريقة أدت إلى انخفاض نمو الحشائش وتدفئة التربة حول الجذور وزيادة مقدرة الجذور على امتصاص الماء والمواد الغذائية مما أدى إلى التبرير فى الإنتاج. مع مراعاة إزالة هذه الأغشية فى جميع الحالات عندما يصل حجم النمو الخضرى فى نباتات الطماطم إلى أكثر من ٦٠٪.

ثاني: المقاومة الكيماوية Chemical control

١- فى الزراعة وقبل بزوغ الحشائش يمكن استعمال أى مبيد تجارى من المركب metam sodium كتدخين للتربة أو مبيد (Dual Magnum) (metolachlor).

٢- فى الزراعة وقبل بزوغ الحشائش يمكن استعمال (Select Max) (clethodim) ويستعمل فى الحشائش الحولية وإذا كانت الحشائش معمرة يجب تكرار المعاملة.

٣- كذلك يستعمل مبيد (Sanda) (halosulfuron) فى مقاومة الحشائش المعمرة فى الحقول المزروعة طماطم. وبيد Sencor 70% WP (ويتبع مجموعة (metribuzin) أو (Matrix 25DF) (rimsulfuron) ويستعملان فى مقاومة الحشائش الحولية عريضة وضيقة الأوراق.

أضرار استعمال مبيدات الحشائش Herbicides Injury

١- أكثر مبيدات الحشائش ضرراً على حقول الطماطم مبيد الحشائش 2,4-D (2,4-Dichloro phenoxy acetic acid) حينما تتعرض هذه الحقول للأبخرة الناتجة عن استعماله فى الحقول المجاورة لها أو الرذاذ أيضاً استعمال الرشاشات التى تم استعمالها فى رش المبيد فى هذه الحقول وأعيد استعمالها فى رش مبيدات أخرى فى حقل الطماطم. وتبدأ الأعراض بانحناء وريقات النموات الطرفية وصغر حجم الأوراق والتفافها وتأخذ العروق اللون الأخضر الباهت وتتكون ظاهرة الجيوب Puffiness فى الثمار وأيضاً مرض Cat face وتخلو الثمار من البذور وتتشوه. ولجناب أضرار استعمال هذا المبيد:

- ١ - عدم استعمال المبيد فى الحقول المجاورة لحقل الطماطم عند اشتداد الرياح.
 - ٢ - عدم استعمال المبيد فى حقل ذات مستوى أعلى من حقل الطماطم.
 - ٣ - عند الرش بالمبيد فى الحقول المجاورة لحقل الطماطم تستعمل الرشاشات ذات الضغط المنخفض.
 - ٤ - عدم استعمال رشاشات هذا المبيد فى رش مبيدات أو أى مركبات فى حقل الطماطم.
- ألمبيد الحشائش Glyphosate عندما تتعرض نباتات الطماطم له أثناء التزهير أو ما قبل التزهير مباشرة يضر ذلك بسجموع الخضرى ضرر من متوسط إلى ضرر شديد وقد يؤدي إلى تساقط الأزهار ويقل عقد الثمار.





Broomrape



Fig (1)
Orobanche sp

Dodder




Fig (2)
Cuscuta sp



Herbicide Damage

Fig (3)
Herbicide damage on tomato plants



المراجع العربية

- ٦ - العروسى، حسين وعماد الدين وصفى (١٩٧٤). الملكة النباتية- دار المطبوعات الجديدة- الإسكندرية ٣١٤ صفحة.
- ٧ - العروسى، حسين وسمير ميخائيل ومحمد على عبد الرحيم (١٩٩٢). أمراض النبات. منشأة المعارف بالإسكندرية- ٥٠٢ صفحة.
- ٣ - المبيدات الزراعية فى مصر (٢٠٠٧).
- (<http://www.apc.gov.eg/ar/products/showall.aspx>) 1 - 34p.
- ٤ - ثابت، كمال على ومحمود ماهر رجب وعبد الله أحمد الشهيدى ومصطفى محمد فهم (١٩٧٦). علم أمراض النبات. مطبعة السعادة القاهرة- ٦٩٣ صفحة.
- ٥ - حسن، أحمد عبد المنعم (١٩٩١). الطماطم. الدار العربية للنشر والتوزيع- ٣٣١ صفحة.
- حماد، شاکر محمد وأحمد أبو النجا ومحمد عبد الفتاح وليبيب شنب (غير معروف سنة النشر) أساسيات علم الحشرات- دار المطبوعات الجديدة- الإسكندرية- ٢٩٥ صفحة.
- ٦ - زكى، سعد على وعصمت خالد علام (١٩٦٩). أمراض النبات البكتيرية والفيروسية- مكتبة الأنجلو المصرية. القاهرة- ٤٩٠ صفحة.
- ٣ - وزارة الزراعة واستصلاح الأراضى- مركز البحوث الزراعية- جمهورية مصر العربية (٢٠٠٣)- زراعة وإنتاج الطماطم. نشرة رقم ٨١٦- ٨٤ صفحة.



المراجع الأجنبية

- 1 - Abang, M.M.; B. Bayaa; and B. Abu Irmaileh (2007). A participatory farming system approach for sustainable broomrape (*Orobanche spp.*) management in the Near East and North Africa. *Crop Prot.* **26** (12): 1723 - 1732.
- 2 - Abawi, G.S.; and R.G. Gorgan (1979). Epidomology of diseases caused by *Sclerotinia* species. *Phytopathology.* **69**: 899 - 904.
- 3 - Abd El Halcem, Soad T. (1998). Biocontrol of cucumber grey mould by *Trichoderma harizianum* and *Bacillus megaterium*. *El Minia Sci. Bull.* **11** (2): 41 - 53.
- 4 - Abd El Halcem, Soad T. (2001). Biological, chemical and biochemical control of cucumber powdery mildew. *El Minia Sci. Bull.* **13** (2) & **14** (1) pp. 229 - 242.
- 5 - Abiko, K. (1978). Influence of temperature and humidity on development of tomato powdery mildew. *Proc. Kansai Pl. Prot. Soc.* **20**: 49 - 52.
- 6 - Abou - Elfotoh, Mohga A. (1991). Studies on exocortis disease of citrus. Msc thesis Fac. Agric. Cairo Univ. 59p.
- 7 - Abouzid, A. M.; J.E. Polston; and E. Hiebert (1992). The nucleotide sequence of tomato mottle virus, a new geminivirus isolated from tomatoes in Florida. *J. Gen. Virol.* **73**: 3225 - 3228.
- 8 - Abu - Gharbich, W.L.; and V. G. Perry (1970). Host differences among populations of *Belonolaimus longicaudatus* Rau. *J. Nematol.* **2**: 209- 216.
- 9 - Abu- Gharbich, W.L.; K.M. Makkouk; and A.R. Saghir (1978). Response of different tomato cultivars to the root- knot nematode, tomato yellow leaf curl virus, and *Orobanche* in Jordan. *Plant Dis. Rep.* **62**: 263 - 266.
- 10 - Accota, G.P.; J. Navas - Castille; E. Noris; E. Moriones; and D. Louro (2000). Typing of tomato yellow leaf curl viruses in Europe. *European J. Pl. Pathol.* **106**: 179 - 186.
- 11 - Adams, P. (1986). Mineral nutrition. In J. G. Atherton and J. Rudich (Eds) «The Tomato crop» pp 281 - 334. Chapman and Hall. London.
- 12 - Akhtar, K. P.; M. Dickenson; G. Sarwar; F.F. Jamil; and M.A. Haq (2008a). First report of the association of a 16 SrII phytoplasma with sesame phyllody in Pakistan. *Pl. Pathol.* **57**: 77 .
- 13 - Almeida, A.M.R.; R.V. Abdelnoor; C.A. Arrabal - Arias; V.P. Calvalho; S.S. Jacond - Filho; S.R.R. Marin; L.C. Benato; M.C. Pinto; and C.G.P. Carvalho (2003). Genotypic diversity among Brazilian isolates of *Macrophomina phaseolina* revealed by RAPD. *Fitopatologia Brasileira.* **28**: 279 - 285.
- 14 - Amadioha, A.C.; and P.N. Uchendu (2003). Post- harvest control of tomato fruit rot caused by *Fusarium solani* with extracts of *Azadirachta indica*. *Discovery and Innovation,* **15** (6): 83 - 86.
- 15 - Amin, A.W.; and M.M.A. Youssef (1998). Effect of organic amendments on the parasitism of *Meloidogyne javanica* and *Rotylenchulus reniformis* and growth of sunflower. *Pakistan J. Nematol.* **16** (1): 63 - 70.
- 16 - Anderson, B.; M. Sandstrom; and A. Stromberg (1998). Indications of soilborne inoculum of *Phytophthora infestans*. *Bot. Res.* **41**: 305 - 310.

- 17 - Andrivon, D. (1995). Biology, ecology and epidemiology of the potato late blight pathogen *Phytophthora infestans* in soil. *Phytopathology*, **85**: 1053 - 1056.
- 18 - Antignus, Y.; O. Lachman; M. Pearlsman; R. Gofman; and M. Bar - Joseph (2002). A new disease of greenhouse tomatoes in Israel caused by distinct strain of tomato apical stunt viroid (TASVd). *Phytoparasitica*, **30** (5): 502 - 510.
- 19 - Antignus, Y.; M. Pearlsman; O. Lachman; and F. Feigelson (2006). Tomato apical stunt viroid (TASVd) a pathogen of greenhouse tomato seedborne and transmitted by bumble bees. *Phytoparasitica*, **34** (3): 306 - 307.
- 20 - Antignus, Y.; O. Lachman; and P. Malenia (2007). Spread of tomato apical stunt viroid (TASVd) in greenhouse tomato crops is associated with seed transmission and bumble bee activity. *Plant Dis.* **91** (1): 47 - 50.
- 21 - Anzai, Y.; H. Kim; J. Y. Park; and H. Wakabayashi (2000). Phylogenetic affiliation of the Pseudomonads based on 16S rRNA sequence. *Int. J. Syst. Evol. Microbiol.* **50**: 1563 - 1589.
- 22 - Apt, W.J.; and E.P. Caswell (1988). Application of nematocides via drip irrigation. *Ann. Appl. Nematol.* **2**: 1 - 10.
- 23 - Arteaga, M.L.; R.C. Emilio; A. Fraile; S. Elisa; and R.C. Fernando (1996). Different tomato bushy stunt virus strains that cause disease outbreaks in solanaceous crops in Spain. *Phytopathology*, **86**: 535 - 542.
- 24 - Asaka, O.; and M. Shoda (1996). Biocontrol of *Rhizoctonia solani* dampingoff of tomato with *Bacillus subtilis* RB - 14. *App. Environ. Microbiol.* **62** (1): 4081 - 4085.
- 25 - Auerswald, H. (1978). Effect of climatic conditions on fruit set of glasshouse tomatoes. (*In De*) *Gartenbau*, **25**: 297 - 299.
- 26 - Avgellis, A.D.; N. Roditakis; C.I. Dovas; N.I. Katis; C. Varvert; N. Vassilakos; and F. Bem (2001). First report of tomato yellow leaf curl virus on tomato crops in Greece. *Plant Dis.* **85**: 678.
- 27 - Babarola, J.O. (1982). Effects of soil amendments on nematodes in tomato cultivation. In: *Proc. Vth. Ann. Confer. Hort. Soc., Nigeria* 21 - 25 Nov., Zaria, Nigeria, 155P.
- 28 - Babu, S.; K. Seetharaman; R. Nonda kumar; and I. Johnson (2000c) Biocontrol efficacy of *Pseudomonas fluorescense* against *Alternaria solani* and tomato leaf blight disease. *Annals. Plant Prot. Sci.*, **8** (2): 252 - 254.
- 29 - Badra, T.; M.A. Salem; and B.A. Oteifa (1979). Nematocidal activity of some organic fertilization and soil amendments. *Revue de Nematologie*, **2**: 29 - 36.
- 30 - Bai, X.; J. Zhang; A. Ewing; S.A. Miller; A. G. Radek; D.V. Shevchenko; K. Tsukeman; J. Walunas; A. Lapidus; J. W. Campbell; and S.A. Hogenhout (2006). Living with genome instability: the adaptation of phytoplasmas to diverse environments of their insect and plant hosts. *J. Bacteriol.* **188**: 3682 - 3696.
- 31 - Balasubramanian, P.; and C. Ramakrishnan (1983). Resistance to the reniform nematode *Rotylenchulus reniformis* in tomato. *Nematologia Mediterranea*. **11**: 203 - 204.

- 32 - Ball - Coelho. B.R.; L.B.Reynolds; A.J. Back; and J.W. Potter (2001). Residue decomposition and soil nitrogen are affected by mowing and fertilization of marigold. *Agron. J.*, **93** (1): 207 - 215
- 33 - Balogh. B.; J.B. Jones; M.T. Momol; S.M. Olson; A. Oradovic; L.King; and L.E. Jackson (2002). Efficacy of bacteriophage formulations for control of bacterial spot on tomato. *Phytopathology*, **92**: 86.
- 34 - Banuelos. G.S.; G.P. Offermann; and E.C. Seim (1985). High relative humidity promotes blossom end rot on growing tomato fruit. *Hort Science*. **20**: 894 - 895.
- 35 - Barbara. D.J.; and E. Clewes (2003). Plant pathogenic *Verticillium* species: how many of them are there?. *Molec. Plant. Pathol.* **4** (4): 297 - 305.
- 36 - Bar - Joseph, M. (2003). Natural history of viroids - horticultural aspects. Pages 246 - 251 in: *Viroids*, A. Hadidi; R. Flores; J.W. Randles; and J.S. Semancik. eds. CSIRO Publishing, Collingwood, Australia.
- 37 - Baul. J.; J. Jones; and B. Jones (1984). Target spot of tomato: epidemiology and control. *Proc. Florida State Hort. Soc.* **97**: 216 - 218.
- 38 - Beachy, R.N.; S. Loeesch - Fries; and N. E. Tumer (1990). Coat protein- mediated resistance against virus infection. *Ann. Rev. Phytopathol.* **28**: 451 - 472.
- 39 - Belanger, R. R.; and W.R. Jarvis (1994). Occurrence of powdery mildew (*Erysiphe sp.*) on greenhouse tomatoes in Canada. *Plant Dis.* **78**: 640.
- 40 - Bell. M.R.; and C.L. Romine (1930). Tobacco budworm field evaluation of microbial control in cotton using *Bacillus thuringiensis* and a nuclear polyhedrosis virus with a feeding adjuvant. *J. Econ. Entomol.* **73**: 427 - 490.
- 41 - Benson, A. P.; and R.P. Singh (1964). Seed transmission of potato spindle tuber virus to tomato. *Amer. Potato. J.* **41**: 294.
- 42 - Bernhardt. J.L.; and J.R. Phillips (1985). Identification of eggs of the bollworm, *Heliothis zea* (Boddie), and the tobacco budworm, *Heliothis virescens* (F). *Southwestern Entomol.* **10**: 236 - 238.
- 43 - Bertagnolli, B.L.; F.K. Dal Soglio; J.B. Sinclair; and D.M. Eastburn (1994). Extracellular enzymes involved in the potential biocontrol of *Rhizoctonia solani* by *Bacillus megaterium* and *Trichoderma harizianum*. *Phytopathology*, **84**: 1136. (Abst.).
- 44 - Bertamini, M.; M.S. Grando; and N. Nedunchezian (2004). Effects of phytoplasma infection on pigments, chlorophyll, protein complex and photosynthetic activities in field grown apple leaves. *Biologia Plantarum*, **47** (2): 237 - 242.
- 45 - Blachinski. D.; D. Shienberg; A. Dincor; U. Kafkafi; L.S. Sujkowski; T.A. Zitter; and W.E. Fry (1996). Influence of foliar application of nitrogen and potassium on Alternaria disease on potato, tomato and cotton. *Phytoparasitica*, **24** (4): 281 - 292.
- 46 - Boll. J.F. (2003). Alfalfa mosaic virus: coat protein - dependent initiation of infection. *Mol. Plant Pathol.* **4** (1): 1 - 8.
- 47 - Boon, J. Van Der (1973). Influence of K/ Ca ratio and drought on physiological disorders on tomato. *Netherlands J. Agric. Sci.* **21**: 56 - 67.

- ☛ - Boonham, N.; L. Gonzalez - Pérez; M.S. Mendez; E. Lilia Peratta; A. Blockley; K. Walsh; I. Barker; and R.A. Mumford (2004). Development of a real - time RT - PCR assay for the detection of potato spindle tuber viroid. *J. Virol. Methods*, **116**: 139 - 146.
- ☛ - Boston, H.; X. Nie; and R.P. Singh (2004). An RT. PCR primer pair for the detection of Pospiviroids and its application in surveying ornamental plants for viroids. *J. Virol. Methods*, **116**: 189 - 203.
- ☛ - Braun, M.; A. M. Gamal Fadl El Din; and H. Burgstaller (1985). Approaches for the control of the parasitic weed *Orobanche ramosa* L. with special references to the use of glyphosate spraying and solar heating techniques on eggplant under the conditions of Sudan. *Acta Horticulturac*, **158**: 335 - 345.
- ☛ - Buschman, L.L. (1980). Parasites of *Nezara viridula* (Hemiptera, Pentatomidae) and other Hemiptera in Florida. *Florida Entomol.* **63**: 154 - 162.
- ☛ - Butler, E.E.; and L. J. Peterson (1970). Sexual reproduction in *Geotrichum candidum*. *Science*, **169**: 481 - 482.
- ☛ - Campbell, R.N.; V.H. Schweers; and D.H. Hall (1982). Corky root of tomato in California caused by *Pyrenochaeta lycopersici* and control by soil fumigation. *Plant Dis.* **66**: 657 - 661.
- ☛ - Candresse. T.; A. Marais; F. Ollivier; E. Verdin; C. Blancard (2007). First report of the presence of tomato apical stunt viroid in Sénégal. *Plant Dis.* **91** (3): 330.
- ☛ - Canfield, M.L.; and L.W. Moore (1992). Control of crown gall in apple (*Malus*) rootstalks using copac E and Terramycin. *Phytopathology*, **82**: 1153 (Abst).
- ☛ - Canthia, M. O. (2006). Tomato (*Lycopersicon esculentum*) early blight. (<http://plant.disease.ippe.orst.edu/disease.cfm?RecordID=1082>) 1 - 2P.
- ☛ - Capinera, J.L. (2001). Handbook of vegetable pests. Academic Press, San Diego. 729 p.
- ☛ - Capinera, J.L. (2001). Vegetable leaf miner. ([http://www.entnemdept.ufl.edu/creatures/veg/Leaf/vegetable leaf miner. htm](http://www.entnemdept.ufl.edu/creatures/veg/Leaf/vegetable%20leaf%20miner.htm)) 1 - 6 p.
- ☛ - Caswell, E.P.; J. de Frank; W.J. Apt; and C.S. Tang (1991). Influence of nonhost plants on population decline of *Rotylenchulus reniformis*. *J. Nematol.* **23**: 91 - 98.
- ☛ - Cavalier - Smith, T. (1998). A revised six kingdom system of life. *Biol. Rev. Camb. Philos. Soc.* **73** (3): 203 - 266.
- ☛ - Cerkauskas, R. (2004). Tomato black leaf mold. AVRDC Publication 04 - 606. 1 - 2 P.
- ☛ - Cerkauskas, R. (2005). Tomato target spot AVRDC Publication 05 - 630. 1 - 2 P.
- ☛ - Chapman, P. J.; and S.E. Lienk (1981). Flight periods of adults of cutworms, armyworms, loopers and others (family Noctuidae) injurious to vegetable and field crops. *New York Agric. Exp. Sta. Search; Agric. Bull.* **14** - 43. pp.
- ☛ - Charles, W.B.; and R.E. Harris (1972). Tomato fruit set at high and low temperatures. *Canada J. Plant Sci.* **52**: 497 - 506.
- ☛ - Chatton, E. (1937). Titres et travaux scientifiques. Sette, Sottano Italy.
- ☛ - Chen, Y.D.; and T.A. Chen (1998). Expression of engineered antibodies in plants: A possible tool for spiroplasma and phytoplasma disease control. *Phytopathology*, **88**: 1367 - 1371.

- 67 - Cherif, M.; and N. Benhamou (1990). Cytochemical aspects of chitin breakdown during the parasitic action of a *Trichoderma* sp. on *Fusarium oxysporum* f. sp. *radicis lycopersici*. *Phytopathology*. **80**: 1406 - 1414.
- 68 - Chiang, B. T., M.K. Nakhla; P. Maxwell; M. Schoenfelder; and S.K. Green (1997). A new geminivirus associated with a leaf curl disease of tomato in Tanzania. *Plant Dis.* **81**: 111.
- 69 - Chienttui, Wu - Wanshi; C.H. Liu; and W.S. Wu (1997). Chemical and biological control of tomato early blight. *Plant Pathol. Bull.* **6** (3): 132 - 140.
- 70 - Chilton, M.D.; M.H. Drummond; D.J. Merio; D. Sciaky; A.L. Montoya; M. P. Gordon; and E.W. Nester (1977). Stable incorporation of plasmid DNA into higher plant cells: the molecular basis of crown gall tumorigenesis. *Cell*, **11** (2): 263 - 271.
- 71 - Christensen, N.M.; M. Nicolaisen; M. Hansen; and A. Schulz (2004). Distribution of phytoplasma in infected plants as revealed by real-time PCR and biomaging. *Mol. Plant Microbe Interact.* **17**: 1175 - 1184.
- 72 - Chunwongse, J.; C. Chunwongse; L. Black; and P. Hanson (2002). Molecular mapping of the Ph-3 gene for late blight resistance in tomato. *J. Hort. Sci. Biotech.* **77** (3): 281 - 286.
- 73 - Cohen, S.; and V. Melamed - Madjar (1978). Prevention by soil mulching of the spread of tomato yellow leaf curl virus transmitted by *Bemisia tabaci* (Gennadius) (Homoptera, Aleyrodidae) in Israel. *Bull. Entom. Res. Israel*, **68**: 465 - 470.
- 74 - Coll, M.; S. Gavish; and I. Dory (2000). Population, biology of the potato tuber moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae) in two potato cropping systems in Israel. *Bull. Entom. Res.* **90**: 309 - 315.
- 75 - Condé, B.; and M.I. Connelly (1994). Australian tomato leaf curl virus: a whitefly-transmitted geminivirus causing a disease of tomatoes in Northern Australia. *Proc. 4th. Inter. Conf. Plant Prot. Kuala Lumpur, Malaysia*, 28 - 31 (3) 1994. PP. 235 - 237.
- 76 - Condé, B.; and M. Connelly (2006). Tomato leafroll - A serious disease in the top end. ([www. nt. gov. au/dpi/](http://www.nt.gov.au/dpi/)) 1 - 4 P.
- 77 - Conover, R.A.; and N.R. Gerhold (1981). Mixture of copper and maneb or mancozeb for control of bacterial spot of tomato and their compatibility for control of fungus diseases. *Proc. Fladelfia State Hort. Soc.* **94**: 154 - 156.
- 78 - Copeland, H.F. (1938). The Kingdoms of organisms. *Quart. Rev. Biol.* **13**: 383 - 420.
- 79 - Copeland, H.F. (1956). The classification of lower organisms. Palo Alto: Pacific Books.
- 80 - Cornelis, P. (ed.) (2008). *Pseudomonas*. *Genom. Molec. Biol.* 1st ed. Caister Academic Press. ISBN 978 - 1 - 904455 - 19 - 6.
- 81 - Costea, M. (2007), Digital Atlas of *Cuscuta* (Convolvulaceae). ([http:// www. wlu. ca/ plant. php? grp- id = 2147 & P= 8968](http://www.wlu.ca/plant.php?grp-id=2147&P=8968)) 1 - 4 P.
- 82 - Cranshaw, W.S. (1999). Flea beetles. Colorado Univ. Fact Sheet no 5 - 592. 1 - 4 P.
- 83 - Cranshaw, W.S. (2002). Hornworms and hummingbird ([http:// www. ext. Colostate. edu. PUBS/ INSECT/ 05517 - html.](http://www.ext.colostate.edu/PUBS/INSECT/05517.html)) 1 - 5 P.

- 54 - Crow, W.T.; and A.S. Brammer (2001). Sting nematode *Belonolaimus longicaudatus*. Univ. Florida, Public. No. : EENY - 239. 1 - 4 P.
- 55 - Crow, W.T. (2004). Stubby root nematode, *Paratrichodorus minor* (Colbran). ([http://creatures.ifas.ufl.edu/nematode/stubby root/ paratrichodorus- minor. htm.](http://creatures.ifas.ufl.edu/nematode/stubby%20root/paratrichodorus-minor.htm)) 1 - 4 p.
- 56 - Culver, J.N.; A.G.C. Lindbeck; and W.O.Dawson (1991). Virus host interaction. Induction of chlorotic and necrotic responses in plants by Tobamoviruses. *Ann. Rev. Phytopathol.* 29: 193 - 217.
- 57 - Cunnington, J. H.; S. Takamatsu; A. C. Lawrie; and I.G. Pascoe (2003). Molecular identification of anamorphic powdery mildews (Erysiphales). *Australian Plant Pathol.* 32: 321 - 428.
- 58 - Czosnek, H.; and H. Latterot (1997). A worldwide survey of tomato yellow leaf curl viruses. *Archives of virology*, 142: 1391 - 1406.
- 59 - Daelemans, A. (1989). Soil solarization in west cameron: effect on weed control, some chemical properties and pathogens of the soil. *Acta Horticulturae.* 255: 169 - 175.
- 60 - Dahlston, D.L.; D. M. Kent; D. L. Rowney; W.A. Cooper; T.E. Young; and R.L. Tassan (1995). Parasitoid shows potential for biocontrol of eugenia psyllid. *California Agric.* 49 (4): 36 - 40.
- 61 - Dahlston, D.L.; K. M. Daane; and T. D. Paine (2005). Imported parasitic wasp helps control red- gum lerp psyllid. *California Agric.* 59 (4): 229 - 234.
- 62 - Daros, J. A.; S.F. Elena, and R. Flores (2006). «Viroids: an Ariadne's thread into the RNA labyrinth». *EMBO Rep.* 7 (6): 593 - 598.
- 63 - Darwesh, Fayza, M. A. (2002). Effect of different fertilizer sources and levels on growth, yield and quality of tomatoes. Ph. D. Thesis Fac. Agric. Cairo Univ. Egypt, 175 pp.
- 64 - Davet, P. (1972). Research on *Colletotrichum coccodes* (Wallr) Hyghes IV. Development of the fungus on tomato in relation to plant and root age. *Phytopath. Mediterr.* 11: 103 - 108.
- 65 - Deacon, J.; A. Robertson; and A. Isbister (2006). The microbial world: Biology and control of crown gall (*Agrobacterium tumefaciens*). ([http:// helios. bto. ed. ac. Uk/ bto/ microbes/ crown. htm.](http://helios.bto.ed.ac.uk/bto/microbes/crown.htm)) 1 - 12 p.
- 66 - Deahl, K.L.; P. De Muth; S.L. Sinder; and A Rivera- Pena (1995b) Identification of mating types and metalxyl resistance in North American populations of *Phytophthora infestans*. *American Potato J.* 72 (1): 35 - 50.
- 67 - De Bokx. J. A.; and P.G. Pirone (1981). Transmission of potato spindle tuber viroid by aphids. *Netherland J. Plant Pathol.* 87: 31.
- 68 - Debolt, J.W.; W.W. Wolf; T.J. Henneberry; and P.V. Vail (1979). Evaluation of light traps and sex pheromone for control of cabbage looper and other lepidopterous insect pests of lettuce. *USDA Technical Bull.* 1606. 39 PP.
- 69 - Decreamer. W. (1991). Stubby root and virus vector nematodes. P. 587 - 625 In: Nickle W.R. (ed.). *Mann. Agric. Nemat.* Marcel Dekker Inc. New York.
- 70 - DEFRA (2007). First UK findings of columnnea latent viroid on tomato production nurseries. Department for Environment, Food and Rural Affairs (UK). ([http:// www. defra. gov. uk/ planth/ newsitem/ clvd. htm.](http://www.defra.gov.uk/plant/newsitems/clvd.htm)) 1p.

- 101 - DEFRA (2007). First UK finding of tomato chlorotic dwarf viroid. Department for Environment, Food and Rural Affairs (UK). (<http://www.defra.gov.uk/plant/newsites/CldVd.htm>) 1 P.
- 102 - Dickinson, M. J. (2007). The phytoplasmas. *Outlooks on Pest Management*, **18**: 160 - 163.
- 103 - Dickson, E.; H.D. Robertson; C.L. Niblett; R. K. Horst; and M. Zaitlin (1979). Minor differences between nucleotide sequences of mild and severe strains of potato spindle tuber viroid. *Nature*, **277**: 60 - 62.
- 104 - Diener, T.O. (1971). Potato spindle tuber virus, IV: A replicating low molecular weight RNA. *Virology*, **45**: 411 - 428.
- 105 - Diener, T. O. (1991). Subviral pathogens of plants: viroids and viroid like satellite RNAs. *The FASEB J.*, **5**: 2808 - 2813.
- 106 - Diener, T.O. (2001). The viroid: biological oddity or evolutionary fossil. *Adv. Virus Res.* **57**: 137 - 184.
- 107 - Diez, J.A.; and D.B. Dusenbery (1989). Preferred temperature of *Meloidogyne incognita*. *J. Nematol.* **21**: 99 - 104.
- 108 - Dillard, H.R. (1987). Tomato anthracnose. (<http://vegetable.mdonline.ppath.cornell.edu/factsheets/tomatoAnth.htm>.) 1 - 3 p.
- 109 - Dillard, H.R. (1989). Effect of temperature, wetness duration, and inoculum density on infection and lesion development of *Colletotrichum coccodes* on tomato fruit. *Phytopathology*, **79**: 1063 - 1066.
- 110 - Dillard, H.R.; and A. C. Cobb (1993). Persistence of *Colletotrichum coccodes* on tomato roots and in soil. *Phytopathology*, **83**: 1345.
- 111 - Dillard, H.R.; and A. C. Cobb (1997). Disease progress of black dot on tomato roots and reduction in incidence with foliar applied fungicides. *Plant Dis.* **81**: 1439 - 1442.
- 112 - Ding, B.; A. Itaya; and X. Zhong (2005). Viroid trafficking: A small RNA makes a big move. *Curr. Opin. Plant Biol.* **8**: 606 - 612.
- 113 - Ding, B.; and A. Itaya (2007). Control of directional macromolecular trafficking across species cellular boundaries: a key to integrative plant biology. *J. Integ. Plant Biol.* **49** (8): 1227.
- 114 - Dinter, Gottlieb, G. (1986). Viroids and virusoids are related to group I introns. *Proc. Natl. Acad. Sci. USA*, **83** (9): 6250 - 6254.
- 115 - Doidge, E.M. (1920). A tomato canker. *J. Dep. Agric. Unions Afr.* **1**: 718 - 721.
- 116 - Doran, J.W.; D.G. Fraser; M.N. Culik; and W.C. Liebhardt (1987). Influence of alternative and conventional agricultural management on soil microbial processes and nitrogen availability. *Amer. J. Altern. Agric.* **2**: 99 - 106.
- 117 - Dorrance, A. E., D.A. Englis; M.L. Derie; C.R. Brown; S. B. Goodwin; W.E. Fry and E.L. Deahl (1999). Characterization of *Phytophthora infestans* populations in western Washington. *Plant Dis.* **83**: 423 - 428.
- 118 - Dreistadt, S.H.; P.A. Phillips; and C.A. O'Donnell (2008). Thrips. (<http://www.ucdavis.edu/PMG/PESTNOTES/pn7429.html>.) 1 - 9 p.

- 11) - Drenth, A.; S.B. Goodwin; W.E. Fry; and L.C. Davidse (1993). Genotypic diversity of *Phytophthora infestans* in the Netherlands revealed by DNA polymorphisms. *Phytopathology*, **83**: 087 - 1092.
- 12) - Drenth, A.; E.M. Janssen; and F. Govers (1995). Formation and survival of oospores of *Phytophthora infestans* under natural conditions. *Plant Pathol.* **44**: 86 - 94.
- 13) - Eisenback, J.D.; and H.H. Triantaphyllou (1991). Root - knot nematodes: *Meloidogyne* species and races. In: Mann, Agric. Nematol., W.R. Nickle. (ED). Marcel Dekker. New York. pp 281 - 286.
- 14) - Eisenberg, H.; Y. Goldwasser; S. Golan; D. Plakhin; and J. Hershenhorn (2004). Egyptian broomrape (*Orobanche aegyptiaca*) control in tomato with sulfonylurea herbicides. *greenhouse studies. Weed Techn.* **18**: 490 - 496.
- 15) - Elad, Y.; J. Kohl; and N.J. Fokkema (1994). Control of infection and sporulation of *Botrytis cinerea* on bean and tomato by saprophytic bacteria and fungi. *European J. Plant Pathol.*, **100** (5): 315 - 336.
- 16) - El Beltagy, A.S.; A.S. Mohamedien; and M. Hafiz (1984). Off - season tomato production. II. Means of improving cold tolerance, frost protection and fruit set for spring tomato production. *Egypt J. Hort.*, **11**: 135 - 149.
- 17) - Ellis, Barbara W.; and F.M. Bradley (1992). The organic gardener handbook of natural insect and disease control. Rodale Press Emmaus, PA. 534 p.
- 18) - Ellis, D. (2008). *Geotrichum candidum*. Univ. Adelaide. Australia. 5005. 1 - 3p.
- 19) - Elmore, C.L.; J.L. Stapleton; C.E. Bell; and J.E. De Vay (1997). Soil solarization: A nonpesticidal method for controlling diseases, nematodes and weeds. *Oakland, Univ. California agric. Nat. Resour Publ.* 21377, 13p.
- 20) - Elphinstone, J.; J. Hennessy; J. Wilson; and D.E. Stead (1996). Sensitivity of different methods for the detection of *Pseudomonas solanaccarum* (Smith) Smith in potato tuber extracts. *EPPO Bull.* **26**: 663 - 678.
- 21) - Erselius, L.J.; M.E. Vega - Sanche; and G.A. Forbes (2000). Stability in population of *Phytophthora infestans* attacked tomato in Ecuador demonstrated by cellulase acetate assessment of Glucose - 6 - phosphate isomerase. *Plant Dis.* **84**: 325 - 327.
- 22) - Erwin, D.C.; and O.K. Ribcro (1996). *Phytophthora* diseases worldwide. *American Phytopath. Soc.* 562 pp.
- 23) - Evans, K.; D.L. Trudgill; and J.M. Webster (1993). *Plant parasitic nematodes in temperate agriculture*. CAB. International, UK, 648 pp.
- 24) - Fagoaga, C.; J.A. Pina; N. Duran - Vila (1994). Occurrence of small RNAs in severely diseased vegetable crops. *Plant Dis.* **78**: 749 - 753.
- 25) - Fagoaga, C. and N. Duran - Vila (1996). Naturally occurring variants of citrus exocortis viroid in vegetable crops. *Plant Pathol.* **45**: 45 - 53.
- 26) - Falk, S.P.; D.M. Gadoury; R.C. Pearson; and R.C. Seem (1995). Partial control of grape powdery mildew by the microparasite *Ampelompses quisqualis*. *Plant Dis.* **79**: 483 - 490.

- 135 - Fay, J.C.; and W.E. Fry (1997). Effects of hot and cold temperatures on the survival of oospores produced by United State strains of *Phytophthora infestans*. American Potato J. **74**: 315 - 325.
- 136 - Fekih - Hassan. I.; F. Gorsané; F. Djilani; H. Fakh - fakh, M. Nakhla; D. Maxwell; and M. Marrakchi (2003). Detection of tomato yellow leaf curl Sardinia virus in Tunisia. EPPO Bull., **33** (2): 347 - 350.
- 137 - Fiume, F. (1995). Control of corky rot caused by *Pyrenochaeta lycopersici* (Schneider et Gerlach) on tomato in protected cultivars. Mededelingen faculteit Landbouwkundige Universiteit Gent **60**: 381 - 392.
- 138 - Fiume, F.; and G. Fiume (2003). Use of culture filtrates of *Pyrenochaeta lycopersici* in tests for selecting tolerant varieties of tomato. J. Plant Pathol. **85** (2): 131 - 133.
- 139 - Flaherty, J.E.; J.B. Jones; B.K. Harbaugh, G.C. Somodi; and L.E. Jackson (2000). Control of bacterial spot on tomato in the greenhouse and field with h - mutant bacteriophages. Plant Science, **35**: 882 - 884.
- 140 - Flores, R.; C. Hernandez; A. E. Martinez de Alba; J.A. Daros; and F. Di Serio (2005) Viroids and viroid host interaction. Ann Rev. Phytopath. **43**: 117- 139.
- 141 - Flynn, p. (2006). White mold on tomato. (http://www.ipm.iastate.edu/ipm/hotnews/2006/6-28/tomato_mold.html) 1- 2p.
- 142 - Foy, C.L.; and R.Rain (1986). Recent approaches for control of parasitic weeds. Annual plant prot **4**: 136 - 144.
- 143 - Fry, W.E.; and S.B Goodwin (1997a). Re- emergence of potato and tomato late blight in the united states. Plant Dis. **81**: 1349 - 1357.
- 144 - Fulling, B.A.; E.C.Tigchelear; and R. Latin (1995) Integration of host resistance and weather-based fungicide scheduling for control of anthracnose of tomato fruit. Plant Dis. **79**: 228- 233.
- 145 - Gaffar, A. (1988). Biological control of sclerotial diseases. pages 153- 175 In: Biocontrol of Plant Diseases. Vol. I, K.G Mukerji and K.L. Garg, eds.CRC Press: Boca Raton, FL, U.S.A.
- 146 - Gandia, M.; L.Bernad, J.Rubio; and N.Duran- Vila (2007). Host effect on the molecular and biological properties of a citrus exocortis viroid isolate from *Vicia faba*. phytopathology, **97**: 1004.
- 147 - Gardan, L.; C.Gouy; R. Christen; and R.Samson (2003). Elevation of three subspecies of *Pectobacterium carotovorum* to species level: *Pectobacterium atrosepticum* sp. nov., *Pectobacterium betavascularum* sp nov. and *Pectobacterium wasabiae* sp. nov. Inter. J. Syst. Evol. Microbiol. **53**: 381 - 391.
- 148 - Gardner, R.G.; and P.B. Shoemaker (1999). Mountain supreme early blight resistant hybrid tomato and its parents, NCEBR - 3 and NCEBR - 4. Hort Science, **34** (3): 745- 746.
- 149 - Gartemann, K.H.; B.Abt; T.Bekel; A. Burger; j. Engemann; M. Fluegel; L.Gaigal; A. Goesmann; I. Graefern; J. kalinowski; O.Kaup O. krichner; L. Krause; B.linke; A.Mehardy; F.Meyer; S.Pohle; C. Ruckert; S.Schneiker; E.M. Cellerman; A Puehler; R. Eichenlatz O.

- Kaiser; and D. Bartels (2008) The genome sequence of the tomato pathogenic actinomycetes *Clavibacter michiganensis* subsp. *michiganensis* NCPPB 382 reveals a large island involved in pathogenicity. J. Bacteriol. **190**: 2138 - 2149.
- 150 - Gavino, P. D.; C. D. Smart; R. W. Sandrock; J.S. Miller; P.B Hamm; T. Yun Lee; R.M. Davis; and W.E.Fry (2000). Implications of sexual reproduction for *Phytophthora infestans* in the United States: Generation of an aggressive lineage. Plant Dis. **84**: 731 - 735.
- 151 - Geraert, E. (2006). Functional and detailed morphology of the Tylenchida (Nematoda). Brill, Leiden ISBN 90 - 04 - 14895 - 7.
- 152 - Ghidui, G.M (1999). Control of insects on tomato. PP 25 - 27 In: Vegetable Entomol. Res Results. Rutgers NJAES Bull. 104 H.
- 153 - Ghidui, G.M; E.M. Hitchner; and J.E.Funderburk (2006). Golden fleck damage to tomato fruit caused by feeding of *Frankliniella occidentalis* (Thysanoptera: Thripidae). Florida Entomol. **89** (2): 279 - 281.
- 154 - Giddings. N.J.: C.W. Bennet; and A.L. Harrison (1951). A tomato disease resembling curly top. Phytopathology, **41**: 415 - 417.
- 155 - Gilreath, J.P.; A.C.Chase; and S.J. Locascio (2001). Crop injury from sublethal rates of herbicides. 1- tomato. Hort Science, **36** (4): 639 - 644.
- 156 - Gomaa, A.E. (2005). The biochemical effect of spinosad for the control of the cotton leafworm, *Spodoptera littoralis* (Boisd). Egypt J. Agric. Res. **83** (1): 33 - 46.
- 157 - Gonzalez - Chavez, M. ;T. Shagarrodsky; D. Barrios; and N. Fraga (2003). Varietal behavior of tomato against early blight under field conditions. Revista de Proteccion Vegetal, **18** (1) : 38 - 41.
- 158 - Goodenough, P.W; and T.H. Thomas (1980).Comparative physiology of field grown tomatoes during ripening on the plant or retard ripening in controlled atmospheres. Ann. Appl. Biol. **94**: 445 - 455.
- 159 - Goodwin, S.B; and A. Drenth (1997). Origin of the A2 mating type of *Phytophthora infestans* outside Mexico. Phytopathology, **87**: 992 - 999.
- 160 - Graham, H.M; and O.T.Robertson (1970). Host plants of *Heliothis virescens* and *Heliothis zea* (Lepidoptera: Noctuidae) in the lower Rio Gande Valley, Texas. Ann. Entmol. Soc. America, **63**: 1261 - 1265.
- 161 - Gravel, V.; C. Martinez; H. Antoun; and R.J.Tweddell (2005). Antagonist microorganisms with the ability to control Pythium damping-off of tomato seeds in Rockwool. Biocontrol, **50** (5) 771 - 786.
- 162 - Gray, S; and F.E.Gildow (2003). Luteovirus - aphid interactions. Ann. Rev. Phytopathol.**41**: 539 - 566.
- 163 - Griffith, G.W.; R.Snell; and D.S.Shaw (1995). Late blight (*Phytophthora infestans*) on tomato in the tropics. Mycologist, **9** (2) : 87 - 89.
- 164 - Grinberger, M.; D. Kadish; and Y. Cohen (1989). Occurrence of the A₂ mating type and oospores of *Phytophthora infestans* in potato crops in Israel. Phytoparasitica, **17**: 197 - 204.

- 165 - Haas, D.; and G. Defago (2005). Biological control of soil - borne pathogens by fluorescent pseudomonads. *Nature Rev. Microbiol.* **3** (4): 307 - 319.
- 166 - Hadidi, A.; R- Flores; J.W. Randles; and J.S. Scmancik (2003). *Viroids*. CSIRO Publishing, Collingwood, Australia. 370p.
- 167 - Hadidi, A.; H. Czosnek; and M. Barba (2004). DNA Microarrays and their potential application for the detection of plant viruses, viroids and phytoplasmas. *J.Plant Pathol.* **5**: 97 -104.
- 168 - Haeckel, E. (1866). *Generelle Morphologie der Organismen*. Reimer, Berlin.
- 169 - Halfeld - Vicira, B.A.; K.L.Nechet; and R.N.T.Barbosa (2006). *Pseudocercospora fuligena* causing leaf mold of tomato in Roraima, Brazil. *Fitopatol. Brasileira*, **31** (3): 320.
- 170 - Halfeld - Vieira, B. A.; K. L. Nechet; and G.R.Souza (2009).influence of different leaf wetness periods in the conidial germination of *Pseudocercospora fuligena*. *Acta Amazonica*, Vol. **39** (2) Manaus. 1- 4 p.
- 171 - Hammond, R.; D.R.smith; and T.O.Diener (1989). Nucleotide sequence and proposed secondary structure of columnea latent viroid: A natural mosaic of viriod sequences. *Nucleic Acids Res.* **17**: 10083 - 10094.
- 172 - Hammond, Rosemaric W.; and R.A Owens (2006). Viroids: New and continuing risks for horticultural and agricultural crops ([http:// www.apsnet.org/online/ feature/ viroids.htm](http://www.apsnet.org/online/feature/viroids.htm)) 1 - 14 P.
- 173 - Hansen, Mary Ann; and R.C Lambe (2000). Septoria leaf spot of tomato. ([http:// www.vt.edu/pubs/plant_disease fs/ 450 -711/ 450- 711. html](http://www.vt.edu/pubs/plant_disease/fs/450-711/450-711.html)) 1- 3p
- 174 - Hartman, G.L.; and T.C.Wang (1992). Black leaf mold development and its effect on tomato yield. *Plant Dis.* **76**: 462 - 465.
- 175 - Hartman, G.L.; and Y.H Huang (1995) Characteristics of *Phytophthora infestans* isolates and development of late blight on tomato in Taiwan. *Plant Dis.* **79**: 849 - 852.
- 176 - Hartstock, A.W; J.A Witz; and D.R.Buck (1979). Moth traps for the tobacco budworm. *Econ. Entomol.***72**: 519 - 522.
- 177 - Hassan, A.A.; M.N stino; S.E. Moustafa; and G.Hanna (1982). Fusarium wilt race identification and screening for resistance in commercial tomato cultivars and wild *Lycopersicon* species. *Egypt. J.Hort.* **9** (2): 125 - 130.
- 178 - Hauben, L.;E.R. Moore; L. Vauterin; M. Steenackers, J. Mergaert; L. Verdonck; and Swings (1998). Phylogenetic position of phytopathogens within the Enterbacteriaceae. *Syst. Appl. Microbiol.* **21**: 384 - 397.
- 179 - Hausbeck, M.K; and S.D.Linderman (1992). Influence of dew period and temperature on infection of tomato foliage by *Colletotrichum coccodes*. *Phytopathology*, **82**: 1091.
- 180 - Hayes, J.L.; and M. Bell (1994) Evaluation of early- season baculovirus treatment for suppression of *Hiliothis virescens* and *Helicoverpa zea* (Lepidoptera: Noctuidae) over a wide area. *J.Econ. Entomol.*, **87**: 58 - 66.

- 188 - Hayslip N.C; and J.R. Iley (1967). influence of potassium fertilizer on severity of tomato graywall. Proc. Amer. Soc. Hort. Sci. **80**: 182 - 186.
- 189 - Henneberry, T. J; G.D. Butler; D.L. Coudriet (1993). Tobacco budworm (Lepidoptera: Noctuidae): Effects of temperature and photoperiod on larval and pupal development, larval mortality and induction of pupal diapause. Southwestern Entomol. **18**: 269 - 279.
- 190 - Hewedy, A.M. (2000). Effect of methods and sources of potassium application on the productivity and fruit quality of some new tomato hybrids. Egypt J. Agric. Res. **78** (1): 227 - 244.
- 191 - Ilijwegen, T. (1992). Biological control of cucumber powdery mildew with *Tilletiopsis minor* under greenhouse conditions. Netherland J. Plant Pathol. **98**: 221 - 225.
- 192 - Hirano, S.S; and C.D Upper (1990). Population biology and epidemiology of *Pseudomonas syringae*. Ann Rev. Phytopathol. **28**: 155 - 177.
- 193 - Hirano, S.S.; and C.D Upprer (2000). Bacteria in the leaf ecosystem with emphasis on *Pseudomonas syringae* - a pathogen ice nucleus and epiphyte. Microbiol. Molec. Biol.Rev. **64**: 624 - 653.
- 194 - Hodgetts, J.; T.Ball; N. Boonham; R.Mumford; and M.Dickinson (2007) Use of terminal restriction fragment length polymorphism (T- RFLP) for identification of phytoplasmas in plants. Plant Pathol. **56** (3): 357 - 365.
- 195 - Hoftman, M. P.; N. Davidson; L.T. Wilson; L.E.Ehler; W.A. Jones; and F.G. Zalom (1991). Imported wasp helps control southern green stink bug. California Agric. **45** (3): 20 - 22.
- 196 - Howell, W.E.; J Burgess; G.I. mink; L.J. Skrzeczkowski; and Y.P. Zhang (1998). Elimination of apple fruit and bark deforming agents by heat therapy. Acta Horticulturae. **472**: 641 - 646.
- 197 - Huang, H.C. (1985). Factors affecting myceliogenic germination of sclerotia of *Sclerotinia sclerotiorum*. Phytopathology, **75**: 433 - 437.
- 198 - Huang - Shao Min; X.Bensheng; T.J.Fang; Z.Y. Ping; J.Q.Wang; and J.L. Qiao (1999). Study on the effect of potassium fertilizer on the prevention and control of tomato early blight. Acta Agriculturae Unversitatalis Henanensis, **33** (2): 138 - 142.
- 199 - Huang, C.C; J.Biesheuvel; P. Lindhout; and R.E. Niks (2000). Host range of *Oidium lycopersici* occurring in the Netherlands. Eur. J. Plant Pathol. **106**: 465 - 473.
- 200 - Hunter, W.B; D.E. Ulman; and A. Moore (1994). Electronic monitoring characterizing the feeding behavior of western flower thrips (Thysanoptera: Thripidae). Entomological Soc. America, Lanham, Maryland. Pp 73 - 85.
- 201 - Hunter, W.B; E.Hiebert; S.E.Webb; J.H. Tsai; and J.E.Polston (1998). Location of geminiviruses in the whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae). Plant Dis. **82**: 1147 - 1151.
- 202 - Hyo Won, J.;H.J.Jung; W.S. Yun; H.J. Kim; Y.Haham, K.H. kim. J.k.Choi (2000). Characterization and partial nucleotide sequence analysis of alfalfa mosaic alfamoviruses isolated from potato and azuki bean in Korea. Plant Pathol. J. **16** (5): 269 - 279.
- 203 - ICTVdB Management (2006). Beet curly top virus. ([http:// www.ncbi.nlm.nih.gov/ ICTVdb/ ICTVdb/...](http://www.ncbi.nlm.nih.gov/ICTVdb/ICTVdb/...) 1- 7P.

- 197 - ICTVdB management (2006). Tomato bushy stunt virus. ([http://www.ncbi.nlm.nih.gov/ICTVdb/ICTVdb/...](http://www.ncbi.nlm.nih.gov/ICTVdb/ICTVdb/) 1- 9P.
- 198 - ICTVdB Management (2006). Tomato spotted wilt virus. (<http://www.ncbi.nlm.nih.gov/ICTVdb/ICTVdb/> 1 - 4p.
- 199 - Infinto, A.; M.Aragona; A. Brunetti; E. Lohaz; A. Oliva; and Porta- Pugalia (2000). Molecular and physiological characterization of Italian isolates of *Pyrenochaeta lycopersi*. Mycol. Res. **107** (6) 707 - 716.
- 200 - Inglis, D.A.; D.A.Johnson; D.E.Ligard; W.E.Fry; and P.B.Hamm (1996) Relative resistance of potato clones in response to new and old populations of *Phytophthora infestans*. Plant Dis. **80**: 575 - 578.
- 201 - Ivanovic, M.; M. Migatovic; D. Antonjvic, G. Paroussi; D. Voyiatzis; and E. Paroussi (2002). Effect of sodium bicarbonate on *Alternaria solani* in tomato. Acta Horticulturae, **579**: 535 - 539.
- 202 - Jackson, C.G.; G.D.Butler; and D.F. Bryan (1969). Time required for development of *Voria ruralis* and its host, the cabbage looper, at different temperatures. J. Econ Entomol. **62** : 69 - 70.
- 203 - Jaim - Garcia, R.;R. Trinidad - Correa; R. Felix Gastelum; T.V Orum; C.C.Wasmann; and M.R. Nelson (2000). Temporal and spatial patterns of genetic structure of *Phytophthora infestans* from tomato and potato in the Del Fuerte Valley. Phytopathology, **90**: 1188: 1195
- 204 - James, C.K.; and B.W Falk (2006). Virus vector interactions mediating nonpersistent and semipersistent transmission of plant viruses. Ann Rev. phytopath. **44** (9): 183 - 212.
- 205 - Jenkins, S.F.; and C.W Averb (1986). Problems and progress in integrated control of southern blight of vegetables. Plant Dis. **70**: 614 - 619.
- 206 - Johnson, M.W; E.R.Oatman; and J.A.Wyman (1980a). Natural control of *Liriomyza sativae* (Diptera: Agromyzidae) in pole tomatoes in southern California. Entomophaga, **25**: 193 - 195
- 207 - Johnson, M.W.; E.R. Oatman; J.A.Wyman; R.A.Van Steen Wyk (1980b) A technique for monitoring *Liriomyza sativae* in fresh market tomatoes. J. Econ. Entomol. **73**: 552 - 555
- 208 - Jones, J.B;K.L. Pohronezny; R.E.Stall; and J.P.Jones (1986). Survival of *Xanthomonas campestris* pv. *vesicatoria* in Florida on tomato crop residues, weeds, seeds and volunteer tomato plants. Phytopathology, **76**: 430 - 434 .
- 209 - Jones, M.E.; J.M. Whipps; B.J. Thomas; T.L.W. Carver; and S.J. Gurr (2000) Initial events in the colonization of tomatoes by *Oidium lycopersici*, a distinct powdery mildew fungus of *Lycopersicon* species. Can. J. Bot. **78**: 1361 - 1366.
- 210 - Jones, H.; J.M Whipps; and S.J Gurr (2001). The tomato powdery mildew fungus *Oidium neolycopersici*. Molec. Plant Pathol. **2** (6) 303 - 309.
- 211 - Jones, M.E. (2001). Identification and development of tomato powdery mildew fungi, *Oidium lycopersici*. Ph.D Thesis Oxford University.
- 212 - Jones, J.B.; A Obradovic; B.Balogh; M. T. Momol; and L.E. Jackson (2002). Control of bacterial spot on tomato with bacteriophages. (Abstr.) Phytopathology, **92**: S108.

- 213 - Jones, J.B.; L.E. Jackson; B. Balogh; A. Obradovic; and M.T. Momol (2007). Bacteriophages for plant disease control. *Ann. Rev. Phytopath.* **45**: 245 - 262.
- 214 - Joseph, Scamrdella M. (1999). Not plants or animals, a brief history of the origin of kingdoms Protozoa, Protista and Protoctesta. *Inter. Microbiol.*, **2**: 207 - 221.
- 215 - Jothi, G.; S. Pugalcndhi; K. Poornima; and G. Rajendran (2003). Management of root- knot nematode in tomato. *Lycopersicon esculentum* Mill. with biogas slurry. *Bioresource Tchn.* **89**: 169 - 170.
- 216 - Juarez - Palacios, C.; R. Felix - Gastelum; R.J.Wakeman; E.J. Paplomatas; and J.E Devay (1991). Thermal sensitivity of three species of *Phytophthora* and the effect of soil solarization on their survival. *Plant Dis.* **75**: 1160 - 1164.
- 217 - Kadder, A.A. (1986). Biochemical and physiological basis for effects of controlled and modified atmospheres on fruits and vegetables. *Food Technol.* **40** (5): 99 - 104.
- 218 - Kadder, A.A (2002). Postharvest biology and technology: an overview. In: Kadder, A.A.(ed) *Postharvest Technology Hort. Crops*, 3rd ed. Pub. no. 3511 Oakland Univ. California PP. 39 - 47.
- 219 - kang, L.H.B; F.W.Plapp; G.W.Elzen; M.L.Wall; J.D. Lopez (1995). Monitoring for resistance to organophosphorus, carbonate, and cyclodiene insecticides in tobacco budworm adults (Lepidoptera: Noctuidae). *J.Econ. Entomol* **88**: 1144 - 1149.
- 220 - kang, B.; l. Yeam; and M.M. Jahn (2005). Genetics of plant virus resistance. *Ann. Rev. Phytopathol.* **43** (9): 581 - 621.
- 221 - kashimoto, k.; Y.Matsuda; K.Matsutani, T.Sameshara; K.Kakutani; T.Nonomura; K.Okada; S. Kusakari; K.Nakata; S.Takamatsu; and H.Toyoda (2003). Morphological and molecular characterization for a japanese isolate of tomato powdery mildew *Oidium neolycopersici* and its host range. *J.Gen. Plant Pathol.* **69**: 176 - 185.
- 222 - Katan, J.(1987). Soil solarization. PP. 77- 105, *Inovative approaches to plant discase control*. I.Chet, ed.. John Wiley & Sons, New york.
- 223 - Kato, M.; E.S.Mizubuti; S.B.Goodwin; and W.E Fry (1997) Sensitivity to protectant fungicides and pathogenic fitness of clonal lineages of *Phytophthora infestans* in the United States. *Phytopathology*, **87**: 973 - 978.
- 224 - Kawaguchi, A.; K.Inoue; and Y. Ichinose (2008). Biological control of crown gall of grapevine, rose, and tomato by nonpathogenic *Agrobacterium vitis* strain VARO 3 - 1. *Phytopathology*, **98** (11): 1218 - 1225.
- 225 - Kay, L. E. (1986). W.M.Stanley's crystallization of the tobacco mosaic virus, 1930 - 1940. *Isis* **77**: 450 - 472.
- 226 - Kaya, C. and D. Higgs (2002). Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. *Scientia Horticulturae*, **93**: 53- 64.
- 227 - Keoreab, E.; and A.J. Murdoch (2001). Stimulation of intergrated control strategies for *Orobanche spp.* based on a life cycle model. *Experim. Agricul.* **37**: 37 - 51.

- 228 - Keinath, A.P.; V.B. DuBose; and P.J. Rathwell (1996). Efficiency and economics of three fungicide application schedules for early blight control and yield fresh market tomato. *Plant Dis.* **80** (11): 1277 - 1282.
- 229 - Kengo, Y.; and A. Kazuo (2002). Race composition of *Fluvial fulva* in Japan during 1997 - 1998. *Ann. Phytopathol. Soc. Japan*, **68** (1): 36 - 38.
- 230 - Khey - Pour, A.; M. Bendahmane; V. Matzeit; G.P. Accoto; S. Crespi; and B. Groenewald (1991). Tomato yellow leaf curl virus from Sardinia is a whitefly transmitted monopartite geminivirus. *Nucleic Acids Res.* **19**: 6763 - 6769.
- 231 - Kim, J.T.; I.H. Park; K.Y. Ryu; J.U. Cheon; and S.H. Yu (2003). Corky root of tomato caused by *Pyrenochaeta lycopersici* in Korea. *Plant Pathol. J.* **19** (3): 181 - 183.
- 232 - Kiss, L. (1996). Occurrence of a new powdery mildew fungus (*Erysiphe sp.*) on tomatoes in Hungary. *Plant Dis.* **80**: 224.
- 233 - Kiss, L.; R.T.A. Cook; G.S. Saenz; J.H. Cunningham; I. Pascoe; M. Bardin; P.C. Neco; S. Takamatsu; Y. Sato; and A.Y. Rossman (2001). Identification of two powdery mildew fungi, *Oidium neolycopersici* sp. nov. and *O. lycopersici*, infected tomato in different parts of the world. *Mycol. Res.* **105**: 684 - 697.
- 234 - Kiss, L.; S. Takamatsu; and J.H. Cunningham (2005). Molecular identification of *Oidium neolycopersici* as the causal agent of the recent tomato powdery mildew epidemics in North America. *Plant Dis.* **89**: 491 - 496.
- 235 - Kleineke, Brochers, A. (1982). Effect of the interaction between *Meloidogyne incognita* and *Fusarium oxysporum* f. sp. *lycopersici* on phytohormone content in tomato plants (CPL Rev. *Plant Pathol.* **61**: 5219).
- 236 - Klug, A. (1999). The tobacco mosaic virus particles: structure and assembly. *Biochemical Science* **354**: 531 - 536.
- 237 - Knight, D. E.; and W.G. Keyworth (1960). Didymella stem rot of outdoor tomatoes. Studies on sources of infection and their elimination. *Ann. Appl. Biol.* **48** (2): 245 - 258.
- 238 - Kondoh, M.; M. Hirai; and M. Shoda (2001). Integrated biological and chemical control of damping-off caused by *Rhizoctonia solani* using *Bacillus subtilis* RB 14 - C and fludioxonil. *J. Biosci. Bioeng.* **91** (2): 173 - 177.
- 239 - Kring, J.B.; D.J. Schuster; J.F. Price; and G.W. Simone (1991). Sweetpotato whitefly - vectored geminivirus on tomato in Florida. *Plant Dis.* **75**: 1186.
- 240 - Kryczynski, S.; E. Paduch - Cichal; and L.J. Skzeczowski (1988). Transmission of three viroids through seed and pollen of tomato plants. *J. Phytopath.* **121**: 51 - 54.
- 241 - Kubota, M.; K. Kishi; and K.A. Biko (2000). Phoma leaf spot, stem and fruit rot caused by *Phoma lycopersici* Cooke in Japan. *Japan J. Phytopath.* **66**: 12 - 17.
- 242 - Kuepper, G. (2003). Flea beetle: organic control options (<http://attra.ncat.org/attra-pub/flea-beetle.html>). 1 - 6 p.
- 243 - Kuske, C.R.; and B.C. Kirkpatrick (1992). Phylogenetic relationships between the western aster yellows mycoplasma like organisms and other Prokaryotes established by 16S rRNA gene sequence. *Inter Sys. Bacteriology* **42**: 75 - 80.

- 244 - Lamp, R.C.; and H.Mary Ann (2000). Buckeye rot of tomato. (<http://pubs.ext.vt.edu/450/450-704.html>) 1- 3p
- 245 - La Mondia, J.A. (1999). Host range of *Oidium lycopersicon* selected solanaceous species in Connecticut. *Plant Dis.* **83**: 341 - 344.
- 246 - Lebeda, A. and B.Miestrova (1999). Identification, occurrence and host range of tomato powdery mildew (*Oidium lycopersici*) in the Czech Republic. *Acta Phytopathol. Entomol. Hung.* **34**: 15 -27.
- 247 - Leccointre, G.; and H.L. Guyader (2006). The tree of life: a phylogenetic classification. Harvard Univ. Press, Cambridge, Massachusetts.
- 248 - Lee, I.M.; D.E.Gundersen - Rindal; and A.Bertaccini (1998). Phytoplasma: Ecology and genomic diversity. *Phytopathology*, **88**: 1359 - 1366.
- 249 - Lee, I.M.; R.E.Davies; and D.E.Gundersen - Rindal (2000). Phytoplasma: phytopathogenic mollicutes. *Ann. Rev. Microbiol.* **54**: 221 - 255.
- 250 - Legard, D.E.; T.Y. Lee; and W.E.Fry (1995). Pathogenic specialization in *Phytophthora infestans* aggressiveness on tomato. *Phytopathology*, **85**: 1356 - 1361.
- 251 - Leite, G. L.D. ; C.A. Costa; C.I.M.ALmeida; and M.Picanco (2003). Effect of fertilization on the incidence of leaf miner and early blight on tomato plants. *Horticultura Brasileira*, **21** (3): 448 - 451.
- 252 - Lema, K.M.; and S.L. Poe (1979). Age specific mortality of *Liriomyza sativa* due to *Thrysonotomyia formosa* and parasitization by *Opius dimidiatus* and *C. formosa*. *Environ. Entomol.* **8**: 935 - 937.
- 253 - Lepka, p.; M.Stitt; E.Moll.; and E.Seemiller (1999). Effect of phytoplasmal infection on concentration and translocation of carbohydrates and amino acids in periwinkle and tobacco. *Physiol. Mol. Plant Pathol.* **55**: 59 - 68.
- 254 - Lewis, T. (1997). Feeding and oviposition injuries to plants, PP 505 - 537. In: T. Lewis (ed), *Trips as Crop Pests*. CAB International NY.
- 255 - Li Youzhou; Chen - Zhiri; Liu - Yong Feng; and Xu - Yang (2003). Effect of antagonistic bacteria for the control of common disease in solanaceous vegetables. *Jiangsu J. Agric. Sci.*, **15** (2) : 96 - 99.
- 256 - Lopez, J.D.; J.L. Goodenough; K.R.Beerwinkle (1994). Comparison of two sex pheromone trap designs for monitoring corn earworm and tobacco budworm (Lepidoptera. Noctuidae). *J.Econ.Entomol.* **87**: 793 - 801.
- 257 - Louws, F.J.; M.Wilson; H.L. Cambell; D.A. Cuppels; J.B.Jones; P.B Shoemaker; F.Sahin; and S.A. Miller (2001). Field control of bacterial spot and bacterial speck of tomato using a plant activator. *Plant Dis.* **85**: 481 - 488.
- 258 - Lye, B. H.; R.N.Storry; and V.L.Wright (1988). Southern green stink bug (Hemiptera: Pentatomidae) damage to fresh market tomatoes. *J. Econ. Entomol.* **81**: 190 - 194.
- 259 - Machad, o, M.A, and K. Zetsche (1990). A structural, functional and molecular analysis of peptides of the holoparasites *Cuscuta reflexa* and *Cuscuta europaea*, *Planta*, **181**: 91 - 96.

- 260-Madula, J.; and D. L. Trudgill (1994). Influence of temperature on *Meloidogyne javanica*. *Nematologica* **40**: 230 - 243.
- 261-Mahovic, M.; S.A Sargent, and J.A. Bartz (2005) Identifying and controlling postharvest tomato diseases in Florida. (<http://edis.ifas.ufl.edu/HS1310>) 1 - 13 p
- 262-Mahovic, M.J., J.D. Tenney; and J.A. Bartz (2007). Application of chlorine dioxide gas for control of bacterial soft rot in tomatoes. *Plant Dis.* **91**(10) 1316 - 1320
- 263-Mai., W.F.,; and P. G. Mullin (1996). Plant parasitic nematode. A Pictorial Key to Genera, 5th Ed. Cornell Univ. Press, Ithaca. New York
- 264-Maiero, M.; T.J. Ng; and T.H. Barksdale (1990). Inheritance of collar rot resistance in the tomato breeding lines C 1943 and NCEBR-2. *Phytopathology* **80**: 1365 - 1368
- 265-Maki, L.R.; E.L. Galyon; M.M. Chang-Chien; and D.R. Coldwell (1974). Ice nucleation induced by *Pseudomonas syringae*. *Appl. Microbiol* **28** (3): 456 - 459
- 266-Marcone, C.; H. Neimark; A.Ragozzino; U.Lauer; and E. Seemiller (1999). Chromosome sizes of phytoplasmas comprising major phylogenetic groups and subgroups. *Phytopathology* **89**: (9) 805 - 810,
- 267-Margulis, L.; K.V. Schwartz; and M. Dotan (1994). The illustrated five kingdoms: A guide to the diversity of life on earth. Harp Collins College Publishers, New York.
- 268-Marko, J.M. and R.E. Stall (1983). Control of bacterial spot of pepper initiated by strains of *Xanthomonas campestris* pv. *vesicatoria* that differ in sensitivity to copper, *Plant Dis* **67**: 779 - 781
- 269-Marois, J.J.; M.Y. Momol; J.W. Kimbrough, R.C. Hochmuth; and W. Danker (2001). First report of powdery mildew on greenhouse tomatoes caused by *Oidium neolycopersicij* in Florida. *Plant Dis.* **85** : 1292
- 270-Martin, P.B; P.D Lingren; G.L Greene; and R.L. Ridgway (1976b). Parasitization of two species of *Plusiinae* and *Heliothis spp.* after releases of *Trichogramma pretiosum* in seven crops. *Environ. Entomol* **5**: 991 - 995
- 271-Martin, R.R; P.K.; Keese; M.J. Young, P.M. Waterhorse; and W.L. Gerlach (1990). Evolution and molecular biology of Luteoviruses. *Ann. Rev. Phytopathol.* **28**: 341 - 363
- 272-Martin, R; C. Arenas; J.A. Daros; A. Covarrubias; J.L. Reyes; N.H. Chua (2007) Characterization of small RNAs derived from citrus exocortis viroid CEVd in infected tomato plants. *Virology*, **367** (1): 135 - 146
- 273-Matsuda, Y., K. Kashimoto; Y. Takikawa; R. Aikami; T. Nonomura; and H. Toyoda (2001). Occurrence of new powdery mildew on greenhouse tomato cultivars. *J. Gen. Plant Pathol.* **67**: 294 - 298
- 274-Matsuda, Y., T. Sameshima; N. Moriura; K. Inoue; T. Nonomura; K. Kakutani, H. Nishimura; S.Kusakari; S. Takamatsu; and H. Toyoda (2005) Identification of individual powdery mildew fungi infecting leaves and direct reaction. *Phytopathology*, **95**: 1137 - 1143

- 275-Matthijs, S.; K.A. Tehrani, G. Laus; R. W. Jackson; R.M. Cooper. and P. Cornelis (2007) Thioquinolobactin, a *Pseudomonas siderphore* with antifungal and anti - *Pythium* activity. *Environ. Microbiol.* **9**: (2) 425 - 434
- 276-Matuszak, J.M.; J. Fernandez- Elquezabel; W.K. Gu; M. Villarreal- Gonzalez; and W.E. Fry (1994). Sensitivity of *Phytophthora infestans* populations to metalaxyl in Mexico: distribution and dynamics. *Plant Dis.* **78**: 911 - 916
- 277 Maul, F., S. A. Sargent; M.O. Balaban; E.A. Baldwin; D.J. Huber; and C.A. Sims (1998) Aroma volatile profiles from ripe tomato fruit are influenced by physiological maturity at harvest: an application for electronic nose technology. *J. Amer. Soc. Hort. Sci.* **123** (6): 1094- 1101.
- 278 Maul, F., S. A. Sargent; C.A. Sims; E.A. Baldwin; M.O. Balaban; and D.J. Huber (2000). Recommended commercial storage temperature effect tomato flavor and aroma quality. *J. Food. Sci.* **65** (7): 1228 - 1237
- 279 Mc Daniel, L. L. and J.H. Tsai (1990). Partial characterization and serological analysis of Pseudo-curly top virus. *Plant Dis* **74**: 17 - 21
- 280 Mc Pherson, R.M.; and L.D. Newsom (1984). Trap crops for control of stink bugs in soybean. *Georgia Entomol-Soc.* **19**: 470 - 489
- 281 Mee Ngan, Y.; J.D. Barak; and A.O. Charkowski (2004). Genomic diversity of *Erwinia caratovora* subs *caratovora* and its correlation with virulence *Appl. Environ Microbiol.* **70**: 3013 - 3023
- 282 Mencarelli, F.; and M.E. Saltveit (1988). Ripening of mature-green tomato fruit slices. *J. Amer. Soc. Hort. Sci.* **113** (5): 742 - 745
- 283 Mendez, R.S.; E. Bustamanta; and F. Merino (1994). The effect of different source and level of calcium on severity of early blight (*Alternaria solani*) on tomato (*Lycopersicon esculentum*). *Manj's Integrado de Plagas*, **33**: 1 - 6
- 284 Meyer, J.M.; V.A. Geoffroy; N. Baida (2002). Siderphore typing, a powerful tool for the identification of the fluorescent and nonfluorescent pseudomonads. *Appl. Environ. Microbiol.* **68** (6): 2745 - 2753
- 285 Miestrova, B.; and A. Lebeda (1999). Taxonomy, distribution and biology of the tomato powdery mildew (*Oidium lycopersici*). *Z. Pflanzenkrankh, Pflanzenschutz*, **106**: 140 - 157
- 286 Mihira, T.; T. Takeuchi; H. Fukuda; T. Kawakami; and F. Yamamoto (1999). Physical control of southern root-knot nematode, *Meloidogyne incognita* and brown root rot caused by *Pyrenochaeta lycopersici* by the combination of root restriction and soil solarization on tomato in plastic greenhouse. *Ann. Rep. Kanto -Tosan. Plant Prot. Soc.* **46** : 145 - 148.
- 287 Mikhaeel, F.T., A.N. Estefanus; S.A. El - Shimi F.W. Riad (1997). Mycorrhizal fungus, slomus aggregation and biogas manure as an integrated control of root knot nematode in tomato. *Bull. Fac. Agric. Univ. Cairo*, **48** (4): 737 - 748.
- 288 Miller, J.S.; and D.A. Johnson (2000). Competitive fitness of *Phytophthora infestans* isolates under semiarid field conditions. *Phytopathology*, **90** (6): 220 - 227

- 289-Miller, S. A.; R.C. Rowe; and R.M. Ricdel (unknown). Fusarium and Verticillium wilt of tomato, potato, pepper, and eggplant. (<http://ohioline.osu.edu/hyg.fact/3000/3122.htm/>) 1- 4 P.
- 290-Mishra, M.D.; Rosemarie W. Hammond, R.A. Owens, D.R. Smith, and T.O. Diener (1991). Indian bunch top disease of tomato plants caused by a distinct strain of citrus exocortix viroid J. Gen. Virol. **72**: 1781 - 1785
- 291-Mishriki, J.F. (1978). Calcium distribution in two cultivars differing in fruit susceptibility to blossom end rot. Ph. D. Thesis Fac. Hort., Acad. Agric., Poznan, Poland 122p
- 292-Mizubuti, E.S. G.; and W.E. Fry (1998). Temperature effects on development stages of isolates from three clonal lineages of *Phytophthora infestans*. Phytopathology, **88**: 837 - 844
- 293-Mizubuti, E.S.G.; D.E. Aylor; and W.E. Fry (2000). Survival of *Phytophthora infestans* sporangia exposed to solar radiation. Phytopathology, **90**: 78 - 84
- 294-Momol, T.; J. Pingsheng; K. Pernezny; R. Mc Govern; and S. Olson (2005). Three soilborne tomato diseases caused by *Ralstonia* and *Fusarium* species and their field diagnostics. (<http://edis.ifas.ufl.edu/PP127>) 1 - 7p
- 295-Moore, L.W. (2008). Crown gall disease of nursery crops. (<http://plantdisease.ippc.org/edu/articles.cfm?articleid=8>) 1 - 9p
- 296-Morretti, C.L.; A.G. Calbo; and G.P. Henz (2000). Postharvest physiology and handling, In: Silva, J.B.C. and L.B. Giordano (eds) Processing tomatoes, Brasilia, Embropa/ SC: 136 - 143
- 297-Morretti, C.L.; E. A. Baldwin, S.A. Sargent; and D.H. Huber (2002) Internal bruising alters aroma volatile profiles in tomato fruit tissues. Hort. Science. **37** (2): 378 - 382
- 298-Moura, M.L. R.; and J. Palminha (unknown). A non-chemical method for the control of *Pyrenochaeta lycopersici* of tomato in the north of Portugal (<http://www.actohort.org/books/366/366-39.htm>) 1 p.
- 299-Mumford, R.A.; K. Walsh; N. Boonham (2000). A comparison of molecular methods for the routine detection of viroids. EPPO Bull. **30**, 431 - 436
- 300-Mumford, R.A.; B. Jarvis; and A. Skelton (2004). The first report of potato spindle tuber viroid (PSTVd) in commercial tomatoes in the UK. Plant Pathol. **53**: 242
- 301-Nakhasi, S.; Schlime; and T. Solomos (1991). Storage potential of tomatoes harvested at breaker stage using modified atmosphere storage. J. Food Sci. **56** (1): 55: 59
- 302-Nakhla, M.K.; D.P. Maxwell; R.T. Martinez; M.G. Carvalho; and R.L. Gilbertson (1999). Wide spread occurrence of the eastern mediterranean strain of tomato yellow leaf curl geminivirus in tomatoes in the Dominican Republic. Plant. Dis. **78**: 926.
- 303-Neshev, G. (1993). Powdery mildew (*Oidium sp.*) on tomatoes in Bulgaria Phytoparasitica, **21**: 339 - 343
- 304-Nevo, E.; and M. Coll (2001). Effect of nitrogen fertilization on *Aphis gossypii* (Homoptera: Aphididae) variation in size, color and reproduction. J. Econ. Entomol. **94**: 27 - 32

- 205-Nielsen, G.R. (1997). Flea beetles. ([www. uvm. edu/extension/ publications/el/el 71. htm,](http://www.uvm.edu/extension/publications/el/el_71.htm)) 1 - 3p
- 206-Noling, J.W. (1999). Nematode management in tomatoes, pepper, and eggplant ([http://edis. ifas. ufl. edu/NG. 032.](http://edis.ifas.ufl.edu/NG.032)) 1 - 18p
- 207-Nono-Womdim. R; I.S. Swai; L.K. Mrosso; M.L. Chada; and R.T. Opea (2002). Identification of root knot nematodes species occurring on tomatoes in Tanzania and resistant lines for their control. *Plant Dis.* **86**: 127 - 130
- 208-Oatman, E.R.; and G.R. Platner (1971). Biological control of the tomato fruit worm, cabbage looper and hornworms on processing tomatoes in Southern California, using mass releases of *Trichogramma pretiosum*. *J. Econ. Entomol.* **64**: 501 - 506
- 209-Obradovic, A.; J.B. Jones; M.T. Momol; B. Balogh; S.M. Olson (2004). Management of tomato bacterial spot in the field by foliar applications of bacteriophages and SAR inducers. *Plant Dis.* **88**: 736 - 740
- 210-Olaya, G. and G.S. Abawi (1996). Effect of water potential on mycelial growth and on production and germination of sclerotia of *Macrophomina phaseolina*, *Plant Dis.* **80**: 1347 - 1350
- 211-Olalla, L.; and J.A. Tores (1998). First report of powdery mildew of tomato caused by *Erysiphe sp.* in Spain. *Plant Dis.* **82**: 592.
- 212-Olson, S.M. (2002). Tomato little leaf. ([http://edis. ifas. ufl. edu/ cv 278](http://edis.ifas.ufl.edu/cv278)) 1 - 5p
- 213-Olson, S.M. (2007). Physiological, nutritional and other disorders of tomato fruits. ([http://www. columbia puplications. com/tomato magazine/april 2007/disorder tomatoes htm](http://www.columbiapublications.com/tomato_magazine/april_2007/disorder_tomatoes.htm)) 1 - 4p
- 214-Oliveira, S. H. F.; C. Sinigaglia, and O. Geracia - Junior (2003). Effects of fungicide adjuvants to control early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) in potato and tomato. *Revista de Agricultura Piracicaba*, **78** (1): 57 - 66
- 215-Owens, R.A.; and T.O. Diener (1981). Sensitive and rapid diagnosis of potato spindle tuber viroid disease by nucleic acid hybridization. *Science* **213**: 670 - 672
- 216-Owens, R.A. (2007). Potato spindle tuber viroid; the simplicity paradigm resolved. *Molec. Plant. Pathol.* **8** (5):549
- 217-Ozbun, J.L.; C.E. Boutonnet; S. Sadik; and P.A. Minges (1967). Tomato fruit ripening. I: Effect of potassium nutrition on occurrence of white tissue. *Proc. Amer. Soc. Hort. Sci.* **91**: 566 - 572
- 218-Pandey, K.K.; P.K. Pandey, G. Kalloo; and M.R. Banerjee (2003). Resistance to early blight of tomato with respect to various parameters of disease epidemics. *J. Gen. Plant Pathol.* **69** (6): 364 - 371
- 219-Papavizas, G.C.C. (1977). Some factors affecting survival of sclerotia of *Macrophomina phaseolina* in soil. *Soil Biol. Bioch.*, **9**: 337 - 341
- 220-Papayiannis, L.C.; A.D. Aveglis; N. Ioannou; and N.I. Katis (2006). First report of tomato yellow leaf curl Sardinia virus (TYLCSV) infecting tomato crops in Greece. ([http:// www. bspp. org. uk/ ndr/ july 2006/ 2006- 51. asp](http://www.bspp.org.uk/ndr/july_2006/2006-51.asp)) 1 - 2 P.

- 321-Parrella, M.P. (1987). Biology of *Liriomyza*. Ann. Rev. Entomol. **32**: 201 - 224
- 322-Parsons, C.S.; R.E. Anderson; and R.W. Penney (1970). Storage of mature green tomatoes in controlled atmospheres. J. Amer. Soc. Hort. Sci **95**: 791 - 794
- 323-Partridge, J.F. (2003). Characteristic of plant pathogenic nematodes. (<http://nu-distance.unl.edu/homer/class/5/inded.html>). 1 - 4p
- 324-Patil, M.J.; S.P. Ukey; and B.T. Raut (2001). Evaluation of fungicides and botanicals for the management of early blight (*Alternaria solani*) of tomato. PKV. Research J. **25** (1): 49 - 51
- 325-Pernezny, K.; L.E. Datnoff; T. Mueller; and J. Collins (1996). Losses in fresh market tomato production in Florida due to target spot and bacterial spot and the benefits of protective fungicides. Plant Dis. **80** (5): 559 - 563
- 326-Pernezny, K.; and R.M. Sonada (1998). Powdery mildew of field - grown tomato in Florida. Plant. Dis. **82**: 262
- 327-Pernezny, K.; P. Stoffella; N. Harranek; J. Sanchez; and A. Beany (2004). Bacterial control of target spot using NOTCH. (<http://www.actahort.org/members/showpdf?booknr=695-19>) 1 - 2p
- 328-Persley, D. and J. Thomas (2007). Tomato leaf curl virus (<http://www2.dpi.qld.gov.au/health/4250.html>.) 1 - 6p
- 329-Phillis, J. (1990). The efficiency of nematode resistant tomato cultivars to *Meloidogyne javanica* under greenhouse conditions. Nematol. Mediter. **18**: 209 - 211
- 330-Picha, D.H.; and C.B. Hall (1981). Influence of potassium, cultivar, and season on tomato graywall and blotchy ripening. J. Amer. Soc. Hort. Sci. **106**: 704 - 708
- 331-Picken, A. J.F.; and, M. Griammatt. (1986). The effect of two fruit setting agents on the yield and quality of tomato fruit in glasshouse in winter. J. Hort. Sci. **61**: 243 - 250
- 332-Pittis, J.E.; and R.C. Shattock (1994). Viability, germination and infection potential of oospores of *Phytophthora infestans*. Plant Pathol. **43**: 387 - 396
- 333-Poe, S.L. (1999). Tomato pinworm. (<http://creatures.ifas.edu/veg/tomato/tomato-pinworm.htm>) 1 - 4p
- 334-Pohronezny, K.L.; M.A. Moss; W.Danker; and J. Schenk (1990). Dispersal; and management of *Xanthomonas campestris* pv. *vesicatoria* during thinning of direct seeded tomato. Plant Dis, **74**: 800 - 805
- 335-Polston, J.E.; E. Hiebert, R.J. Mc Govern; P.A. Stansly, and D.J. Schuster (1993). Host range of tomato mottle virus, a new geminivirus infecting tomato in Florida. Plant Dis. **77**: 1181 - 1184
- 336-Polston, J.E.; R.J. Mc Govern, and L.G. Brown (1999). Introduction of tomato yellow leaf curl virus in Florida and implications for the spread of this and other geminiviruses of tomato. Plant Dis. **83**: 983 - 984
- 337-Pracros, P.; J. Renaudin; S. Eveillard; A. Mouras, and M. Hernmould (2006). Tomato flower abnormalities induced by stolbur phytoplasma infection are associated with changes of expression of floral development genes. Molec. Plant Microb, Interac. **19** (1): 62 - 68

- 333-Puchta, H.; T. Herold, K. Verhoeven; A. Roenhorst; K.Ramm, W.Schmidt - Puchta; and H.L. Sanger (1990). A new strain of potato spindle tuber viroid (PSTVd-N) exhibits major sequences differences as compared to all other PSTVd strains sequenced so far. *Plant Mol. Biol.* **15**: 509 - 511
- 339-Pujos, A.; and M. Morard (1997). Effects of potassium deficiency on tomato growth and mineral nutrition at the early production stage. *Plant and Soil.* **189**: 189 - 196
- 340-Punja, Z.K. (1985). The biology, ecology and control of *Sclerotium rolfsii*. *Ann. Rev. Phytopath.* **23**: 97 - 127
- 341.-Purdey, L.H. (1979). *Sclerotinia sclerotiorum*: history, diseases and symptomatology, host range, geographic distribution, and impact. *Phytopathology.* **69**: 875 - 880
- 342-Purkayastha, S.; B. Kaur; N.Dilbahi; and A. Chaudkury. (2006). Characterization of *Macrophomina phaseolina*, the charcoal rot pathogen of cluster bean, using conventional techniques and PCR - based molecular markers. *Plant Pathol.* **55**: 106 - 116
- 343.-Raccah, B. (2000). Plant virus transmission by insects. In: *Encyclopedia of Life Science.* John Wiley & Sons, LTD. Chichester
- 344-Rao, M.H. (1994). Growth, yield and quality of tomato, carrot and cauliflower as influenced by levels and sources of potassium. *J. Potassium Res.* **10** (4): 402 - 406
- 345.-Rebois, R.V. (1973). Effect of soil temperature on infectivity and development of *Rotylenchulus reniformis* on resistant and susceptible soybeans. *J. Nematol.* **5**: 10 - 13
- 346-Reynolds, L.B.; J.W. Potter and B.R. Ball - Coelho (2000). Crop rotation with *Tagetes sp.* is an alternative to chemical fumigation for control of root lesion nematodes. *Agronomy J.* **92**: 957 - 966
- 347-Rich, J.R.; and G.W. Bird (1973). Inhibition of *Rotylenchulus reniformis* penetration of tomato and cotton roots with foliar applications of oxamyl. *J. Nematol.* **5**: 221 - 224
- 348-Rich, C.M. (1978). The tomato. *Sci. American*, **239**(2): 76 - 87
- 349-Ristaino, J.B.; K.B. Perry, and R.D. Lumsden (2001). Effect of solarization and *Gliocladium virens* on sclerotia of *Sclerotium rolfsii*, soil microbiota, and the incidence of southern blights of tomato. *Phytopathology*, **91**: 1117 - 1124
- 350-Ritenour, M.A.; S.A. Sargent; and J.A. Bartz (2002). Chlorine use in produce packing lines. (<http://edis.ifas.ufl.edu/pdffiles/CH/CH16000.pdf>) 1 - 4p
- 351-Roberts, P.D.; R.J. Mc Govern; and L.E. Datnolf (2001). Fusarium crown and root rot of tomato in Florida. (<http://edis.ifas.ufl.edu/PG.082>). 1 - 4p
- 352-Robinson, J.E.; K.E. Browne; and W.G. Burton (1975). Storage characteristics of some vegetables and soft fruits. *Ann. Appl. Biol.* **81**: 399 - 408
- 353-Robinson. A.F; C.M. Heald; S.L. Flanagan; W.H. Thames; and J. Amador (1987). Geographical distribution of *Rotylenchulus reniformis*, *Meloidogyne incognita* and *Tylenchulus semipetrans* in the lower Rio Grand Valley as related to soil texture and land use. *Ann Appl. Nematol.* **1**. 20 - 25.

- 354-Robinson, A.F., R.N. Inserra, E.P. Caswell - Chen; N. Volvas; and A. Troccoli (1997). *Rotylenchulus* species: Identification, distribution, host ranges, and crop plant resistance. *Nematropica*, **27**: 127 - 180
- 355-Robson, M.C. (2000). Tomato problems can be avoided (<http://ring.wsu.edu/column/17-16-00.htm>). 1 - 2p
- 356-Rocheleau, L.; and M. Pelchat (2006). The subviral RNA data base: a toolbox for virologists, the hepatitis delta virus and satellite RNAs. *Research BMC Microbiol* **6**: 24
- 357-Rojas - Martinez, R.I.; E. Zavaleta - Mejia; and R.O. Gomez. (1999). Effect of association of marigold - tomato on the injury caused by *Alternaria solani* in tomato. *Fitopatologica*. **4** (2): 83 -89
- 358-Ryder, M.H.; and D.A. Jones (1991). Biological control of crown gall using *Agrobacterium* strains. K 84 and K1026. *Australian J. Plant Physiol.* **18**: 571 - 579
- 359-Sabaratnam, S. (2008). Tomato chlorotic dwarf viroid (TCDVd) (<http://www.agf.gov.ca/cropprot/tcdvd.htm>) 1 - 4p
- 360-Salazar, L.F.; M. Querci; I. Bartolini; and V.Lazarte (1995). Aphid transmission of potato spindle tuber viroid assisted by potato leaf roll virus. *Fitopatologica* **30**: 50 - 58
- 361-Salaguero Navas, V.E; J. E. Funderburk; S.M. Olson; and R.J. Beshear (1991). Damage to tomato fruit by the western flower thrips. *J. Entomol. Sci.* **26** (4): 436 - 442
- 362-Sano, T.; R. W. Hammond; and R.A. Owens (2003). Biotechnological approaches for controlling viroid diseases. Pages 343 - 349 in: *Viroids*. A. Hadidi; R. Flores; J.W. Ranby; and J.S. Semancik, eds. CSIRO Publishing, Collingwood, Australia
- 363-Sanogo, S.; and S.P. Pennypacker (1997). Factors affecting sporangial and mycelial germination of sclerotia of *Colletotrichum coccodes*. *Plant Dis.* **81**: 333 - 336
- 364-Sargent, S.A.; J.K. Brecht; and J.J. Zollner (1992). Sensitivity of tomatoes at mature - green and breaker ripeness stages to internal bruising. *J. Amer Soc.Hort. Sci.* **117** (1): 119 - 123
- 365-Sargent, S.A.; and J.J. Van Sickle (1996). Premium - quality tomato program for Florida. *Proc. Florida tomato Inst., C.S. Vaverna (ed) Publ. PRD 108. Univ. Florida Inst. Food & Agric. Sci Gainesville.* pp 8 - 9
- 366-Sawant, G.C.; P.V. Desai; and R.G. Padhiar (1999). Effect of different spacing, fertilizer dose, use of *Trichoderma viride* and different fungicidal formulations on the occurrence of early blight disease of tomatoes. *Indian J. Environ. Toxic.* **9** (2): 84 - 87
- 367-Schwartz, H.F.; and F.H. Gent (2007). Eggplant, pepper and tomato white mold ([http://highplainsipm.org/Hp_IPM_search/Does/White mold - eggplant pepper tomato. tm](http://highplainsipm.org/Hp_IPM_search/Does/White_mold_eggplant_pepper_tomato.htm)) 1 - 3p
- 368-Sclar, C. (2000). An overview of western flower thrips management. *Rutgers Coop Ext. Newsletter.* **10** (5): 2 - 3
- 369-Scemuller, E.; C. Marcone; U. Louer; A. Ragozzino, and M. Goschi (1998). Current status of molecular classification of the phytoplasmas. *J. Plant Pathol.* **80**: 3 - 26

- 370-Sheikh, A.; and A. Ghaffar (1979). Relation of sclerotial inoculum density and soil moisture to infection of field crops by *Machrophomina phaseolina*. Pakistan J.Bot. 11: 185 - 189
- 371-Shishkoff, N.; and R.N. Campbell (1990). Survival of *Pyrenochaeta lycopersici* and the influence of temperature and cultivar resistance on the development of corky root of tomato Plant. Plant Dis. 74: 889 - 894
- 372-Shizue, S.; K. Eiko; I. Koji; T. Koji; D. Hirotsuka; and N. Hideo (2004). Chemical control of *Corynespora* target spot of tomato caused by benzimidazol resistant strain of *Corynespora cassicola*. (http://sciencelinks.in/east/article/2004_24/000020042404_A_0816498nhn) 1p
- 373-Siddiqui, I.A.; and S.S. Shaikat (2002). Mixtures of plant disease suppressive bacteria enhance biological control of multiple tomato pathogens. Biol. Fertil. Soil. 36 (4): 260 - 268
- 374-Sijmons, P.G.; H.J. Atkinson; and U. Wyss (1994). Parasitic strategies of root nematodes and associated host cell responses. Ann, Rev. Phytopath. 32: 235 - 259
- 375-Sikora, E.J. (2007). Virus diseases of tomato (<http://www.aces.edu/pubs/docs/A/ANR-08361>) 1 - 6p
- 376-Sinisterra, X.H.; C.L. McKenzie; W.B. Hunter; R.G. Shatters (2005). Transcript expression of Begomovirus in the whitefly vector *Bemisia tabaci* (Hemiptera: Aleyrodidae). J. Gen. Virol., 86: 1525 - 1532
- 377-Simons, J.N.; and D.M. Coe (1958). Transmission of pseudo - curly top virus in Florida by a treehopper. Virology, 6: 43 - 48
- 378-Simons, J.N. (1962b). Life history and behavioral studies on *Micrutalis malleifera*, a vector of pseudo- curly top virus. J. Econ Entomol. 55: 363 - 365
- 379-Singh, R.P. (1970). Seed transmission of potato spindle tuber virus in tomato and potato. American Potato. J. 47: 225 - 227
- 380 Singh, R.P.; X. Nie; and M. Singh (1999). Tomato chlorotic dwarf viroid: an evolutionary link in the origin of pospiviroids. J. Gen. Virol. 60: 2323 - 2328
- 381 Singh, R.P.; M.K. Rana; and S. Singh (2001). Effect of host nutrition on early blight of tomato. J. Mycol. Plant Pathol. 31 (2): 248 - 250.
- 382- Singh. R.P.; and A. D. Oilworth (2006). Detection of citrus exocortis viroid. Iresine viroid and tomato chlorotic dwarf viroid in new ornamental host plants in India. Plant Dis. 90: 1457.
- 383-Skoric, D.; M. Conerly; J.A. Szychowski; and J.S. Semancik (2001). Citrus exocortis viroid - induced symptom modification as a response to a host specific temperature - sensitive reaction. Virology 280: 115 - 123.
- 384 Smart, G.C.; and K.B. Nguyen (1991). Sting and awlnematodes. Pp. 627- 668. In Nickle, W.R. (ed.), Man. Agric. Nematol. Marcel Dekker Inc., Ny.
- 385 Smart, C.D.; M.R. Willman; H. Mayton; E.S.G. Mizubuti; R.W. Sandrock; A.E. Muldoon; and W.E. Fry (1998). Self fertility in two clonal lineages of *Phytophthora infestans*. Fung. Gen. Biol. 25: 134 - 142.
- 386-Smith. V.L.; S. M. Douglas; and J.A. La Mondia (1997). First report of powdery mildew of tomato caused by *Erysiphe* sp. in Connecticut. Plant Dis. 81: 229 - 263.

- 387-Squitier, J.M. (2007). Southern green stink bug, *Nezara viridula* (Linnaeus). (<http://entemdept.ifas.ufl.edu/creatures/veg/bean/southern-green-stink-bug.htm>) 1 - 4 p
- 388-Stapleton, J.J.; and J.E. De Vay (1986). Soil solarization a non-chemical approach to disease and pest management. *Crop Protection* 5: 190 - 198.
- 389-Steadman, J.R. (1979). Control of plant diseases caused by *Sclerotinia* species. *Phytopathology*, 69: 904 - 907.
- 390-Stone, T.B.; and S.R. Sims (1993). Geographic susceptibility of *Heliothis virescens* and *Helicoverpa zea* (Lepidoptera: Noctuidae) to *Bacillus thuringiensis*. *J. Econ. Entomol.* 86: 989 - 994.
- 391-Surdiman, A.N.; and J.M. Webster (1995). Effect of ammonium ions on egg hatching and second stage juveniles of *Meloidogyne incognita* in axenic tomato root culture. *J. Nematol.* 27: 346 - 352.
- 392-Susmu, T.; T. Koji; S. Yoko; and T. Yoshihiro (2005). Control of the vegetable leafminer, *Liriomyza sativa* Blanchard, in tomato greenhouse by using yellow sticky roller traps. *Ann. Rep. Kansai Plant Prot. Soc.* 47: 133 - 135.
- 393-Swift, C.E. (1996). *Cuscuta* and *Grammica* species - Dodder: A plant parasite. (<http://www.colostate.edu/Depts/coopExt/dodder.html>) 1- 4p.
- 394-Symons, R.H. (1991). Intriguing viroids and virusoids: What is their information content and how did they evolve. *Mol. Plant Microb. Interact.* 4 (2): 111 - 121.
- 395-Tantius, P.H.; A.M. Fyfe; D.S Shaw; and R.C. Shattock (1986). Occurrence of the A2 mating type and self-fertile isolates of *Phytophthora infestans* in England and Wales. *Plant Pathol.* 35: 578 - 581.
- 396-Tenllado, F.; and J. Bol (2000). Genetic dissection of the multiple functions of alfalfa mosaic virus coat protein in viral RNA replication, encapsidation, and movement. *Virology*, 268: 29 - 40.
- 397-Thomma, B.P.H, P.H. Van Esse; P.W. Crous; and P.J. De Wit (2005). *Cladosporium fulvum* (syn. *Passalora fulva*) a highly specialized plant pathogen as a model for functional studies on plant pathogenic *Mycosphaerellaceae*. *Mol. Plant. Pathol.* 6 (4): 379 - 393.
- 398-Tisserat, N. (2006). Leaf spot of tomato. (<http://www.oznet.kstat.edu/path-ext/act sheets/tomato/early 20% blight 20% Septoria 1-3P>).
- 399-Todd, J.W. (1989). Ecology and behavior of *Nezara viridula*. *Ann. Rev. Entomol.* 34: 273 - 292.
- 400-Tofoli, J.G.; R.J. Domingues; O. Garcia - Junior; and C. kurosawa (2003). Tomato early blight control by fungicides and its effects on yield. *Summa Phytopathologica*, 29 (3): 225 - 235
- 401-Tsror, L. and B.J. Bieche (1999). Biological control of early blight in tomatoes. *Acta Horticulturalae*, 487: 271 - 273.
- 402-USDA (1966). US. Standards for Grades of Greenhouse Tomatoes USDA. Agric. Mktg. Serv. Washington, D.C. (<http://www.ams.usda.gov/standards/stanfrfv.htm>) 1P.
- 403-USDA (1991). U.S. Standards for Grades of Fresh Tomatoes. USDA. Agric. Mktg. Serv., Washington, D.C. (www.ba.ars.usda.gov/hb66/138tomato.pdf).

- 414- Vakalounakis, D.J. and A. Papadakis (1992). Occurrence of a new powdery mildew of greenhouse tomato in Greece, caused by *Erysiphe* sp. *Plant Pathol.* **41**: 372 - 373.
- 415- Vattanatumgum, A.; and W. Ruchtapakoranchai (1979). Principal insects which attack tomato in the tropics and their control. In «Proc. 1st Inter. Symp. Tropical Tomato» pp 132 - 135, Shanhua, Taiwan.
- 416- Vegt, W. and H. Buchenauer (1997). Enhancement of biological control by combination of antagonistic fluorescent *Pseudomonas* strains and resistance inducers against damping-off and powdery mildew in cucumber. *Z. Pflanzenkrankh. Pflanzenschutz.* **104** (3): 272 - 280.
- 417- Veloukas, I.; J.A. Bardas; G.S. Karaoglou; and K. Tzavella Klonari (2007). Management of tomato leaf mould caused by *Cladosporium fulvum* with trifloystrobin. *Crop Protect* **26** (6): 845 - 851.
- 418- Verhoeven, J.Th. J.; C.C.C. Jonson; T.M. Willemer; L.F.F. Kox; R.A. Owens; and J.W. Roenhorst (2004). Natural infections of tomato by citrus exocortis viroid, columnea latent viroid, potato spindle tuber viroid and tomato chlorotic dwarf viroid. *Eur. J. Plant Pathol.* **110**: 823 - 831.
- 419- Verhoeven, C.; C.C. Jonson; and J.W. Roenhorst (2006). First report of tomato apical stunt viroid in tomato in Tunisia. *Plant Dis.* **90**: 528.
- 420- Walker, J.C. (1950). *Plant Pathology*. Mc Graw - Hill Book Company. New York.
- 421- Wallace, H.R. (1968). The influence of soil moisture on survival and hatch of *Meloidogyne javanica*. *Nematologica.* **14**: 231 - 242.
- 422- Watterson, J.C. (1986). Disease. In J.G. Atherton and J. Rudich (Eds) «The Tomato Crop» pp. 443 - 484. Chapman and Hall. London.
- 423- Weingartner, D.P.; J.R. Shumaker; and G.C. Smart (1983). Why soil fumigation fails to control potato corky ringspot disease in Florida. *Plant Dis.* **67**: 130 - 134.
- 424- Whipps, J.M.; S.P. Budge; and J.S. Fenlon (1998). Characteristics and host range of tomato powdery mildew. *Plant Pathol.* **47**: 36 - 48.
- 425- White, C. (1999). *Handbook of chlorination and alternative disinfectants*. 4th Edition. John Wiley & Sons, Inc. New York, NY.
- 426- Whittaker, R.H. (1968). New concepts of kingdoms of organisms. *Science.* **163**: 150 - 160.
- 427- Wilcox, G.E., J.E. Hoff; and C.M. Jones (1973). Ammonium reduction of calcium and magnesium content of tomato and sweet corn leaf tissue, and influence on incidence of blossom end rot of tomato fruit. *J. Amer. Soc. Hort. Sci.*, **98**: 86 - 89.
- 428- Wills, R.B.H.; B. Mc Glasson; D. Graham; and D. Joyce (1998). *Postharvest: An introduction to the physiology and handling of fruits, vegetables and ornamentals*. Univ. New South Wales Press. 262pp.
- 429- Wilson, M.; H.L. Campbell; J.B. Jones; and D.A. Cuppels (2002). Biological control of bacterial speck of tomato under field conditions at several locations in North America. *Phytopathology*, **92** (12): 1284 - 1292.
- 430- Woese, C.; and G. Fox (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. USA.*, **74** (11): 5088 - 5090.

- 421-Woese, C.R.; W.E. Balch; L.J. Magrum, G.E. Fox; and R.S. Wolfe (1977). An ancient divergence among the bacteria. *J. Molec. Evolut.*, **9**: 305 - 311.
- 422-Woese, C.; O. Kandler; M. Wheelis (1990). Towards a natural system of organisms: proposal for the domains: Archaea, Bacteria and Eucarya. *Proc. Natl. Acad. Sci. USA.*, **87** (12): 4576 - 4579.
- 423-Wokocha, R.C. (1990). Integrated control of *Sclerotium rolfsii* infection on tomato in the Nigerean Savana: Effect of *Trichoderma viride* and some fungicides. *Crop Protect.*, **9**: 231 - 234.
- 424-Workneh, F.; A.H.C. Van Bruggen; L.E. Drinkwater; and C. Shenan (1993). Varieties associated with corky root and *Phytophthora* root rot of tomatoes in organic and conventional farms. *Phytopathology*, **83**: 581 - 589.
- 425-Yarden, D.; T. Ben - Yophet; J. Katan; and N. Aharonson (1986). Fungicidal control of *Sclerotinia sclerotiorum* in soil with a combination of benomyl and thiram. *Plant Dis.* **70**: 738- 742.
- 426-Yoshim, Y.; and H.B. Scholthof (2005). Tomato bushy stunt virus: a resilient model system to study virus- plant interactions. *Mol. Plant Pathol.*, **6** (5): 491 - 502.
- 427-Zehnder, G.W.; and J.T. Trumble (1984). Spatial and diel activity of *Liriomyza* species (Diptera: Agromyzidae) in fresh market tomatoes. *Environ. Entom.*, **13**: 1411 - 1416.
- 428-Zhu, Y.; L. Green; Y.M. Woo; R. Owens; and B. Ding (2001). Cellular basis of potato spindle tuber viroid systemic movement. *Virology*, **279** (1): 69 - 77.
- 429-Zupan, J.; T.R. Muth; O. Draper; and P. Zambryski (2001). The transfer of DNA from *Agrobacterium tumefaciens* into plants: a feast of fundamental insights. *Plant J.*, **23**: 1 - 28.



صور ملونة



الفصل الرابع

Tomato Late Blight

Fig (1)

Life cycle of *P. infestans*
(a) Sporangiohores; (b) sporangia;
(c) sporangial contents dividing up
to form zoospores; (d) zoospores; (e)
germinating zoospores




Fig (2)

Late blight on tomato leaf



Fig (3)

Late blight lesions on susceptible tomato
stem





Fig (4)

Tomato late blight on green fruits



Fig (5)

Tomato late blight on red fruits

Tomato Buckeye




Fig (6)

Phytophthora parasitica, a common buckeye rot fungus: sporangiophores and sporangia in various stages of forming zoospores (drawing by Lenore Gray)



Fig (7)

Fruit lesions with concentric ring pattern typical of buckeye rot (Photo by P. Warren)



Tomato Early Blight and Collar Rot (caused by *A. solani*)

Fig (8)
Alternaria solani spores




Fig (9)
Tomato collar rot on seedling stem



Fig (10)
Tomato Early Blight
Stem lesions caused by *A. solani*






Fig (11)
Tomato early blight
Target- ring lesions leaf



Fig (12)
Tomato early blight on fruit and leaves

Alternaria Stem Canker (*A. alternata*)



Fig (13)
Conidia and conidiophore of
A. alternata



Fig (14)
Alternaria stem canker



Alternaria tenuis

Fig (15)
Conidia and conidiophores of *A. tenuis*



Septoria Leaf Spot

Fig (16)
Pycnidia of *Septoria lycopersici*




Fig (17)
Septoria lesions have tan or whitish centers





Black Leaf Mold

Pseudocercospora fuligena conidiophores and conidium



Fig (18)
Fascicle of divergent
conidiophores



Fig (19)
Solitary cylindrical-
obclavate conidium

Symptoms of Black Leaf Mold on Tomato Leaves



Fig (20)

Black sooty patches develop on both upper and lower leaf surfaces (left)
The soot-covered leaves wilt, dry, and usually remain hanging on the vine (right)



Tomato Leaf Mold

Fig (21)

Fulvia fulva the tomato leaf mold fungus as it might appear under a high- power microscope: (a) a fascicle of conidiophores emerging from the leaf surface; (b) conidia (drawing by Lenore Gray)




Fig (22)

Tomato leaf mold symptoms on upper leaf surface



Fig (23)

Tomato leaf mold symptoms on lower leaf surface





Tomato Leaf Mold



Fig (24)
Tomato leaf mold symptoms on fruit and stem

Didymella Stem Rot



Fig (25)
Black stem cankers on seedling



Fig (26)
Large black stem canker



Tomato Target Spot



Fig (27)

Corynespora cassiicola; conidiophore with linked cylindrical cells




Fig (28)

C. cassiicola; conidium with pseudo-septa



Fig (29)

Symptom of target spot on tomato leaves



Fig (30)

Symptom of target spot on young fruit

Fig (31)

Target spot symptom on ripening fruit





Grey Leaf Spot



Fig (32)

Conidium spore of *Stemphylium*




Fig (33)

Stemphylium solani, *S. lycopersici*
symptoms on tomato leaf

Tomato Phoma Rot



Fig (34)

Tomato phoma rot on leaf



Fig (35)

Tomato phoma rot on fruit



Tomato Anthracnose



Fig (36)
Spores of *C. coccodes*



Fig (37)
Acervulus: conidia and conidiophores



Fig (38)
Sunken lesions, slightly paler than the healthy tissue. black fruiting bodies are visible on this closed lesion



Tomato Fusarium wilt



Fig (39)

Fusarium wilt of tomato. Note yellowing and death of leaves on one side of the stem

Verticillium and Fusarium Vascular Discoloration




Fig (40)

Dark brown vascular discoloration in tomato caused by Fusarium wilt (lower). Verticillium wilt (upper) causes a lighter tan discoloration of the vascular tissue



Tomato Verticillium Wilt

Fig (41)

V. Albo atrum conidiophore




Fig (42)

V. dahliae: conidiophore and spores




Fig (43)

Typical V- shaped lesions on tomato leaves associated with Verticillium wilt





Tomato Crown and Root Rot



Fig (44)
Canker on stem at soil line



Fig (45)
Internal discoloration of the crown and
root rot. (Note missing taproot)

White Mold on Tomato



Fig (46)
Black sclerotia in stem of infected
tomato vines



Fig (47)
White mold on tomato fruit



Tomato Grey Mold

Fig (48)

Conidiphore and conidia of *B. cinerea*



Fig (49)

Grey mould on flower



Fig (50)

Grey mould on tomato fruit



Fig (51)

Early symptoms of grey mould on pruning wound





Tomato Powdery Mildew




Fig (52)

Conidia (A) and conidiophore (B) of
Oidium neolycopersici



Fig (53)

Symptoms of *Oidium taurica* consist
of chlorotic spots on the upper leaf
surface and profuse fungal sporulation
on the lower surface (left photo). For
Oidium neolycopersici, Powdery white
colonies appear on the upper surface
(right photo)



Tomato Corky Root rot




Fig (54)

Tomato corky root rot



Tomato Southern Blight

Fig (55)

White mycelial growth and sclerotia
near the soil surface



Tomato Damping off

Fig (56)

Affected plants occur in batches in
nursery beds



Fig (57)

Seedlings affected by damping off





Sour Rot




Fig (58)

Anthroconidium development in
Geotrichum candidum



Fig (59)

Tomato fruit with *Geotrichum* sour- rot

Rhizopus Rot on Tomato Fruit



Fig (60)

A carton of tomatoes with nested
Rhizopus rot (and some secondary
fungi)

Black Mold on Tomato Fruit




Fig (61)

A fruit with a fingernail wound (arrow)
that later developed into black mold



Tomato Soil Rot

Fig (62)
Soil rot on tomato fruit caused by
R. solani



Pythium Rot

Fig (63)
Pythium rot



Fusarium rot

Fig (64)
Fusarium originating from the stem-
scar





الفصل الخامس

Tomato Bacterial Wilt



Sudden & permanent wilt occur



Fig (1)

Brown discoloration of the vascular system



An ooze will flow from a cut infected stem (right tube)

Tomato Bacterial Soft Rot



Fig (2)

Vines wilt and die; the inner stem becomes brown and slimy



Fig (3)
Bacterial soft rot on tomato fruit



Fig (4)
Sliced tomato showed infected tissue with bacterial soft rot through the blossom end



Tomato Pith Necrosis

Fig (5)
The stem pith of tomato plant affected with pith necrosis turns dark brown and develops hollow cavities





Tomato Bacterial Spot




Fig (6)
Initial symptoms



Fig (7)
Lesions may coalesce, causing blighted areas on leaves



Fig (8)
Immature fruit show brown, slightly sunken, scabby spots. Lesions on stems are elliptical in shape



Tomato Bacterial Speck


Fig (9)

Leaves from a greenhouse- grown tomato seedling infected with bacterial speck



Fig (10)

Fruit infected with bacterial speck



Bacterial Canker

Fig (11)

Bacterial canker symptoms on tomato leaflet, showing yellow border between live and dead tissue



Fig (12)

Bacterial canker symptoms on tomato fruit showing bird's eye. On leaflet, showing distinctive upward curling of leaf edges





Tomato Crown Gall



Fig (13)
Tomato infected with crown gall



الفصل السادس

Tomato Aster Yellows

Fig (1)
Infected and healthy tomato plants



Tomato Big Bud

Fig (2)
Enlarged buds and growth distortion
is characteristic of tomato big- bud
phytoplasma





Tomato Stolbur Phytoplasma



Absence of petals stamens and carpels



Mature flower

Fig (3)



Sepal hypertrophy green petals and stamens



Leaf structure of sepals, absence of petals



الفصل السابع

Tobacco Mosaic Virus




Fig (1)

1- Genomic RNA 2 - Capsomer
3 - Capsid



Fig (2)

Symptoms of tobacco mosaic virus on tomato

A- Symptoms on leaves

B- Gray wall symptoms

C- Necrotic spots on fruit



Tomato Mosaic Virus



Fig (3)

Leaf symptoms of tomato mosaic virus



Fig (4)

Fruit symptoms of tomato mosaic virus

Cucumber Mosaic Virus



Fig (5)

Healthy leaf (left) and infected leaf (right)




Fig (6)

Shoestring, tendril-like foliage



Tomato Yellow Leaf Curl Virus


Fig (7)

Infected plant (left) is yellowing and becoming stunted as compared to healthy plant (right)



Fig (8)

Leaf curling and erect growth



Tomato Yellow Leaf Curl Sardinia Virus

Fig (9)

Yellow leaf curl disease on tomato caused by yellow leaf curl Sardinia virus





Tomato Leaf Curl Virus



Fig (10)

Leaves curling and distortion



Fig (11)

(A) 21 days- leaves curling and early stages of interveinal yellowing



(B) 35 day- marked leaf curling and interveinal yellowing



(C) 56 day- plant distortion and loss of leaf development



Potato Virus Y



Fig (12)

Foliage shows mild roughness, faint mottling and slight distortion (upper photo). Later, the foliage may curl downward, giving the plant a drooping appearance (under photo)



Tobacco Etch Virus

Fig (13)

Leaves show mottling, crinkling, distortion and downward curling





Tomato Spotted Wilt Virus



Fig (14)

Purple flecking of young leaves caused by tomato spotted wilt




Fig (15)

Fruit symptoms of tomato spotted wilt

Tomato Alfalfa Mosaic Virus



Fig (16)

Foliage damaged by alfalfa mosaic virus



Fig (17)

Alfalfa mosaic virus symptoms on tomato fruits



Tomato Curly Top Virus

Fig (18)
Foliage damaged by beet curly top virus



Tomato Bushy Stunt Virus

Fig (19)
Upper leaves of plants infected by tomato bushy stunt virus are yellow and curled



Fig (20)
Tomato bushy stunt virus on tomato fruits





Tomato Mottle Virus



Fig (21)
Tomato mottle virus

Pseudo Curly Top Virus on Tomato Plant



Fig (22)
Infected tomato with healthy
comparison




Fig (23)
Top of plant infected with pseudo curly
top virus



Tomato Yellow Top Virus

Fig (24)
Drawed tomato plant





الفصل الثامن

Potato Spindle Tuber Viroid (PSTVd)



Fig (1)

Tomato infected with PSTVd showing purpling and chlorosis, down-curling, stunting and distortion

Tomato Apical Stunt Viroid (TASVd)



Fig (2)

Symptoms of tomato apical stunt viroid on tomato leaves



Tomato Chlorotic Dwarf Viroid (TCDVd)



Fig (3)
Greenhouse tomato plants infected
with TCDVd



Citrus Exocortis Viroid on Tomato Plants (CEVd)

Fig (4)
Tomato plants infected by CEVd





Tomato Bunchy Top Viroid (TBTVD)



Fig (5)
Symptoms of TBTVD on tomato plant

Tomato Planta Macho Viroid (TPMVD)



Fig (6)
Symptoms of tomato planta macho viroid on tomato plant



الفصل التاسع

Plant Pathogenic Nematodes




Fig (1)

Female and male nematodes

Root- Knot Nematode

Fig (2)

Adult female of the root-knot nematode (*Meloidogyne sp.*) with attached egg mass



Fig (3)

Adult male of root-knot nematode





Root-Knot Nematode




Fig (4)

Root system infected with *M. incognita*

Reniform Nematode




Fig (5)

Male and young female of reniform nematode. stages typically found in soil




Fig (6)

Life stages of reniform nematode. ranging from left to right is juvenile. young female with swollen body. and mature female in kidney shape



Sting Nematode

Fig (7)
The adult female of sting nematode





Fig (8)
Greatly enlarged adult female
sting nematode showing slender,
elongated stylet (center) used for
reaching deep inside root tissue





Stubby Root Nematode




Fig (9)
Stubby root nematode



Fig (10)
Onchio style of stubby root nematode

Root Lesion Nematode




Fig (11)
Tomato root lesion nematode



الفصل العاشر

Whitefly

Fig (1)
Nymphs stage of whitefly



Fig (2)
Adult stage of whitefly



Aphids

Fig (3)
Green peach aphid colony





Thrips




Fig (4)
Western flower thrips

Cabbage Looper



Fig (5)
Cabbage looper moth



Fig (6)
Cabbage looper larva



Tomato Psyllid

Fig (7)
Adult stage of tomato psyllid




Fig (8)
Tomato psyllid numphs on tomato leaf



Fig (9)
Psyllid sugar on tomato leaves





Tomato Pinworm




Fig (10)

Tomato Pinworm. A. Adult. B. Eggs and new larva. C. Larva. D. Pupa. E. Early damage. F. Later damage



Fig (11)

Larva of tomato pinworm

Flea Beetles



Fig (12)

Potato flea beetle



Fig (13)

Tobacco flea beetle



Fig (14)
Corn flea beetle



Fig (15)
Eggplant flea beetle



Tomato Fruitworms

Fig (16)
Life cycle of tomato fruitworm





Fig (17)
Life cycle of tobacco budworm





Tomato and Tobacco Hornworms



Fig (18)
Adult tomato hornworm



Fig (19)
Adult tobacco hornworm




Fig (20)
Larva of tomato hornworm



Fig (21)
Larva of tobacco hornworm



Beet Leafhopper

Fig (22)
Adult beet leafhopper



Tomato Cutworms

Fig (23A)
Larva of black cutworm




Fig (23B)
Moth of black cutworm





Fig (24A)
Larva of granulate cutworm



Fig (24B)
Moth of granulate cutworm



Fig (25A)
Larva of variegated cutworm



Fig (25B)
Moth of variegated cutworm



Tomato Cutworms Life Cycle

Fig (26)
Black cutworm




Fig (27)
Granulate cutworm





Fig (28)
Variegated cutworm





Cotton Leafworm



Fig (29A)

Larva of cotton leafworm



Fig (29B)

Moth of cotton leafworm

Potato Tuberworm



Fig (30A)

Larva of potato tuberworm



Fig (30B)

Moth of potato tuberworm



Vegetable Leafminer

Fig (31A)
Adult *L. sativae* (vegetable
leafminer)



Fig (31B)
Larva of *L. sativae*



Fig (31C)
Leafminer damage on greenhouse
tomato leaf



Mole Cricket

Fig (32)
Grylotalpa grylotalpa





Tomato Stink Bugs



Fig (33)

Adults of tomato stink bugs



Fig (34)

Tomato fruit infected with stink bugs

Spider Mite



Fig (35A)

Two spotted spider mite




Fig (35B)

Red spider mite



Tomato Borers

Fig (36)
Tuta absoluta moth



Fig (37)
Tuta absoluta larva



Fig (38)
Tuta absoluta infestation on fruits





الفصل الحادى عشر

Effect of Pesticides on Tomato Leaf



Fig (1)

Malathion damage on tomato leaf

Tomato Physiological Diseases



Fig (2)

Tomato fruits infected with BER
Tomato Blossom End Rot (BER)



Tomato Catface

Catfaced fruit with hole into fruit



Fig (3)

Fruit showing catfacing on blossom end



Tomato Fruit Cracking

Fig (4A)

Radial cracking of tomato fruit




Fig (4B)

Concentric cracking of tomato fruit





Physiological Tomato Leaf Roll



Fig (5)

A) Severe physiological leaf roll symptoms on a tomato plant. B) Tomato plants with physiological leaf roll on the older (lower) leaves with normal new (top) growth that developed after air temperatures cooled. C) Some tomato cultivars are less susceptible to physiological leaf roll than others

Tomato Sun Scald



Fig (6)

Lethal sun scald. note sunken area



Puffiness



A) Fruit severely affected by puffiness. note larg open areas



B) Note absence of seed in gel area caused by puffiness

Fig (7)

Yellow and Green Shoulders



A
Yellow shoulder



B
Green shoulder

Fig (8)



Internal White Tissue



Fig (9)

Internal white tissues

Tomato Fruit Pox




Fig (10)

White spots on tomato fruit



Tomato Gold Fleck



A

Mottle spots on tomato fruit



B

Circular ring spot on tomato fruit

Fig (11)

Irregular Ripening

Fig (12)

Irregular ripening on tomato fruits



Gray Wall

Fig (13)

Gray wall on tomato fruits





Cloudy Spots



Fig (14)

Cloudy spot on tomato fruit

Rain Check



Fig (15)

Rain check on tomato fruit

Zippering



Fig (16)

Zipper strip spots on tomato fruit

Dimpling



Fig (17)

Oviposition dimples persisting on ripe fruit



Symptoms of Elements Deficiency



Fig (18)

Nitrogen deficiency




Fig (19A)

Phosphorus deficiency



Fig (19B)



Fig (20)
Potassium deficiency



Fig (21)
Calcium deficiency




Fig (22)
Magnesium deficiency



Fig (23)
Sulfur deficiency




Fig (24)
Iron deficiency



Fig (25)
Copper deficiency




Fig (26)
Boron deficiency



Fig (27)
Zinc deficiency



Fig (28)
Manganese deficiency



Fig (29)
Molybdenum deficiency



Fig (30)
Chloride deficiency



الفصل الثاني عشر

Broomrape

Fig (1)
Orobanche sp



Dodder

Fig (2)
Cuscuta sp





Herbicide Damage



Fig (3)
Herbicide damage on tomato plants