

RECOMMENDATIONS

RECOMMENDATIONS

From the current study, it is recommended that:

1. Oral dexmedetomidine premedication can be used successfully for pediatric sedation.
2. Proper dose, formulation and timing of oral dexmedetomidine intake make it more effective with fewer side effects.
3. Oral dexmedetomidine alone can be not enough to preform painfull procedures.
4. Future studies should emphasize on the antrograde amnesia noticed with the dexmedetomidine with comparison to other drugs or using a larger number of patients.
5. In other studies, the combination of oral ketamine with other sedative agents should be studied.

REFERENCES

REFERENCES

1. Sumiya K, Homma M, Watanabe M, Baba Y, Inomata S, Kihara S, et al. Sedation and plasma concentration of clonidine hydrochloride for pre-anesthetic medication in pediatric surgery. *Biological & pharmaceutical bulletin* 2003;26(4):421-3.
2. Mellon RD, Simone AF, Rappaport BA. Use of anesthetic agents in neonates and young children. *Anesthesia and analgesia* 2007;104(3):509-20.
3. Lasky T, Ernst FR, Greenspan J. Use of analgesic, anesthetic, and sedative medications during pediatric hospitalizations in the United States 2008. *Anesthesia and analgesia* 2012;115(5):1155-61.
4. Banchs RJ, Lerman J. Preoperative Anxiety Management, Emergence Delirium, and Postoperative Behavior. *Anesthesiology clinics* 2014;32(1):1-23.
5. Kain ZN, Mayes LC, Caldwell-Andrews AA, Karas DE, McClain BC. Preoperative anxiety, postoperative pain, and behavioral recovery in young children undergoing surgery. *Pediatrics* 2006;118(2):651-8.
6. MacLaren JE, Thompson C, Weinberg M, Fortier MA, Morrison DE, Perret D, et al. Prediction of preoperative anxiety in children: who is most accurate? *Anesthesia and analgesia* 2009;108(6):1777-82.
7. Cuzzocrea F, Gugliandolo MC, Larcian R, Romeo C, Turiaco N, Dominici T. A psychological preoperative program: effects on anxiety and cooperative behaviors. *Paediatric anaesthesia* 2013;23(2):139-43.
8. Paul M. CAPD news. *Journal of the Canadian Dental Association* 2003;69(5):281-4.
9. Johnson JV, Jeske AH, McDaniel HK, Herrera GQ. Development of an in vivo model for assessment of drug-induced vascular injury. *J Oral Maxillofac Surg* 1989;47(8):819-22.
10. Keith T. CAPD news. *Journal of the Canadian Dental Association* 2003;69(5):281-4.
11. Nelson T, Nelson G. The role of sedation in contemporary pediatric dentistry. *Dental clinics of North America* 2013;57(1):145-61.
12. Hicks CG, Jones JE, Saxon MA, Maupome G, Sanders BJ, Walker LA, et al. Demand in pediatric dentistry for sedation and general anesthesia by dentist anesthesiologists: a survey of directors of dentist anesthesiologist and pediatric dentistry residencies. *Anesthesia progress* 2012;59(1):3-11.
13. de Castro AM, de Oliveira FS, de Paiva Novaes MS, Araujo Ferreira DC. Behavior guidance techniques in Pediatric Dentistry: attitudes of parents of children with disabilities and without disabilities. *Special care in dentistry : official publication of the American Association of Hospital Dentists, the Academy of Dentistry for the Handicapped, and the American Society for Geriatric Dentistry* 2013;33(5):213-7.
14. Wells JJ, Lamprecht DE. AAP Committee on Drugs. *Clinical pediatrics* 1999;38(10):621-2.
15. Cote CJ. American Academy of Pediatrics sedation guidelines: are we there yet? *Archives of pediatrics & adolescent medicine* 2012;166(11):1067-9.

16. Clinical policy for procedural sedation and analgesia in the emergency department. American College of Emergency Physicians. *Annals of emergency medicine* 1998;31(5):663-77.
17. American Academy of Pediatrics. Committee on Psychosocial Aspects of C, Family H, Task Force on Pain in Infants C, Adolescents. The assessment and management of acute pain in infants, children, and adolescents. *Pediatrics* 2001;108(3):793-7.
18. Kaplan RF, Yang CI. Sedation and analgesia in pediatric patients for procedures outside the operating room. *Anesthesiology clinics of North America* 2002;20(1):181-94, vii.
19. Catalano K. What is in a name--sedation or anesthesia. *AORN journal* 2002;75(3):550-3.
20. Pacheco GS, Ferayorni A. Pediatric procedural sedation and analgesia. *Emergency medicine clinics of North America* 2013;31(3):831-52.
21. Cravero JP, Blike GT. Review of pediatric sedation. *Anesth Analg* 2004;77:11-5.
22. Fallah R, Nakhaei MH, Behdad S, Moghaddam RN, Shamszadeh A. Oral chloral hydrate vs. intranasal midazolam for sedation during computerized tomography. *Indian pediatrics* 2013;50(2):233-5.
23. Varughese AM, Hagerman N, Patino M, Wittkugel E, Schnell B, Salisbury S, et al. A comparison of inhalational inductions for children in the operating room vs the induction room. *Paediatric anaesthesia* 2012;22(4):327-34.
24. Mahoney G. Moderate sedation in paediatric dental patients. *Australian dental journal* 2012;57(4):522-3; author reply 3-4.
25. Beringer RM, Segar P, Pearson A, Greampet M, Kilpatrick N. Observational study of perioperative behavior changes in children having teeth extracted under general anesthesia. *Paediatric anaesthesia* 2014.
26. Cohen-Salmon D. [Perioperative psychobehavioural changes in children]. *Annales francaises d'anesthesie et de reanimation* 2010;29(4):289-300.
27. Kain ZN, Caldwell-Andrews AA, Mayes LC, Wang SM, Krivutza DM, LoDolce ME. Parental presence during induction of anesthesia: physiological effects on parents. *Anesthesiology* 2003;98(1):58-64.
28. Kain ZN, Sevarino F, Pincus S, Alexander GM, Wang SM, Ayoub C, et al. Attenuation of the preoperative stress response with midazolam: effects on postoperative outcomes. *Anesthesiology* 2000;93(1):141-7.
29. Kuttner L. Pediatric hypnosis: pre-, peri-, and post-anesthesia. *Paediatric anaesthesia* 2012;22(6):573-7.
30. Ashley PF, Williams CE, Moles DR, Parry J. Sedation versus general anaesthesia for provision of dental treatment in under 18 year olds. *The Cochrane database of systematic reviews* 2012;11:CD006334.
31. Riva J, Lejbusiewicz G, Papa M, Lauber C, Kohn W, Da Fonte M, et al. Oral premedication with midazolam in paediatric anaesthesia. Effects on sedation and gastric contents. *Paediatric anaesthesia* 1997;7(3):191-6.

32. Almenrader N, Passariello M, Coccetti B, Haiberger R, Pietropaoli P. Premedication in children: a comparison of oral midazolam and oral clonidine. *Paediatric anaesthesia* 2007;17(12):1143-9.
33. Hamid M, Khan MA, Khatri A, Akhtar I. Effectiveness of premedication at the time of separation from parent and mask induction in paediatric patients coming for congenital heart disease surgery. *Journal of the College of Physicians and Surgeons--Pakistan : JCPS* 2012;22(5):280-4.
34. Abback PS, Ben Sallah T, Hilly J, Skhiri A, Silins V, Brasher C, et al. [Opioid-sparing effect of ketamine during tonsillectomy in children]. *Annales francaises d'anesthesie et de reanimation* 2013;32(6):387-91.
35. Eghbal MH, Taregh S, Amin A, Sahmeddini MA. Ketamine improves postoperative pain and emergence agitation following adenotonsillectomy in children. A randomized clinical trial. *Middle East journal of anesthesiology* 2013;22(2):155-60.
36. Larenza MP, Landoni MF, Levionnois OL, Knobloch M, Kronen PW, Theurillat R, et al. Stereoselective pharmacokinetics of ketamine and norketamine after racemic ketamine or S-ketamine administration during isoflurane anaesthesia in Shetland ponies. *British journal of anaesthesia* 2007;98(2):204-12.
37. Fassoulaki A, Theodoraki K, Melemenis A. Pharmacology of sedation agents and reversal agents. *Digestion* 2010;82(2):80-3.
38. Mani V, Morton NS. Overview of total intravenous anesthesia in children. *Paediatric anaesthesia* 2010;20(3):211-22.
39. Roelofse JA. The evolution of ketamine applications in children. *Paediatric anaesthesia* 2010;20(3):240-5.
40. White M, de Graaff P, Renshof B, van Kan E, Dzoljic M. Pharmacokinetics of S(+) ketamine derived from target controlled infusion. *British journal of anaesthesia* 2006;96(3):330-4.
41. Anderson BJ, Palmer GM. Recent developments in the pharmacological management of pain in children. *Current opinion in anaesthesiology* 2006;19(3):285-92.
42. Lee JH, Kim K, Kim TY, Jo YH, Kim SH, Rhee JE, et al. A randomized comparison of nitrous oxide versus intravenous ketamine for laceration repair in children. *Pediatric emergency care* 2012;28(12):1297-301.
43. Sondhi V, Patnaik SK, Khullar A. Procedural sedation associated displacement of sharp oesophageal foreign body. *BMJ case reports* 2012;2012.
44. Kunisawa T, Nagata O, Iwasaki H. [Pharmacokinetic simulation of high-dose administration of dexmedetomidine for decubitus treatment]. *Masui The Japanese journal of anesthesiology* 2006;55(8):995-8.
45. Farag E, Argalious M, Abd-Elsayed A, Ebrahim Z, Doyle DJ. The use of dexmedetomidine in anesthesia and intensive care: a review. *Current pharmaceutical design* 2012;18(38):6257-65.
46. Hammer GB, Sam WJ, Chen MI, Golianu B, Drover DR. Determination of the pharmacodynamic interaction of propofol and dexmedetomidine during esophagogastroduodenoscopy in children. *Paediatric anaesthesia* 2009;19(2):138-44.

47. Bol C, Danhof M, Stanski DR, Mandema JW. Pharmacokinetic-pharmacodynamic characterization of the cardiovascular, hypnotic, EEG and ventilatory responses to dexmedetomidine in the rat. *The Journal of pharmacology and experimental therapeutics* 1997;283(3):1051-8.
48. Chrysostomou C, Schmitt CG. Dexmedetomidine: sedation, analgesia and beyond. *Expert opinion on drug metabolism & toxicology* 2008;4(5):619-27.
49. Petroz GC, Sikich N, James M, van Dyk H, Shafer SL, Schily M, et al. A phase I, two-center study of the pharmacokinetics and pharmacodynamics of dexmedetomidine in children. *Anesthesiology* 2006;105(6):1098-110.
50. McMorrow SP, Abramo TJ. Dexmedetomidine sedation: uses in pediatric procedural sedation outside the operating room. *Pediatric emergency care* 2012;28(3):292-6.
51. Potts AL, Warman GR, Anderson BJ. Dexmedetomidine disposition in children: a population analysis. *Paediatric anaesthesia* 2008;18(8):722-30.
52. Tobias JD, Gupta P, Naguib A, Yates AR. Dexmedetomidine: applications for the pediatric patient with congenital heart disease. *Pediatric cardiology* 2011;32(8):1075-87.
53. Carollo DS, Nossaman BD, Ramadhyani U. Dexmedetomidine: a review of clinical applications. *Current opinion in anaesthesiology* 2008;21(4):457-61.
54. Eisenach JC, Lavand'homme P, Tong C, Cheng JK, Pan HL, Virtanen R, et al. Antinociceptive and hemodynamic effects of a novel alpha2-adrenergic agonist, MPV-2426, in sheep. *Anesthesiology* 1999;91(5):1425-36.
55. Carroll CL, Krieger D, Campbell M, Fisher DG, Comeau LL, Zucker AR. Use of dexmedetomidine for sedation of children hospitalized in the intensive care unit. *Journal of hospital medicine : an official publication of the Society of Hospital Medicine* 2008;3(2):142-7.
56. Czaja AS, Zimmerman JJ. The use of dexmedetomidine in critically ill children. *Pediatric critical care medicine : a journal of the Society of Critical Care Medicine and the World Federation of Pediatric Intensive and Critical Care Societies* 2009;10(3):381-6.
57. Su F, Hammer GB. Dexmedetomidine: pediatric pharmacology, clinical uses and safety. *Expert opinion on drug safety* 2011;10(1):55-66.
58. Ihmsen H, Saari TI. [Dexmedetomidine. Pharmacokinetics and pharmacodynamics]. *Der Anaesthesist* 2012;61(12):1059-66.
59. Tobias JD. Controlled hypotension in children: a critical review of available agents. *Paediatric drugs* 2002;4(7):439-53.
60. Tobias JD, Berkenbosch JW, Russo P. Additional experience with dexmedetomidine in pediatric patients. *Southern medical journal* 2003;96(9):871-5.
61. Kulikov AS, Lubnin AY. [Dexmedetomidine: new opportunities in anesthesiology]. *Anesteziologiya i reanimatologiya* 2013(1):37-41.
62. Darlong V, Shende D, Subramanyam MS, Sunder R, Naik A. Oral ketamine or midazolam or low dose combination for premedication in children. *Anaesthesia and intensive care* 2004;32(2):246-9.

63. Zub D, Berkenbosch JW, Tobias JD. Preliminary experience with oral dexmedetomidine for procedural and anesthetic premedication. *Paediatric anaesthesia* 2005;15(11):932-8.
64. Song D, Joshi GP, White PF. Fast-track eligibility after ambulatory anesthesia: a comparison of desflurane, sevoflurane, and propofol. *Anesthesia and analgesia* 1998;86(2):267-73.
65. Davis PJ, Tome JA, McGowan FX, Jr., Cohen IT, Latta K, Felder H. Preanesthetic medication with intranasal midazolam for brief pediatric surgical procedures. Effect on recovery and hospital discharge times. *Anesthesiology* 1995;82(1):2-5.
66. Shane SA, Fuchs SM, Khine H. Efficacy of rectal midazolam for the sedation of preschool children undergoing laceration repair. *Annals of emergency medicine* 1994;24(6):1065-73.
67. Macnab AJ, Levine M, Glick N, Susak L, Baker-Brown G. A research tool for measurement of recovery from sedation: the Vancouver Sedative Recovery Scale. *Journal of pediatric surgery* 1991;26(11):1263-7.
68. Mortensen M, McMullin C. Discharge score for surgical outpatients. *The American journal of nursing* 1986;86(12):1347-9.
69. Meechan JG, Seymour RA. The use of third molar surgery in clinical pharmacology. *The British journal of oral & maxillofacial surgery* 1993;31(6):360-5.
70. Meechan JG, Seymour RA. The use of third molar surgery in clinical pharmacology. *Brit J Oral Maxillofac Surg* 1993;31:360-5.
71. Hatava P, Olsson GL, Lagerkranser M. Preoperative psychological preparation for children undergoing ENT operations: a comparison of two methods. *Paediatr Anaesth* 2000;10:477-86.
72. Kain ZN, Mayes LC, Wang SM, Caramico LA, Hofstadter MB. Parental presence during induction of anesthesia versus sedative premedication: which intervention is more effective? *Anesthesiology* 1998;89:1147-56.
73. Feld LH, Negus JB, White PF. Oral midazolam preanesthetic medication in pediatric outpatients. *Anesthesiology* 1990;73:831-4.
74. Singh N, Pandey RK, Saksena AK, Jaiswal JN. A comparative evaluation of oral midazolam with other sedatives as premedication in pediatric dentistry. *J Clin Pediatr Dent* 2002;26:161-4.
75. Sekerci C, Donmez A, Ates Y, Okten F. Oral ketamine premedication in children (placebo controlled double-blind study). *Eur J Anesthesiol* 1996; 13: 60661 1.
76. Knox JW, Boviil JG, Clarke RS, Dundee JW. Clinical studies of induction agents, ketamine. *Br J Anaesth* 2002;42:875-85.
77. Munro HM, Tirotta CF, Felix DE, Lagueruela RG, Madril DR, Zahn EM, et al. initial experience with dexmedetomidine for diagnostic and interventional cardiac catheterization in children . *Paediatr Anaesth* 2007;17:109-12.
78. Knox JW, Boviil JG, Clarke RS, Dundee JW. Clinical studies of induction agents, ketamine. *Br J Anaesth* 2002;42:875-86.

79. Gupta PK, Sehgal R, Gupta M, Kawatra R. Attenuation of cardiovascular responses to ketamine: Comparative evaluation of lignocaine and clonidine Indian J Anaesth 2002;46:457-9.
80. Yuen VM, Hui TW, Irwin MG, Yuen MK. A comparison of intranasal dexmedetomidine and oral midazolam for premedication in pediatric anesthesia: a double blinded randomized controlled trial. Anesth Analg 2008;106:1715-21.
81. Talke P, Chen R, Thomas B, Aggarwall A, Gottlieb A, Thorborg P, et al. the hemodynamic and adrenergic effects of perioperative dexmedetomidine infusion after vascular surgery Analg 2000;90:834-9.
82. Akin A, Bayram A, Esmoğlu A, Tosun Z, Aksu R, Altuntas R, et al. Dexmedetomidine vs midazolam for premedication of pediatric patients undergoing anesthesia. Paediatr Anaesth 2012;22:871-6.
83. Humphries Y, Melson M, Gore D. Superiority of oral ketamine as an analgesic and sedative for wound care procedures in pediatric patients with burns. J Burn Care Rehabil 1997;18:34-6.
84. Gutstein HB, Johnson KL, Heard MB, Gregory GA. Oral ketamine preanaesthetic medication in children. Anesthesiology 1992;76: 28-33.
85. Kamal K, Soliman D, Zakaria D. Oral dexmedetomidine versus oral midazolam as premedication in children. Ain Shams J Anesthesiol 2008;1.
86. Munro HM, Tirotta CF, Felix DE, Lagueruela RG, Madril DR, Zahn EM, et al. Initial experience with dexmedetomidine for diagnostic and interventional cardiac catheterization in children. Paediatr Anaesth 2007;17:109-12.
87. Yuen VM, Hui TW, Irwin MG, Yuen MK. A comparison of intranasal dexmedetomidine and oral midazolam for premedication in pediatric anesthesia: A double-blinded randomized controlled trial. Anesth Analg 2008;106:1715-21.
88. Schmidt AP, Valinetti EA, Bandeira D, Bertacchi MF, Simões CM, Auler JO Jr. Effects of preanesthetic administration of midazolam, clonidine, or dexmedetomidine on postoperative pain and anxiety in children. Paediatr Anaesth 2007; 17:667-74.
89. Tobias JD, Berkenbosch JW. Initial experience with dexmedetomidine in paediatric-aged patients. Pediatric Anesthesia 2002; 12: 171-5.
90. Jalowiecki P, Rudner R, Gonciarz M, Kawecki P, Petelencz M, Dziurdzik P. Sole use of dexmedetomidine has limited utility for conscious sedation during outpatient colonoscopy Anesthesiology 2005; 103: 269-73.
91. Alhashemi JA. Dexmedetomidine vs. midazolam for monitored anaesthesia care during cataract surgery. British Journal of Anaesthesia 2006; 96: 722-6.
92. Munro HM, Tirotta CF, Felix DE, et al. Initial experience with dexmedetomidine for diagnostic and interventional cardiac catheterization in children. Pediatric Anesthesia 2007; 17: 109-12.
93. Tobias JD, Phipps S, Smith B, Mulhern RK. Oral ketamine premedication to alleviate the distress of invasive procedures in paediatric oncology patients Pediatrics 1992; 90: 537-541.
94. Kain ZN, Mayes LC, Wang SM, Hofstadter MB. Postoperative behavioral outcomes in children: effects of sedative premedication. Anesthesiology 1999;90:758-65.

PROTOCOL

ORAL KETAMINE VERSUS ORAL
DEXMEDETOMIDINE PREMEDICATION FOR OUT-
PATIENT PAEDIATRIC DENTAL SURGERIES

الكيتامين عن طريق الفم مقابل الديكسميتوميدين عن طريق الفم ما قبل التخدير

للاطفال الخاضعين لعمليات الاسنان

٩٧

Protocol of a thesis submitted
to the Faculty of Medicine
University of Alexandria
In partial fulfillment of the
requirements of the degree of
**Master of Anaesthesia
And Surgical Intensive Care**

خطة بحث مقدمة
لكلية الطب
جامعة الإسكندرية
إيفاء جزئياً
لشروط الحصول على درجة
الماجستير في التخدير
والعناية المركزة الجراحية

By

من

Manar Mahmoud Elwi
MBBCh , Alex.
Resident
Alexandria University Hospitals
Department of Anaesthesia
And Surgical Intensive Care
Faculty of Medicine
University of Alexandria
2013

منار محمود علوى
بكالوريوس الطب والجراحة, الإسكندرية
طبيب مقيم
مستشفيات جامعة الإسكندرية
قسم التخدير
والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية
٢٠١٣

Handwritten signature of Manar Mahmoud Elwi

SUPERVISORS

المشرفون

Prof. Dr. Emad Aldin Abdel Moniem Arida أ.د. عماد الدين عبد المنعم عريضة
 Professor of Anaesthesia أستاذ التخدير
 And Surgical Intensive Care, والعناية المركزة الجراحية
 Faculty of Medicine, كلية الطب
 University of Alexandria. جامعة الإسكندرية

Prof. Dr. Wafaa Mohamed Shafshak أ.د/ وفاء محمد شفشق
 Professor of Anaesthesia أستاذ التخدير
 And Surgical Intensive Care, والعناية المركزة الجراحية
 Faculty of Medicine, كلية الطب
 University of Alexandria. جامعة الإسكندرية

Prof. Dr. Aly Sharaf أ.د/ علي شرف
 Professor of Paediatric Dentistry أستاذ طب اسنان الأطفال
 Faculty of dentistry, كلية اسنان
 Univeristy of Alexandria. جامعة الاسكندرية

Dr. Ola Mohy El deen Zanaty د. علا محيى الدين زنتى
 Assistant Professor of Anesthesia أستاذ مساعد التخدير
 And Surgical Intensive Care والعناية المركزة الجراحية
 Faculty of Medicine, كلية الطب
 University of Alexandria. جامعة الإسكندرية

ASSISTANT RESEARCHER

الباحث المساعد

Sameh Fathi Ali

سامح فتحى على

Sixth grade student

طالب بالفرقة السادسة

Faculty of Medicine,

كلية الطب

University of Alexandria.

جامعة الاسكندرية

Mobile phone: 0113412340

E-mail : same7oldlion33@yahoo.com

INTRODUCTION


Children undergoing surgical procedures are subjected to stressful events while preparing for surgery, such as the admission process, having blood drawn, receiving injections or having other medications administered. The children's stress continues while they are being transported, during induction and when they awaken before their parents arrive in the postanesthesia care unit. ⁽¹⁾

One of the challenges for paediatric anesthesiologists is to minimize distress for children in the perioperative period and to facilitate a smooth induction of anesthesia. ⁽²⁾

Maladaptive behavioral responses such as general anxiety, night time crying, enuresis and separation anxiety occur up to 44% of children two weeks after surgery and about 20% of these children will continue to demonstrate negative behavior 6 months after surgery. ⁽³⁾

Preoperative anxiety stimulates the sympathetic, parasympathetic and endocrine system leading to an increase in heart rate, blood pressure and cardiac excitability. ⁽⁴⁾

Anticipation of dental procedures induces a number of physiologic changes characteristic of the anxious state. Anxiety and fear associated with dental procedures have been well documented and thought to represent a model of acute stressor. ⁽⁵⁾




Dental treatment rendered under general anesthesia (GA) includes amalgam restorations, extractions, stainless steel crowns, composite/glass ionomer restorations, formocresol pulpotomy, fissure sealants, strip crowns and pulpectomy. Amalgam restorations and extractions constituted the most frequent dental procedure performed on child dental patients under GA.⁽⁶⁾

Unpremedicated children frequently object inhalational induction, and they often feel that the use of needles is one of the most worrisome aspects of the hospital stay. Children aged two to six years are especially vulnerable to this problem. Therefore, these children should be premedicated, in order to allow smooth induction, decrease anxiety and to prevent postoperative psychological and behavioral changes.⁽⁷⁾

various drugs have been advocated as premedication to allay anxiety and facilitate the smooth separation of children from parents. The ideal premedicant in children should be readily acceptable and should have a rapid and reliable onset with minimal side effects.⁽⁸⁾

Oral premedication are widely used in paediatric anaesthesia to reduce emotional trauma and ensure smooth induction.⁽⁹⁾

Ketamine is an arylcycloalkylamine that produces a state of sedation, anesthesia, immobility, analgesia, amnesia and dissociation from the environment. Pharmacologically, ketamine is classified as an N-methyl-d-aspartate (NMDA) receptor antagonist. It induces a state referred to as dissociative anesthesia by reducing or blocking signals to the conscious mind from other parts of the brain.^(10,11)

Handwritten signature and initials in black ink, appearing to be 'S. H. S.' followed by 'INSA'.


Dexmedetomidine (Dex) has recently been added to the anesthesia. It belongs to the class of α_2 agonists and possesses the properties of sedation, analgesia and opioid sparing effect. It differs from clonidine in being 16 times more specific for α_2 receptors and with a more selective action and a shorter half-life. (12,13)



AIM OF THE WORK

The present study is designed to compare orally administered ketamine and orally administered dexmedetomidine as a sedative premedication in children undergoing out-patient dental surgery under general anesthesia as regards:

- 1) Sedative effect
- 2) Reaction to separation from the parents
- 3) Ease of venipuncture
- 4) Recovery
- 5) Discharge times
- 6) Postoperative amnesia.



12/12/20

PATIENTS

This study will be carried out on 80 ASA I and II patients aged between 4 and 6 years admitted to Department of paediatric dentistry, Alexandria Main University Hospital, scheduled for elective out-patient dental surgery under general anesthesia.

Sample size is statistically approved by the biostatistics department of the High Institute Of Public Health, Alexandria University.

Exclusion criteria

Patients suffering from:

- Cardiac disease.
- Neuromuscular disease.
- Pulmonary disease (e.g. asthma).
- Anticipated difficult airway.
- Any condition with increased risk of regurgitation and aspiration of gastric contents.
- Patients suffering from pharyngeal pathology e.g. abscess and haematoma.
- known allergy or hypersensitivity reaction to any of the study drugs.



METHODS

After approval of Medical Ethics Committee of the Faculty of Medicine and taking an informed written consent from the parents, every patient will be subjected to a careful pre-anesthetic assessment including:


Pre-operative evaluation:

- History of medical illness and drugs.
- Complete physical examination.
- Routine laboratory investigation: Complete blood count, bleeding time, clotting time, prothrombin time, partial thromboplastin time and International normalized ratio .

Pre-medication

- Children will be randomly allocated to one of the two equal study groups using computer-generated random numbers.
 - I. Group K :will receive 5 mg/kg oral ketamine⁽¹⁴⁾
 - II. Group D: will receive oral dexmedetomidine 3 µg/kg .⁽¹⁵⁾

Both oral premedications will be diluted in 3 ml of apple juice to be given thirty minutes before induction of anaesthesia.
- Children will be given premedication in the preoperative holding area in the presence of one parent.
- Emla cream will be applied to the place of possible venipuncture.



Monitoring:

- All patients in the operating theater will be connected to multichannel monitor. (Trakmon-Kontron-Limited- England) for display of:
 - Non invasive blood pressure.
 - HR and lead II ECG
 - Oxygen saturation.

Anesthesia:

- On arrival to operating room, an intravenous cannula will be inserted.
- All patients will be subjected to the same anesthetic protocol

a) Induction:


Atropine (0.02mg/kg IV) followed by propofol (2-3 mg/kg IV)
Endotracheal intubation will be facilitated by atracurium (0.5mg/kg IV).
Intra-operative analgesia will be maintained by ketorolac (0.5mg/kg IV)

b) Maintenance:

- Anesthesia will be maintained by isoflurane (1-2%) in 100% oxygen. Ventilation will be mechanically controlled. Atracurium increments will be given as required.

c) Recovery:

At the end of the procedure isoflurane will be discontinued .
Neuromuscular block will be antagonized by neostigmine



(0.04mg/kg IV) and atropine (0.02mg /kg IV). Children will be extubated after return of protective reflexes.

d) **Discharge:**

Children will be discharged home on meeting discharge criteria.⁽¹⁶⁾

MEASUREMENTS:

1) Demographic data:

Age, sex and weight

2) Duration of surgery in minutes.

3) Vital signs

- Heart rate (beats/min).
- Mean arterial blood pressure (mmHg).
- Oxygen saturation percentage (SaO₂%).

Will be recorded at the following times:

- Base line (before sedation).
- Before induction of general anesthesia (30 min after sedation).
- During the procedure: Every 15 min.
- Post-operative: Every 30 min till discharge



Handwritten signatures and initials, including a large signature on the left, a smaller signature in the middle, and initials 'N.S.' on the right.

4) Assessment of sedation

will be recorded after admission to OR (30 min after sedation) using sedative score: ⁽¹⁷⁾

1	Sleeping, no response to patting the shoulder.
2	Sleeping, no response to call. Responds to patting on the shoulder.
3	Eyes closed, dull reaction. Responds to verbal stimulus.
4	Eyes open and closed by turns, dull reaction. Responds to verbal
5	Eyes open, dull reaction. Responds to verbal stimulus.
6	Normal reaction.
7	Irritable with body movement.

5) Assessment of ease of Separation and venipuncture

Will be graded according to Ease of Induction Score System: ⁽¹⁸⁾

1	Excellent	Patient unafraid, cooperative or asleep.
2	Good	Slight fear and/or crying, quiet with reassurance.
3	Fair	Moderate fear and crying, not quiet with reassurance.
4	Poor	Crying, need for restraint.

6) Assessment of recovery:⁽¹⁹⁾

will be assessed as base line then postoperative every 10 min till baseline score regained, using "Vancouver sedative recovery scale for children". Recovery time from discontinuation of anesthesia till regaining baseline score will be recorded.

	Parameter	Finding	Points
(A)	Awake-asleep	Awake and alert.	4
		Awake but drowsy.	3
		Asleep but easily aroused.	2
		Asleep and difficult to arouse.	1
		Asleep and unable to arouse.	0
(B)	Response to stimuli	Responds fully to stimuli.	2
		Delayed response to stimuli.	1
		Absent response to stimuli.	0
(C)	Facial expression	Alert.	1
		Flat.	0
(D)	Appearance of eyes	Bright eyes.	1
		Dull eyes; glazed.	0
(E)	Feeling how looks at you	Looks "at you"	1
		Looks "through you"	0
(F)	Visual stimulus	Recognition of stimulus.	1
		Limited or no recognition of stimulus.	0
(G)	Eye movement	Purposeful and spontaneous.	1
		Little or no spontaneous or purposeful.	0
(H)	Activity	Spontaneous & varied central activity.	4
		Spontaneous & varied peripheral activity.	3
		Central activity in response to stimuli.	2
		Peripheral activity in response to stimuli.	1
		No movement.	0
(I)	Tremor or ataxia	Absence of tremor or ataxia.	1
		Ataxia or tremor on being moved.	0
(J)	Spontaneous movement	Coordinated spontaneous movement.	2
		Weak/coarse spontaneous movement.	1
		No purposeful spontaneous movement.	0
(K)	Hand movements	Age-appropriate manual dexterity.	2
		Awkward or clumsy hand movement.	1
		No fine hand movement.	0

7) Discharge time: ⁽²⁰⁾


Discharge time from discontinuation of anesthesia until the patient is discharged to home, using "The Short-Stay Surgery Discharge Score" will be recorded (A score of 10-12 was needed for discharge).

Parameter	Finding	Points
Temperature	Severe chills and/or increased perspiration.	0
	Oral temperature > 38°C or < 36°C.	1
	Oral temperature 36-38°C.	2
Circulation	Blood pressure and heart rate changed more than 20 % from preoperative values.	0
	Standing blood pressure < supine blood pressure. Pulse increases >10 beats/min when standing.	1
	Standing blood pressure about the same as supine. Pulse increase < 10 beats/min when standing.	2
Activity	Unable to ambulate, sleepy.	0
	Out of bed with assistance.	1
	Out of bed without assistance and able to states discharge instructions.	2
Pain	Needs pain medicine administered IM.	0
	Needs pain medicine administered PO.	1
	Free of pain.	2
Bleeding	Large amount of surgical bleeding.	0
	Moderate amount of surgical bleeding.	1
	No surgical bleeding and no hematoma.	2
Intake Output	No PO fluids; unable to void.	0
	PO ice chips, sips fluids; voids into urinal.	1
	PO fluids without nausea or vomiting for 1 hour; voids quantity sufficient in bathroom.	2

[Handwritten signatures and initials]

8) Assessment of amnesia:⁽²¹⁾

The anterograde amnesia will be assessed after 24 hours by a recall questionnaire of the parents, which will include the questions about events which takes place after the administration of sedative.



RESULTS

The results obtained from this study will be tabulated and statistically analyzed using the standard statistical methods with the aid of different ways of presentation: numerical, mathematical and graphical.

[Handwritten signature]
[Handwritten signature]

DISCUSSION

The results of the study will be discussed in details and compared with any available and published data.



Handwritten signature and date: 12/1/2024

REFERENCES


1. Kain ZN, Caldwell-Andrews AA, Krivutza DM, Weinberg ME, Wang SM, Gaal D. Trends in the practice of parental presence during induction of anesthesia and the use of preoperative sedative premedication in the United States, 1995-2002: results of a follow-up national survey. *Anesth Analg* 2004;98(5):1252-9.
2. Kain ZN, Caldwell-Andrews AA, Krivutza DM, Weinberg ME, Wang SM, Gaal D. Trends in the practice of parental presence during induction of anesthesia and the use of preoperative sedative premedication in the United States, 1995-2002: results of a follow-up national survey. *Anesth Analg* 2004;98(5):1252-.
3. Hosey MT, Macpherson LM, Adair P, Tochel C, Burnside G, Pine C. Dental anxiety, distress at induction and postoperative morbidity in children undergoing tooth extraction using general anaesthesia. *Br Dent J* 2006;200(1):39-43.
4. Hosey MT, Asbury AJ, Bowman AW, Millar K, Martin K, Musiello T, et al. The effect of transmucosal 0.2 mg/kg midazolam premedication on dental anxiety, anaesthetic induction and psychological morbidity in children undergoing general anaesthesia for tooth extraction. *Br Dent J* 2009;207(1): 32-3.
5. Brand H, Abraham-Inpijn L. Cardiovascular responses induced by dental treatment. *Eur J Oral Sci* 1996;104:245-52
6. Al-Malik MI, Al-Sarheed MA. Comprehensive dental care of pediatric patients treated under general anesthesia in a hospital setting in Saudi Arabia. *J Contemp Dent Pract* 2006;7:79-88.



7. Kain ZN, Caldwell-Andrews AA, Maranets I, et al. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesth Analg* 2004;99:1648-54.
8. Sheta SA, Alsarheed M. Oral midazolam premedication for children undergoing general anaesthesia for dental care. *Int J Pediatr* 2009.
9. Richmond MN, Daum RE. Premedication with oral slow release morphine in dental anaesthesia. A comparison with temazepam. *Anaesthesia* 1988;43(8):694-6.
10. Cortinas M, Oya B, Caparros P, Cano G, Ibarra M, Martinez L. [Oral ketamine-midazolam premedication of uncooperative patients in major outpatient surgery]. *Rev Esp Anesthesiol Reanim* 2010;57(8):479-85.
11. Banerjee B, Bose A, Pahari S, Dan AK. A comparative study of paediatric oral premedication: midazolam, ketamine and low dose combination of midazolam and ketamine. *J Indian Med Assoc* 2011;109(6):386-8.
12. Yuen VM, Hui TW, Irwin MG, Yuen MK. A comparison of intranasal dexmedetomidine and oral midazolam for premedication in pediatric anesthesia: a double-blinded randomized controlled trial. *Anesth Analg* 2008;106(6):1715-21.
13. Mowafi HA, Aldossary N, Ismail SA, Alqahtani J. Effect of dexmedetomidine premedication on the intraocular pressure

A handwritten signature in black ink, appearing to be 'M. A. M.', is written across the bottom of the page. The signature is stylized and somewhat cursive.

- changes after succinylcholine and intubation. *Br J Anaesth* 2008;100(4):485-9.
14. Darlong V, Shende D, Subramanyam MS, Sunder R, Naik A. Oral ketamine or midazolam or low dose combination for premedication in children. *Anaesth Intensive Care* 2004;32(2):246-9.
 15. Zub D, Berkenbosch JW, Tobias JD. Preliminary experience with oral dexmedetomidine for procedural and anesthetic premedication. *Paediatr Anaesth* 2005;15(11):932-8.
 16. Song D, Joshi GP, White PF. Fast track eligibility after ambulatory anaesthesia: a comparison of desflurane, sevoflurane, and propofol. *Anesth, Analg* 1998;86:267-73.
 17. Peter J, Tome JA, Francis X, et al. Preanaesthetic medication with intranasal midazolam for brief pediatric surgical procedures: Effect on recovery and hospital discharge times. *Anesthesiology* 1995;82:2-5.
 18. Shane SA, Fuchs SM, Khine H. Efficacy of rectal midazolam for the sedation of preschool children undergoing laceration repair. *Ann Emerg Med* 1994;24:1065-73.
 19. Macnab AJ, Levine M, Glick N, Susak L, Baker-Brown G. A research tool for measurement of recovery from sedation: the Vancouver Sedative Recovery Scale. *J Pediatr Surg* 1991;26:1263-7.
 20. Mortensen M, McMullin C. Discharge scores for surgical outpatients. *Am J Nurs* 1986;86:1347-9.



21. Meechan JG, Seymour RA. The use of third molar surgery in clinical pharmacology. *Brit J Oral Maxillofac Surg* 1993;31:360-5.

A handwritten signature in black ink, consisting of several fluid, overlapping strokes.A handwritten signature in black ink, featuring a prominent loop and several sharp, angular strokes.A handwritten signature in black ink, with a series of connected, somewhat horizontal strokes.A handwritten signature in black ink, appearing as a series of overlapping, slightly curved lines.

ARABIC SUMMARY

الملخص العربي

من الواضح أن عدم وجود القدر الكافي من التهدئة قبل العمليات الجراحية في الأطفال ينتج عنه شعور هؤلاء الأطفال و عائلاتهم بالقلق خلال الفترة المحيطة بالعملية و أن زياده التوتر و القلق أثناء مرحله الإدخال التخديري من الممكن أن يؤدي لزياده الضلالات أثناء الأفاقه.

ان القلق المرتبط بعمليات الاسنان يمثل حالة من التوتر و توقع الخطر، و أن توقعها يحفز مجموعه من التغيرات الفسيولوجية المميزه لحاله القلق.

ان أدويه ما قبل التخدير في الأطفال عاده ما تكون في صوره ادويه مساعده للتخفيف من التوتر و الخوف المصاحب للعمليات الجراحية ، كما يسهل فصل الابناء عن الاباء و يحقق السهوله في الإدخال التخديري.

ان أفضل عقار للتهدئة قبل التخدير و طريقه إعطاءه لا يزال محل خلاف.

هدف البحث

كان الهدف من هذا البحث دراسة و مقارنة عقار الكيتامين و عقار الديكسميتوميدين عن طريق الفم في الأطفال للتهدئة قبل عمليات الاسنان الصغرى تحت تأثير التخدير الكلى من حيث التأثير على العلامات الحيوية، درجة التهدئة، سهوله الانفصال بين الطفل و ابويه ، سهوله ادخال قسطره وريديه طرفيه ، التأثير على الفتره الزمنية اللازمه للأفاقه من التخدير و الخروج للمنزل، و كذلك قدره العقار على أحداث فقدان للذاكره.

أجريت هذه الدراسة على ثمانين طفلا سليما ، تتراوح أعمارهم بين اربع و ست سنوات و كانوا بالوضع الصحى الاول او الثاني حسب تصنيف الجمعيه الامريكيه للتخدير، و الذى حدّد لهم سلفاً جراحات أسنان صغرى تحت تأثير التخدير الكلى.

وقد تم تقسيم المرضى إلى مجموعتين متساويتين حسب نوع العقار المستخدم كدواء مهدء قبل العملية:

المجموعة K: تم إعطائهم شراب عقار الكيتامين عن طريق الفم ، ثلاثون دقيقه قبل دخول حجره العمليات.

المجموعة D: تم إعطائهم شراب عقار الديكسميتوميدين عن طريق الفم ، ثلاثون دقيقه قبل دخول حجره العمليات.

تم تقييم كلّ المرضى قبل العمليات الجراحية عن طريق التعرف على التاريخ المرضي، الفحص السريري الكامل، تقييم الممرات الهوائية، و الإختبارات المعملية الروتينية. عند وصول المرضى إلى غرفة العمليات تم توصيلهم بأجهزة المراقبة القياسية و التي تشمل علي؛ رسم القلب، قياس ضغط الدم، و قياس نسبة تشبع الدم بالأوكسجين، ثم تركيب قسطره وريديه طرفيه. جميع المرضى الذين شاركوا في هذه الدراسة تم إخضاعهم لنفس نظام التخدير.

القياسات

لكل المرضى المشتركين في هذه الدراسة تم قياس معدل ضربات القلب و متوسط ضغط الدم الشرياني و نسبة تشبع الدم الشرياني بالأوكسجين باستمرار مع تسجيلها قبل العملية الجراحية وكل ١٥ دقيقه أثناء العملية وبعد العملية كل ٣٠ دقيقه. كذلك تم قياس درجة التهدئة قبل العملية الجراحية باستخدام مقياس التهدئة، و تم قياس سهوله الحصول علي وريد بعد استخدام الدواء بنصف ساعه باستخدام مقياس سرعه الإدخال التخديري و سرعه دخول القسطره الوريديه، كما تم قياس نسبه وعى الطفل بعد العملية بالمقارنه لقبول استخدام الدواء باستخدام مقياس التعافى التخديري، و قياس الوقت اللازم للإفاقة الكاملة و أماكنه الخروج الى المنزل، كما تم التحقق من قدره العقار على أحداث فقدان للذاكره عن طريق اسئله مباشره للاهل و الطفل.

نتائج البحث

- لا توجد فروق ذات دلالة إحصائية بين المجموعتين المشتركتين في هذه الدراسة من حيث السن والنوع ووزن الجسم، وقت العملية.
- لا توجد فروق ذات دلالة إحصائية بين المجموعتين المشتركتين في هذه الدراسة من حيث العلامات الحيوية.
- يوجد فروق ذات دلالة إحصائية بين معدل ضربات القلب بين المجموعتين عند ٣٠ دقيقه بعد اعطاء العقار و في لدقيقه ٠ و ٣٠ ما بعد العملية.
- يوجد فروق ذات دلالة إحصائية بين متوسط ضغط الدم الشرياني بين المجموعتين عند ٣٠ دقيقه بعد اعطاء العقار و في الدقيقه ٣٠ و ٦٠ بعد العملية.

- لا توجد فروق ذات دلالة احصائية ما بين المجموعتين فيما يتعلق بنسبه تشبع الاكسجين بالدم علي مدار الدراسه.
- درجه التهده كانت أفضل في المجموعه D مع وجود فارق ذو دلالة إحصائية بين المجموعتين.
- سهوله انفصال الطفل عن والديه و تركيب قسطره وريديه طرفيه كانت افضل في المجموعه D مع وجود فارق ذو دلالة احصائية بين المجموعتين.
- لا توجد فروق ذات دلالة إحصائية بين المجموعتين المشتركتين في هذه الدراسه من حيث الوقت اللازم للنهوض من التخدير و لأمكانيه الخروج الى المنزل.
- قدره العقار على أحداث فقدان للذاكره تحقق بنسبه أكبر في المجموعه D مع وجود فارق ذو دلالة إحصائية بين المجموعتين.

نستخلص من هذا البحث:

١. عقارى الكيتامين و الديكسميتوميدين عن طريق الفم ممكن استخدامهم لتهده الاطفال قبل العمليات الجراحيه بنجاح.
٢. اعطاء عقار الديكسميتوميدين عن طريق الفم وحده قبل العمليه الجراحيه كافى لتهده الطفل بصوره مرضيه و لا يحتاج بجانبه الي اى محاولات تعديل اخرى لتهده الطفل سلوكيا.
٣. اعطاء عقار الكيتامين عن طريق الفم وحده بجرعه ٥ مجم/كجم غير كافى لتهده الطفل تماما.

التوصيات

١. استخدام عقار الديكسميتوميدين عن طريق الفم قبل التخدير فى الأطفال بالجرعه و الوقت الامثل لتفادى اي اعراض جانبيه للعقار.
٢. عقار الديكسميتوميدين وحده غير كافى للقيام بعمليات مؤلمه نسبيا كما يجب ضروره دراسه تاثيره في فقدان الذاكره .
٣. ضروره دراسه استخدام عقار الكيتامين قبل التخدير فى الاطفال مع أدويه مهدئه أخرى.

الملخص العربي

لجنة الإشراف

أ.د./ عماد الدين عيد المنعم عريضه
أستاذ التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

أ.د./ وفاء محمد شفشق
أستاذ التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية

أ.د./ على عبد العزيز شرف
أستاذ طب اسنان أطفال
كلية طب الأسنان
جامعة الإسكندرية

د./ علا محيي الدين زنتاى
استاذ مساعد التخدير والعناية المركزة الجراحية
كلية الطب
جامعة الإسكندرية



الكيتامين عن طريق الفم مقابل الديكسميتوميدين عن طريق الفم ما قبل التخدير للاطفال
الخاضعين لعمليات الاسنان

مقدمة من

منار محمود علوى

بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٨

للحصول على درجة

الماجستير

فى

التخدير والعناية المركزة الجراحية

موافقون

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د/ سلوى شعبان شعراوى

أستاذ التخدير والعناية المركزة الجراحية

كلية الطب

جامعة الإسكندرية

أ.د/ عمرو محمد حلمى

أستاذ التخدير والعناية المركزة الجراحية

كلية الطب

جامعة قناة السويس

د/ علا محيى الدين زناتى

استاذ مساعد التخدير والعناية المركزة الجراحية

كلية الطب

جامعة الإسكندرية



الكيتامين عن طريق الفم مقابل الديكسميتوميدين عن طريق الفم ما قبل التخدير للاطفال الخاضعين لعمليات الاسنان

رسالة علمية

مقدمة إلى كلية الطب- جامعة الإسكندرية
إستيفاء للدراسات المقررة للحصول على درجة

الماجستير

فى

التخدير والعناية المركزة الجراحية

مقدمة من

منار محمود علوى

بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٨
طبيب مقيم مستشفيات جامعة الإسكندرية - قسم التخدير والعناية المركزة الجراحية
كلية الطب - جامعة الإسكندرية