

# CONCLUSION

## **CONCLUSION**

Choroidal thickness is decreased in diabetics relative to age-matched normal individuals. Choroid thickness is decreased as diabetic retinopathy progresses from mild NPDR to PDR. Presence of diabetic macular edema is associated with a significant decrease in the choroidal thickness.

# REFERENCES

## REFERENCES

- 1- Retina RSJ. 4<sup>th</sup> ed. Vol 1. Philadelphia, PA: Elsevier Mosby, 2006. 33-4
- 2- Cao J, McLeod S, Merges CA, Luty GA. Choriocapillaris degeneration and related pathologic changes in human diabetic eyes. *Arch Ophthalmol* 1998; 116: 589-97.
- 3- Gemenetzi M, De Salvo G, Lotery AJ. Central serous chorioretinopathy: an update on pathogenesis and treatment. *Eye (Lond)* 2010; 24: 1743-56.
- 4- Spaide RF. Age-related choroidal atrophy. *Am J Ophthalmol* 2009; 147: 801-10.
- 5- Torres VL, Brugnoli N, Kaiser PK, Singh AD. Optical coherence tomography enhanced depth imaging of choroidal tumors. *Am J Ophthalmol* 2011; 151: 586-93.
- 6- Maruko I, Iida T, Sugano Y, Saito M, Sekiryu T. Subfoveal retinal and choroidal thickness after verteporfin photodynamic therapy for polypoidal choroidal vasculopathy. *Am J Ophthalmol* 2011; 151: 594-603.
- 7- Inoue R, Sawa M, Tsujikawa M, Gomi F. Association between the efficacy of photodynamic therapy and indocyanine green angiography findings for central serous chorioretinopathy. *Am J Ophthalmol* 2010; 149: 441-6.
- 8- Shiraki K, Moriwaki M, Kohno T, Yanagihara N, Miki T. Age-related scattered hypofluorescent spots on late-phase indocyanine green angiograms. *Int Ophthalmol* 1999; 23: 105-9.
- 9- Nagaoka T, Kitaya N, Sugawara R, Yokota H, Mori F, Hikichi T, et al. Alteration of choroidal circulation in the foveal region in patients with type 2 diabetes. *Br J Ophthalmol* 2004; 88: 1060-3.
- 10- Rosen RB, Hathaway M, Rogers J, Pedro J, Garcia P, Dobre GM, et al. Simultaneous OCT/SLO/ICG imaging. *Invest Ophthalmol Vis Sci* 2009; 50: 851-60.
- 11- Spaide RF, Yannuzzi LA, Slakter JS, Sorenson J, Orlach DA. Indocyanine green videoangiography of idiopathic polypoidal choroidal vasculopathy. *Retina* 1995; 15: 100-10.
- 12- Yannuzzi LA. Indocyanine green angiography: a perspective on use in the clinical setting. *Am J Ophthalmol* 2011; 151: 745-51.
- 13- Desmettre T, Devoisselle JM, Mordon S. Fluorescence properties and metabolic features of indocyanine green (ICG) as related to angiography. *Surv Ophthalmol* 2000; 45: 15-27.
- 14- Pemp B, Schmetterer L. Ocular blood flow in diabetes and age-related macular degeneration. *Can J Ophthalmol* 2008; 43: 295-301.
- 15- Luty G, Grunwald J, Majji AB, Uyama M, Yoneya S. Changes in choriocapillaris and retinal pigment epithelium in age-related macular degeneration. *Mol Vis* 1999; 5: 35.
- 16- Falsini B, Anselmi GM, Marangoni D, D'Esposito F, Fadda A, Di Renzo A, et al. Subfoveal choroidal blood flow and central retinal function in retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 2011; 52: 1064-9.

- 17- Malhotra A, Minja FJ, Crum A, Burrowes D. Ocular anatomy and cross-sectional imaging of the eye. *Semin Ultrasound CT MR* 2011; 32: 2-13.
- 18- Branchini L, Regatieri CV, Flores-Moreno I, Baumann B, Fujimoto JG, Duker JS. Reproducibility of choroidal thickness measurements across three spectral domain optical coherence tomography systems. *Ophthalmology* 2012; 119: 119-23.
- 19- Margolis R, Spaide RF. A pilot study of enhanced depth imaging optical coherence tomography of the choroid in normal eyes. *Am J Ophthalmol* 2009; 147: 811-5.
- 20- Fujiwara T, Imamura Y, Margolis R, Slakter JS, Spaide RF. Enhanced depth imaging optical coherence tomography of the choroid in highly myopic eyes. *Am J Ophthalmol* 2009; 148: 445-50.
- 21- Imamura Y, Fujiwara T, Margolis R, Spaide RF. Enhanced depth imaging optical coherence tomography of the choroid in central serous chorioretinopathy. *Retina* 2009; 29: 1469-73.
- 22- Regatieri T, Manjunath V, Taha M, Fujimoto JG, Duker JS. Choroidal thickness in normal eyes measured using Cirrus HD optical coherence tomography. *Am J Ophthalmol* 2010; 150: 325-9.
- 23- Huang D, Swanson EA, Lin CP, Schuman JS, Stinson WG, Chang W, et al. Optical coherence tomography. *Science* 1991; 254: 1178-81.
- 24- Sull AC, Vuong LN, Price LL, Srinivasan VJ, Gorczynska I, Fujimoto JG, et al. Comparison of spectral/Fourier domain optical coherence tomography instruments for assessment of normal macular thickness. *Retina* 2010; 30: 235-45.
- 25- de Boer JF, Cense B, Park BH, Pierce MC, Tearney GJ, Bouma BE. Improved signal-to-noise ratio in spectral-domain compared with time-domain optical coherence tomography. *Opt Lett* 2003; 28: 2067-9.
- 26- Leitgeb R, Hitzinger C, Fercher A. Performance of fourier domain vs. time domain optical coherence tomography. *Opt Express* 2003; 11: 889-94.
- 27- Spaide RF, Koizumi H, Pozzoni MC. Enhanced depth imaging spectral-domain optical coherence tomography. *Am J Ophthalmol* 2008; 146: 496-500.
- 28- Kim YT, Ikuno Y, Maruko I, Yasuno Y, Miura M, Sekiryu T, et al. Reproducibility of retinal and choroidal thickness measurements in enhanced depth imaging and high-penetration optical coherence tomography. *Invest Ophthalmol Vis Sci* 2011; 52: 5536-40.
- 29- Nandakumar N, Branchini L, Regatieri CV, Walsh AC, Durbin MK, Keane PA, et al. Characterization of choroidal morphology in healthy eyes using spectral domain optical coherence tomograph. *Invest Ophthalmol Vis Sci* 2011; 52: 7019-26.
- 30- Manjunath V, Fujimoto JG, Duker JS. Cirrus HD-OCT high definition imaging is another tool available for visualization of the choroid and provides agreement with the finding that the choroidal thickness is increased in central serous chorioretinopathy in comparison to normal eyes. *Retina* 2010; 30: 1320-1.
- 31- Kim SW, Oh J, Kwon SS, Yoo J, Huh K. Comparison of choroidal thickness among patients with healthy eyes, early age-related maculopathy, neovascular age-related macular degeneration, central serous chorioretinopathy, and polypoidal choroidal vasculopathy. *Retina* 2011; 31: 1904-11.

- 32- Kim YT, Kang SW, Bai KH. Choroidal thickness in both eyes of patients with unilaterally active central serous chorioretinopathy. *Eye (Lond)* 2011; 25: 1635-40.
- 33- Maruko I, Iida T, Sugano Y, Ojima A, Sekiryu T. Subfoveal choroidal thickness in fellow eyes of patients with central serous chorioretinopathy. *Retina* 2011; 31: 1603-8.
- 34- Maruko I, Iida T, Sugano Y, Ojima A, Ogasawara M, Spaide RF. Subfoveal choroidal thickness after treatment of central serous chorioretinopathy. *Ophthalmology* 2010; 117: 1792-9.
- 35- Pryds A, Larsen M. Choroidal thickness following extrafoveal photodynamic treatment with verteporfin in patients with central serous chorioretinopathy. *Acta Ophthalmol Acta Ophthalmol* 2012; 90: 738-43.
- 36- Maruko I, Iida T, Sugano Y, Furuta M, Sekiryu T. One-year choroidal thickness results after photodynamic therapy for central serous chorioretinopathy. *Retina* 2011; 31: 1921-7.
- 37- Klein R, Klein BE, Linton KL. Prevalence of age-related maculopathy. The Beaver Dam Eye Study. *Ophthalmology* 1992; 99: 933-43.
- 38- Fung AE, Lalwani GA, Rosenfeld PJ, Dubovy SR, Michels S, Feuer WJ, et al. An optical coherence tomography-guided, variable dosing regimen with intravitreal ranibizumab (Lucentis) for neovascular age-related macular degeneration. *Am J Ophthalmol* 2007; 143: 566-83.
- 39- Koizumi H, Yamagishi T, Yamazaki T, Kawasaki R, Kinoshita S. Subfoveal choroidal thickness in typical age-related macular degeneration and polypoidal choroidal vasculopathy. *Graefes Arch Clin Exp Ophthalmol* 2011; 249: 1123-8.
- 40- Hidayat AA, Fine BS. Diabetic choroidopathy. Light and electron microscopic observations of seven cases. *Ophthalmology* 1985; 92: 512-22.
- 41- Weinberger D, Kramer M, Priel E, Gatton DD, Axer-Siegel R, Yassur Y. Indocyanine green angiographic findings in nonproliferative diabetic retinopathy. *Am J Ophthalmol* 1998; 126: 238-47.
- 42- Esmaelpour M, Povazay B, Hermann B, Hofer B, Kojic V, Hale SL, et al. Mapping choroidal and retinal thickness variation in type 2 diabetes using three-dimensional 1060-nm optical coherence tomography. *Invest Ophthalmol Vis Sci* 2011; 52: 5311-6.
- 43- Schocket LS, Brucker AJ, Niknam RM, Grunwald JE, DuPont J, Brucker AJ. Foveolar choroidal hemodynamics in proliferative diabetic retinopathy. *Int Ophthalmol* 2004; 25: 89-94.
- 44- Shiragami C, Shiraga F, Matsuo T, Tsuchida Y, Ohtsuki H. Risk factors for diabetic choroidopathy in patients with diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol* 2002; 240: 436-42.
- 45- Gupta V, Gupta A, Gupta P, Sharma A. Spectral-domain cirrus optical coherence tomography of choroidal striations seen in the acute stage of Vogt-Koyanagi-Harada disease. *Am J Ophthalmol* 2009; 147: 148-53.
- 46- Diniz B, Regatieri C, Andrade R, Maia A. Evaluation of spectral domain and time domain optical coherence tomography findings in toxoplasmic retinochoroiditis. *Clin Ophthalmol* 2011; 5: 645-50.

- 47- Bordaberry MF. Vogt-Koyanagi-Harada disease: diagnosis and treatments update. *Curr Opin Ophthalmol* 2010; 21: 430-5.
- 48- Herbort CP, Mantovani A, Bouchenaki N. Indocyanine green angiography in Vogt-KoyanagiHarada disease: angiographic signs and utility in patient follow-up. *Int Ophthalmol* 2007; 27:173-82.
- 49- Fong AH, Li KK, Wong D. Choroidal evaluation using enhanced depth imaging spectral-domain optical coherence tomography in Vogt-Koyanagi-Harada disease. *Retina* 2011; 31: 502-9.
- 50- Yasuno Y, Okamoto F, Kawana K, Yatagai T, Oshika T. Investigation of multifocal choroiditis with panuveitis by three-dimensional high-penetration optical coherence tomography. *J Biophotonics* 2009; 2: 435-41.
- 51- Vance SK, Khan S, Klancnik JM, Freund KB. Characteristic spectral-domain optical coherence tomography findings of multifocal choroiditis. *Retina* 2011; 31: 717-23.
- 52- Fretton A, Finger PT. Spectral domain-optical coherence tomography analysis of choroidal osteoma. *Br J Ophthalmol* 2012; 96: 224-8.
- 53- Say EA, Shah SU, Ferenczy S, Shields CL. Optical coherence tomography of retinal and choroidal tumors. *J Ophthalmol* 2010; 2011: 385058.
- 54- Witkin AJ, Ko TH, Fujimoto JG, Chan A, Drexler W, Schuman JS, et al. Ultra-high resolution optical coherence tomography assessment of photoreceptors in retinitis pigmentosa and related diseases. *Am J Ophthalmol* 2006; 142: 945-52.
- 55- Yeoh J, Rahman W, Chen F, Hooper C, Patel P, Tufail A, et al. Choroidal imaging in inherited retinal disease using the technique of enhanced depth imaging optical coherence tomography. *Graefes Arch Clin Exp Ophthalmol* 2010; 248: 1719-28.
- 56- Ho J, Branchini L, Regatieri C, Krishnan C, Fujimoto JG, Duker JS. Analysis of normal peripapillary choroidal thickness via spectral domain optical coherence tomography. *Ophthalmology* 2011; 118: 2001-7.
- 57- Maul EA, Friedman DS, Chang DS, Boland MV, Ramulu PY, Jampel HD, et al. Choroidal thickness measured by spectral domain optical coherence tomography: factors affecting thickness in glaucoma patients. *Ophthalmology* 2011; 118: 1571-9.
- 58- Mwanza JC, Hochberg JT, Banitt MR, Feuer WJ, Budenz DL. Lack of association between glaucoma and macular choroidal thickness measured with enhanced depth-imaging optical coherence tomography. *Invest Ophthalmol Vis Sci* 2011; 52: 3430-5.
- 59- Ehrlich JR, Peterson J, Parlitsis G, Kay KY, Kiss S, Radcliffe NM. Peripapillary choroidal thickness in glaucoma measured with optical coherence tomography. *Exp Eye Res* 2011; 92: 189-94.
- 60- Usui S, Ikuno Y, Miki A, Matsushita K, Yasuno Y, Nishida K. Evaluation of the choroidal thickness using high-penetration optical coherence tomography with long wavelength in highly myopic normal-tension glaucoma. *Am J Ophthalmol* 2011.
- 61- Park SC, De Moraes CG, Teng CC, Tello C, Liebmann JM, Ritch R. Enhanced depth imaging optical coherence tomography of deep optic nerve complex structures in glaucoma. *Ophthalmology* 2012; 119: 3-9.

- 62- Unterhuber A, Povazay B, Hermann B, Sattmann H, Chavez-Pirson A, Drexler W. In vivo retinal optical coherence tomography at 1040 nm-enhanced penetration into the choroid. *Opt Express* 2005; 13: 3252-8.
- 63- Povazay B, Bizheva K, Hermann B, Unterhuber A, Sattmann H, Fercher A, et al. Enhanced visualization of choroidal vessels using ultrahigh resolution ophthalmic OCT at 1050 nm. *Opt Express*. 2003; 11: 1980-6.
- 64- Povazay B, Hermann B, Unterhuber A, Hofer B, Sattmann H, Zeiler F, et al. Three-dimensional optical coherence tomography at 1050 nm versus 800 nm in retinal pathologies: enhanced performance and choroidal penetration in cataract patients. *J Biomed Opt* 2007; 12: 041211.
- 65- Chinn SR, Swanson EA, Fujimoto JG. Optical coherence tomography using a frequency-tunable optical source. *Opt Lett* 1997; 22: 340-2.
- 66- de Boer JF, Milner TE, van Gemert MJ, Nelson JS. Two-dimensional birefringence imaging in biological tissue by polarization-sensitive optical coherence tomography. *Opt Lett* 1997; 22: 934-6.
- 67- Drexler W, Fujimoto JG. State-of-the-art retinal optical coherence tomography. *Prog Retin Eye Res* 2008; 27: 45-88.
- 68- Choma MA, Yang C, Izatt JA. Instantaneous quadrature low-coherence interferometry with  $3 \times 3$  fiber-optic couplers. *Opt Lett* 2003; 28: 2162-4.
- 69- Choma MA, Hsu K, Izatt JA. Swept source optical coherence tomography using an all-fiber 1300-nm ring laser source. *J Biomed Opt* 2005; 10: 44009.
- 70- Srinivasan VJ, Huber R, Gorczynska I, Fujimoto JG, Jiang JY, Reisen P, et al. High-speed, high-resolution optical coherence tomography retinal imaging with a frequency-swept laser at 850 nm. *Opt Lett* 2007; 32: 361-3.
- 71- Huber R, Adler DC, Srinivasan VJ, Fujimoto JG. Fourier domain mode locking at 1050 nm for ultra-high-speed optical coherence tomography of the human retina at 236,000 axial scans per second. *Opt Lett* 2007; 32: 2049-51
- 72- Izatt JA, Kulkarni MD, Yazdanfar S, Barton JK, Welch AJ. In vivo bidirectional color Doppler flow imaging of picoliter blood volumes using optical coherence tomography. *Opt Lett* 1997; 22: 1439-41.
- 73- Leitgeb R, Schmetterer L, Drexler W. Real-time assessment of retinal blood flow with ultrafast acquisition by color Doppler Fourier domain optical coherence tomography. *Opt Express* 2012; 73.
- 74- White B, Pierce M, Nassif N, Cense B, Park B, Tearney G, et al. In vivo dynamic human retinal blood flow imaging using ultrahigh-speed spectral domain optical coherence tomography. *Opt Express* 2003; 11: 3490-7.
- 75- Wang Y, Lu A, Gil-Flamer J, Tan O, Izatt JA, Huang D. Measurement of total blood flow in the normal human retina using Doppler Fourier-domain optical coherence tomography. *Br J Ophthalmol* 2009; 93: 634-7.
- 76- Leitgeb RA, Schmetterer L, Hitzenberger CK, Fercher AF, Berisha F, Wojtkowski M, et al. Real-time measurement of in vitro flow by Fourier-domain color Doppler optical coherence tomography. *Opt Lett* 2004; 29: 171-3.

- 77- Makita S, Hong Y, Yamanari M, Yatagai T, Yasuno Y. Optical coherence angiography. *Opt Express* 2006; 14: 7821-40.
- 78- Miura M, Makita S, Iwasaki T, Yasuno Y. Three-dimensional visualization of ocular vascular pathology by optical coherence angiography in vivo. *Invest Ophthalmol Vis Sci* 2011.
- 79- Srinivasan VJ, Adler DC, Chen Y, Gorczynska I, Huber R, Duker JS, et al. Ultrahigh-speed optical coherence tomography for threedimensional and en face imaging of the retina and optic nerve head. *Invest Ophthalmol Vis S* 2008; 49: 5103-10.

# **PROTOCOL**

موافق  
2014/05/12  
ب. ب. ب.

العنوان  
د. ب. ب.

**CHOROIDAL THICKNESS IN PATIENTS WITH  
DIABETIC RETINOPATHY ANALYZED BY  
SPECTRAL-DOMAIN OPTICAL COHERENCE  
TOMOGRAPHY**

تحليل سمك المشيما في مرضى اعتلال الشبكية السكري باستخدام التصوير  
المقطعي بالإتساق البصري

Protocol of a thesis submitted  
to the Faculty of Medicine  
University of Alexandria  
In partial fulfillment of the  
requirements of the degree of  
**Master of Ophthalmology**

خطة بحث مقمنة  
لكلية الطب  
جامعة الإسكندرية  
إيفاء جزئياً  
لشروط الحصول على درجة  
الماجستير في طب وجراحة العين

By

من

**Essam Mohamed Mahmoud  
Bakour**  
MBBCh, Alex.  
Resident  
Damanhour Eye Hospital  
Department of Ophthalmology  
Faculty of Medicine  
University of Alexandria  
2014

عصام محمد محمود بكور  
بكالوريوس الطب والجراحة ، الإسكندرية  
طبيب مقيم  
مستشفى العيون بدمهور  
قسم طب و جراحة العين  
كلية الطب  
جامعة الإسكندرية  
٢٠١٤

ك. ب. ب.

د. ب. ب.

د. ب. ب.

د. ب. ب.

مواد  
الطب  
١٢/٥/٢٠٢٢

2.

### SUPERVISORS

### المشرفون

**Prof. Dr. Ahmed Magdy Beda**  
Professor of Ophthalmology,  
Faculty of Medicine,  
University of Alexandria.




أ.د/ أحمد مجدى بده  
أستاذ طب و جراحة العين  
كلية الطب  
جامعة الإسكندرية

### CO-SUPERVISOR

### المشرف المشارك

**Dr. Amir Ramadan Ahmed**  
Lecturer in Ophthalmology,  
Faculty of Medicine,  
University of Alexandria,  
For his experience in the field of retina.



د/ أمير رمضان أحمد  
مدرس طب و جراحة العين  
كلية الطب  
جامعة الإسكندرية  
وذلك لخبرته في مجال الشبكية

**ASSISTANT RESEARCHER**

الباحث المساعد

**Mahmoud Mohamed Awaly**

محمود محمد عوالي

Sixth grade student,

طالب بالفرقة السادسة

Faculty of Medicine,

كلية الطب

University of Alexandria,

جامعة الإسكندرية

**Mobile phone:** 01006213906

**E-mail:** weml.ec@hotmail.com

عوالي

ابراهيم عوالي

## INTRODUCTION

Diabetic retinopathy is a leading cause of vision loss worldwide.<sup>(1)</sup> The development of macular edema and proliferative retinopathy are major causes of visual impairment.<sup>(1,2)</sup> Clinical and experimental findings suggested that choroid vasculopathy in diabetes may play a role in the pathogenesis of diabetic retinopathy.<sup>(3,4,5)</sup> Various choroid abnormalities including obstruction of the choriocapillaris, vascular degeneration, choroid aneurysms, and choroidal neovascularization have been reported in histopathologic studies of diabetic eyes.<sup>(3,6,7)</sup> In addition, loss of alkaline phosphatase activity in the diabetic choriocapillaris that represents loss of viable endothelial cells and concurrent degeneration of the choriocapillaris.<sup>(3,4)</sup>

There are few clinical studies on choroidal angiopathy in diabetes. This is because of the difficulty of imaging the choroid in vivo. Indocyanine green angiography reveals both hyper and hypofluorescent spots in diabetic eyes, although the significance is unknown.<sup>(4)</sup> It has been proposed that the hypofluorescent spots result from ischemic changes of the choroidal vessels and represent either a dye filling delay or a defect of the choriocapillaris.<sup>(3,4)</sup> It is further postulated that the hyperfluorescent spots may be secondary to the presence of choroidal neovascularization, intra-choroidal microvasculature abnormalities, or nodules at the level of the choriocapillaris or underlying stroma.<sup>(4,5,6)</sup> Concurrently, the risk factors associated with diabetic choroidopathy include the presence of severe diabetic retinopathy, poor glycemic control, and the nature of the treatment regimen.<sup>(6)</sup>

CS

اسماء

Additionally, studies assessing the choroidal blood flow beneath the fovea with the use of laser Doppler flowmetry indicate a reduction of choroidal blood flow and volume in patients with nonproliferative and proliferative diabetic retinopathy.<sup>(6)</sup> There was a more prominent decrease in flow in the case of proliferative diabetic retinopathy. A better clinical understanding of choroidal damage might be important for an accurate assessment of diabetic eye disease, but adequate visualization of the choroid using optical coherence tomography (OCT) has not been possible until recently, owing to its posterior location and the presence of pigmented cells that attenuate the incident light. Recent reports showed successful examination and measurement of choroidal thickness in normal and pathologic states using the Heidelberg Spectralis OCT (Heidelberg Engineering, Heidelberg, Germany).<sup>(7)</sup>

with  
ilept

### AIM OF THE WORK

The aim of the work is to compare between choroidal thickness in patients with diabetic retinopathy and normal persons using enhanced-depth imaging spectral –domain optical coherence tomography.

### SUBJECTS

The study will include 40 adult eyes of adults recruited from the ophthalmology department at Alexandria main hospital.

The eye will be equally divided into two groups:

Group (1) will include 20 eyes having either non-proliferative diabetic retinopathy or proliferative diabetic retinopathy.

Group (2) will include 20 eyes of normal subjects as control.

**Exclusion criteria:**

1. Central retinal vein occlusion.
2. Branch retinal vein occlusion.
3. Secondary optic atrophy.
4. Myopic degeneration.
5. Choroidal neovascularization.
6. Media Opacity.

*Handwritten signature and date*  
12/12

## METHODS

The diabetic patients and the healthy adults will be subjected to the following:

- Full history taking including age, sex, past ocular and systemic history.
- In the diabetic group, further inquiry will be done on diabetic mellitus type, duration and previous treatment applied to the eye.
- Full ophthalmic examination including visual acuity assessment, IOP measurement, anterior segment and posterior segment examination.
- Investigation will include:
  - Fluorescein angiography to determine the stage of diabetic retinopathy. (in the diabetic group only).
  - Optical coherence tomography of the macula area and of the choroid using enhanced depth imaging of the Heidelberg Spectralis OCT. (in both groups).

ic/pt

CSJ

o b e . c o m


### RESULTS

The results of this study will be assessed, tabulated and statistically analyzed in appropriate figures and tables.

*Handwritten marks:*  
A signature and the word "desp" are visible in the lower middle section of the page.

## DISCUSSION

The results will be discussed in view of achievement of the aim, their significance and their comparison with previous related researches.



## REFERENCES

1. Weinberger D, Kramer M, Priel E, et al. "Indocyanine green angiographic findings in nonproliferative diabetic retinopathy". *Am J Ophthalmol* 1998; 126:238-47.
2. Kobayashi S, Fukuta M, Kontani H, et al. "A quantitative assay for angiogenesis of cultured choroidal tissues in streptozotocin-diabetic Wistar and spontaneously diabetic GK rats". *Jpn J Pharmacol* 1998; 78:471-8.
3. Mac Gregor LC, Rosecan LR, Laties AM, Matschinsky FM. "Altered retinal metabolism in diabetes. I. Microanalysis of lipid, glucose, sorbitol, and myo-inositol in the choroid and in the individual layers of the rabbit retina". *J Biol Chem* 1986; 261:4046-51.
4. Nagaoka T, Kitaya N, Sugawara R, et al. "Alteration of choroidal circulation in the foveal region in patients with type 2 diabetes". *Br J Ophthalmol* 2004; 88:1060-3.
5. Yang L, Gong H, Wang Y, et al. "Nicotine alters morphology and function of retinal pigment epithelial cells in mice". *Toxicol Pathol* 2010; 38:560-7.
6. Fujiwara T, Imamura Y, Margolis R, et al. "Enhanced depth imaging optical coherence tomography of the choroid in highly myopic eyes". *Am J Ophthalmol* 2009; 148:445-50.
7. Spaide RF. Age-related choroidal atrophy. *Am J Ophthalmol* 2009;147:801-10.



\_\_\_\_\_

# **ARABIC SUMMARY**

## المخلص العربي

اعتلال الشبكية السكري هو السبب الرئيسي لفقدان البصر في جميع أنحاء العالم. التأثير على البقعة الصفراء بالشبكية واعتلال الشبكية السكري هي الأسباب الرئيسية لضعف البصر.

اعتلال وعائي المشيمة في مرض السكري قد تلعب دوراً في التسبب في اعتلال الشبكية السكري.

والهدف من هذا العمل هو المقارنة بين سمك المشيمة في المرضى الذين يعانون من اعتلال الشبكية السكري والأشخاص العاديين باستخدام التصوير المقطعي الأكثر عمقاً.

وشملت الدراسة أربعين عين من الكبار البالغين من قسم طب وجراحة العيون بمستشفى جامعة الإسكندرية الرئيسية.

وتضمنت المجموعة الأولى عشرين عين مصابة بمرض اعتلال الشبكية السكري.

وتضمنت المجموعة الثانية عشرين عين طبيعية لأشخاص لا تعاني من المرض.

وقد تم دراسة تأثير مرض اعتلال الشبكية السكري على المشيمة باستخدام التصوير المقطعي الأكثر عمقاً عن طريق خط في اتجاهين الرأسى والأفقى وقد تم قياس سمك المشيمة يدوياً من الحافة الخارجية إلى الحافة الداخلية للمشيمة.

وقد تم قياس سمك المشيمة في مرض السكري وقد وجد أنه له صلة بمرض اعتلال الشبكية السكري وكما زاد شدة المرض وجد انخفاض ملحوظ في سمك المشيمة.

الأوعية الدموية للمشيمة هي المسؤولة عن وظيفة الشبكية وعن حجم الدم في المشيمة وأى تأثير على هذه الأوعية يؤثر على تدفق الدم للمشيمة والشبكية وبالتالي يؤدي إلى التأثير على البصر.

لقد وضعت نظرية أن انخفاض سمك المشيمة قد يكون ذات صلة إلى نقص الأكسجين بأنسجة الشبكية.

والمشيمة هي المصدر الرئيسى لتغذية الشبكية وطبقات الشبكية الخارجية وأى تعرض لنقص تدفق الدم والأكسجين يؤثر على شبكية العين في مرض السكري.

وهذه الدراسة من عيوبها قلت عدد الحالات المستخدمة.

وفى الختام أن سمك المشيمة يقل بشكل ملحوظ في مرض اعتلال الشبكية السكري كلما اشتد المرض وسوف يكون هناك حاجة إلى مزيد من الدراسة والأبحاث على سمك المشيمة في مرض اعتلال الشبكية السكري.

# الملخص العربي

## لجنة الإشراف

.....  
أ.د./ أحمد مجدى بده  
أستاذ طب وجراحة العين  
كلية الطب  
جامعة الإسكندرية

.....  
د./ امير رمضان احمد  
مدرس طب وجراحة العين  
كلية الطب  
جامعة الإسكندرية

تحليل سمك المشيمة فى مرضى اعتلال الشبكية السكرى باستخدام التصوير المقطعى  
بالإتساق البصرى

مقدمة من

عصام محمد محمود بكور

بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٩

للحصول على درجة

الماجستير

فى

طب وجراحة العيون

موافقون

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د./ دلال أحمد شوقي

أستاذ طب وجراحة العيون

كلية الطب

جامعة الإسكندرية

أ.د./ أحمد مجدي بده

أستاذ طب وجراحة العيون

كلية الطب

جامعة الإسكندرية

أ.د./ ياسر رجب سراج

أستاذ طب وجراحة العيون

كلية الطب

جامعة طنطا



# تحليل سمك المشيمة فى مرضى اعتلال الشبكية السكرى باستخدام التصوير المقطعى بالإتساق البصرى

رسالة علمية

مقدمة إلى كلية الطب- جامعة الإسكندرية  
إستيفاء للدراسات المقررة للحصول على درجة

الماجستير

فى

طب وجراحة العيون

مقدمة من

عصام محمد محمود بكور

بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٩