

LIST OF ABBREVIATIONS

ABCDE	: Airway, breathing, circulation, disability, exposure & environment.
AIS	: Abbreviated Injury Score.
AMPLE	: Allergies, medications, previous medical/surgical history, last meal (Time), events environment surrounding the injury.
APRS	: Anterior pararenal space.
ARF	: Anterior renal fascia.
ATLS	: Advanced Trauma Life Support.
AVPU	: Alert, Voice, Pain, Unresponsive.
BAT	: Blunt abdominal trauma.
CNS	: Central nervous system.
CT	: Computed tomography.
D	: Duodenum.
	DIAGNOSTIC PERITONEAL LAVAGE.
DPL	:
ED	: Emergency department.
ERCP	: Endoscopic retrograde cholangio-pancreatogram.
FAST	: Focused assessment with sonography for trauma.
GCS	: Glasgow coma scale.
GU	: Genitourinary.
IV	: Intravenous.
IVC	: Inferior vena cava.
L	: Lumbar.
LCF	: Lateroconal fascia.
MI	: Milliliter.
MRCP	: Magnetic retrograde cholangio-pancreatogram.
MVC	: Motor vehicle crash..
O₂	: Oxygen.
OIS	: Organ Injury Scale.
PP	: Posterior peritoneum.
PPRS	: Posterior pararenal space.
PRF	: Posterior renal fascia.
RPH	: Retroperitoneal hematoma.
RPI	: Retroperitoneal injuries.
RTA	: Road traffic accident.
TF	: Transversalis fascia
US	: Ultrasonography.
WBCS	: White blood cells.

List of Figures

	Page
Figure (1): An illustration of fascial planes and spaces of the retroperitoneum.	3
Figure (2): Right renal contusion .Axial enhanced CT image shows a focal area of decreased enhancement (arrow) in the right kidney.	5
Figure (3): Abdominal CT scan with IV contrast. (Pancreatic transaction). (a) At the level of the pancreas shows a full-thickness linear low-attenuation laceration of the pancreatic neck (arrow) just anterior to the proximal portal vein. (b) Inferior level of the pancreas to shows a focal hematoma at the transection site that extends into the pancreatic head (arrow).	8
Figure (4): Abdominal CT scan obtained with IV and oral contrast. (a) extra luminal air (arrows) adjacent to the duodenum(Duodenal perforation) (b) The third portion of the duodenum is thickened and edematous (arrowheads) (Duodenal hematoma).	9
Figure (5): abdominal CT scan obtained with IV contrast (Traumatic adrenal hemorrhage) shows a right adrenal hemorrhage (arrowheads) with periadrenal hemorrhage (arrow).	11
Figure (6): An illustration of the various grades in severity of kidney injury.	13
Figure (7): An illustration of anatomic division of the ureter.	15
Figure (8): Zones of retroperitoneum. 1- Central-medial retroperitoneal zone. 2- Lateral retroperitoneal zone. 3- Pelvic retroperitoneal zone.	17
Figure (9): Abdominal CT scan with IV contrast (Small zone I retroperitoneal hematoma) shows a small amount of intermediate-attenuation fluid (arrowheads) between the aorta and IVC.	18
Figure (10): Paraspinal hematoma due to a burst fracture. (a) Contrast-enhanced CT scan shows a paraspinal hematoma (arrowheads) that surrounds the L5 vertebral body. (b) CT scan (bone window) shows a burst fracture of L5 (arrow).	19
Figure(11): Abdominal CT scan (Retroperitoneal fluid without retroperitoneal organ injury) shows fluid within the anterior pararenal space (arrowheads) diffusely surrounding the pancreas.	20
Figure (12): An illustration of the 4 scanning windows of the FAST examination.	28
Figure (13): Percentage of patients with BAT.	30
Figure (14): Shows age incidence in the present study.	31
Figure (15): Shows patient's gender in the present study	32
Figure (16): Mode of injury in the present study.	33
Figure (17): The physical examination in the present study.	34
Figure (18): Ultrasound diagnosis of the RPI patients.	35
Figure (19): CT finding of patients with proven RPI.	36
Figure (20): Abdominal CT scan with IV contrast (Retroperitoneal air) shows extra luminal air (arrowheads) in the right perirenal space, in the anterior pararenal space, and along the fascial planes of the abdominal musculature.	37

Figure (21): Contrast-enhanced CT scan shows right-sided perirenal and posterior pararenal retroperitoneal hematoma (white arrowheads) from a right renal laceration, with active extravasation of IV contrast material (black arrow) from a renal parenchymal injury. A laceration of the right kidney (black arrowhead). hyperenhancing, thickened small bowel (white arrows).	37
Figure (22): Associated non retroperitoneal lesions with patients with proven RPI detected by CT.	38
Figure (23): Associated injury with patients with proven RPI.	39
Figure (24): pelvic radiograph shows widening of symphysis pubis (long arrow), widening of left sacroiliac joint (short arrow) and a displaced fracture of the right femoral shaft (arrowhead).	40
Figure (25): Chest radiograph with pneumothorax on the left side with left lung collapse.	40
Figure (26): Management of patients with proven RPI.	41
Figure (27): Blood transfusion of patients with proven RPI.	42
Figure (28): Duration of hospital stay of patients with proven RPI.	43
Figure (29): Morbidity of patients with proven RPI.	44
Figure (30): Mortality of patients with proven RPI.	45

List of Tables

	Page
Table (1): Anatomic spaces of the retroperitoneum.	2
Table (2): American Association for the Surgery of Trauma Pancreatic OIS.	8
Table (3): American Association for the Surgery of Trauma duodenal OIS.	10
Table (4): American Association for the Surgery of Trauma Adrenal OIS.	11
Table (5): American Association for the Surgery of Trauma Kidney OIS.	12
Table (6): American Association for the Surgery of Trauma Ureter OIS.	15
Table (7): Anatomic Zones of the Retroperitoneum.	16
Table (8): Shows Glasgow Coma Scale.	24
Table (9): Shows age incidence in the present study.	31
Table (10): Shows patient's gender in the present study.	32
Table (11): Mode of injury in the present study.	33
Table (12): The physical examination in the present study.	34
Table (13): Ultrasound diagnosis of the RPI patients.	35
Table (14): CT finding of patients with proven RPI.	36
Table (15): Associated non retroperitoneal lesions with patients with proven RPI detected by CT.	38
Table (16): Associated injury with patients with proven RPI.	39
Table (17): Management of patients with proven RPI.	41
Table (18): Duration of hospital stay of patients with proven RPI.	43
Table (19): Morbidity of patients with proven RPI.	44
Table (20): Mortality of patients with proven RPI.	45