

## LIST OF FIGURES

Figures	Page
<b>Figure (1):</b> Bones of the elbow	4
<b>Figure (2):</b> Ossification centres of the distal end of the humerus	5
<b>Figure (3):</b> Capsule and ligaments of the elbow joint.	6
<b>Figure (4):</b> Anastamosis around elbow joint	8
<b>Figure (5):</b> Ulnar nerve crossing elbow.	11
<b>Figure (6):</b> Radial nerve crossing elbow	12
<b>Figure (7):</b> Median nerve crossing elbow	13
<b>Figure (8):</b> Brachial artery crossing elbow.	14
<b>Figure (9):</b> a) humeral-ulnar angle b) metaphyseal diaphyseal angle	16
<b>Figure (10):</b> Baumann angle	17
<b>Figure (11):</b> a: Anterior humeral line b: Humerotrochlear angle	18
<b>Figure (12):</b> : Mechanism of injury in extension type of supracondylar fracture of the humerus	20
<b>Figure (13):</b> a): Posteromedially displaced distal fragment with liability to radial nerve injury b): Posterolaterally displaced distal fragment with liability to medial nerve and Brachial artery injury	22
<b>Figure (14):</b> Motor neurological assessment: A) Normal "OK" sign. B) Normal thumb up sign. C) Normal starfish sign. D) Abnormal "OK" sign.	25
<b>Figure (15):</b> Flow chart to guide decision making in cases in which a possible vascular injury is associated with a displaced supracondylar humeral fracture	28
<b>Figure (16) :</b> Fishtail deformity	30
<b>Figure (17):</b> Manipulative reduction of supracondylar fracture of the humerus in children	32
<b>Figure (18):</b> Avoid elbow hyperflexion	32
<b>Figure (19):</b> Skin traction	33
<b>Figure (20):</b> a) Modified Palmer olecranon screw b) Overhead skeletal traction	34

<b>Figures</b>		<b>Page</b>
<b>Figure (21):</b>	Schematic representation of lateral external fixation	<b>35</b>
<b>Figure (22):</b>	Antegrade flexible intramedullary nailing	<b>37</b>
<b>Figure (23):</b>	patient position for closed pinning	<b>38</b>
<b>Figure (24):</b>	skin incision of anterior approach	<b>39</b>
<b>Figure (25):</b>	skin incision of medial approach	<b>40</b>
<b>Figure (26):</b>	skin incision of lateral approach	<b>41</b>
<b>Figure (27):</b>	skin incision of posterior approach.	<b>42</b>
<b>Figure (28):</b>	Distribution of the studied cases according to age	<b>44</b>
<b>Figure (29):</b>	Distribution of the studied cases according to sex	<b>45</b>
<b>Figure (30):</b>	Distribution of the studied cases according to side affected	<b>45</b>
<b>Figure (31):</b>	Distribution of the studied cases according to lag to surgery	<b>46</b>
<b>Figure (32):</b>	Distribution of the studied cases according to Skin condition	<b>47</b>
<b>Figure (33):</b>	skin incision	<b>49</b>
<b>Figure (34):</b>	identification of the ulnar nerve	<b>50</b>
<b>Figure (35):</b>	triceps muscle flap	<b>50</b>
<b>Figure (36):</b>	fixation of the fracture	<b>51</b>
<b>Figure (37):</b>	a. closure of the triceps muscle, b. closure of the skin.	<b>51</b>
<b>Figure (38):</b>	Distribution of the studied cases according to result	<b>55</b>
<b>Figure(39-42):</b>	Case 1	<b>69-71</b>
<b>Figure (43-46):</b>	Case 2	<b>72-73</b>
<b>Figure (47-50):</b>	: Case 3	<b>74-75</b>
<b>Figure (51-54):</b>	: Case 4	<b>76-77</b>

## LIST OF TABLES

<b>Tables</b>	<b>Page</b>
<b>Table (I):</b> Distribution of the studied cases according to age.	<b>44</b>
<b>Table (II):</b> Distribution of the studied cases according to lag to surgery	<b>46</b>
<b>Table(III):</b> Flynn's criteria	<b>52</b>
<b>Table (IV):</b> Distribution of the studied cases according to result.	<b>55</b>
<b>Table (V):</b> Distribution of the studied cases according to operation time	<b>56</b>
<b>Table (VI):</b> Distribution of the studied cases according to heal time.	<b>56</b>
<b>Table (VII):</b> Distribution of the studied cases according to loss of the range of motion after removal of the cast	<b>57</b>
<b>Table (VIII):</b> Distribution of the studied cases according to loss of the range of motion after the end of follow up.	<b>57</b>
<b>Table (IX):</b> Distribution of the studied cases according to loss of carrying angle at the end of the follow up	<b>58</b>
<b>Table (X):</b> Distribution of the studied cases according to infection after removal of the cast	<b>58</b>
<b>Table (XI):</b> Distribution of the studied cases according to Baumann angle immediately post-operative	<b>59</b>
<b>Table (XII):</b> Distribution of the studied cases according to frontal humeral line immediately post-operative.	<b>59</b>
<b>Table (XIII):</b> Distribution of the studied cases according to humerotrochlear angle immediately post- operative	<b>60</b>
<b>Table (XIV):</b> Distribution of the studied cases according Stability of the k-wires	<b>60</b>
<b>Table (XV):</b> Distribution of the studied cases according to Baumann angle at the end of the follow up	<b>61</b>
<b>Table (XVI):</b> Distribution of the studied cases according to frontal humeral line at the end of follow up.	<b>61</b>
<b>Table (XVII):</b> Distribution of the studied cases according to humerotrochlear angle after the end of follow up.	<b>62</b>
<b>Table (XVIII):</b> Relation between result with age (years)	<b>63</b>
<b>Table (XIX):</b> Relation between results with sex	<b>63</b>
<b>Table (XX)</b> Relation between results with the side affected	<b>64</b>
<b>Table (XXI)</b> Relation between results with operation time	<b>64</b>
<b>Table (XXII)</b> Relation between result with heal time	<b>65</b>
<b>Table (XXIII)</b> Relation between results with time elapsed before surgery	<b>65</b>

<b>Tables</b>		<b>Page</b>
<b>Table (XXIV)</b>	Relation between results with position of the k-wires	<b>66</b>
<b>Table (XXV)</b>	Relation between result and affected baumann's angle at the end of follow up	<b>66</b>
<b>Table (XXVI)</b>	Relation between results and position of the frontal humeral line at the end of follow up	<b>67</b>
<b>Table (XXVII)</b>	Relation between results and humerotrochlear angle at the end of the follow up	<b>67</b>

## LIST OF ABBREVIATIONS

<b>SCFH</b>	: Supracondylar fracture humerus
<b>K-wires</b>	: Kirschner wires
<b>FIN</b>	: Flexible Intramedullary Nailing
<b>ROM</b>	: Range of Motion
<b>CT</b>	: Computerized Tomography
<b>CTA</b>	: Computerized Tomographic Angiography
<b>CRPP</b>	: Closed Reduction and Percutaneous Pinning
<b>ORIF</b>	: Open Reduction and Internal Fixation
<b>AP</b>	: Anteroposterior
<b>AVN</b>	: Avascular Necrosis
<b>OR</b>	: Operating Room