

## **Acknowledgment**

I wish first to thank Allah for helping me to accomplish this research and for providing me with such supportive supervisors.

I would like to express my profound gratitude and appreciation to Professor Dr. Shaaban Mohammad Al Assal, Prof. of Clinical Oncology and Nuclear Medicine, Faculty of Medicine, Alexandria University who devoted great efforts and lot of his time and interest in supervising this work together with his guidance and careful instructions.

I wish to express my deep appreciation to Professor Dr. Yousry Abdelmaguid Rostom, Professor of Clinical Oncology and Nuclear Medicine, Faculty of Medicine, Alexandria University, for his kind supervision, continuous guidance, and constructive remarks.

My special thanks to Dr. Azza Mohammad Darwish Lecturer in Clinical Oncology and Nuclear Medicine Faculty of Medicine, Alexandria University, for her continuous help, valuable advice and support.

I'm also thankful to all staff members and to my colleagues in the Clinical Oncology and Nuclear Medicine Department for their help.

# LIST OF CONTENTS

<b>ACKNOWLEDGMENT</b> .....	<b>i</b>
<b>LIST OF CONTENT</b> .....	<b>ii</b>
<b>LIST OF TABLES</b> .....	<b>iii</b>
<b>LIST OF FIGURES</b> .....	<b>v</b>
<b>I. INTRODUCTION</b> .....	<b>1</b>
<b>II. AIM OF THE WORK</b> .....	<b>27</b>
<b>III. MATERIALS AND METHODS</b> .....	<b>28</b>
<b>IV. RESULTS</b> .....	<b>30</b>
<b>V. DISCUSSION</b> .....	<b>65</b>
<b>VI. SUMMARY</b> .....	<b>71</b>
<b>VII. CONCLUSIONS AND RECOMMENDATIONS</b> .....	<b>73</b>
<b>VIII. REFERENCES</b> .....	<b>74</b>
<b>PROTOCOL</b>	
<b>ARABIC SUMMARY</b>	

## LIST OF TABLES

Table	Page
(1) Distribution of the studied cases in relation to all head and neck cancer and all body tumors:	30
(2) Distribution of the studied cases according to primary site (n=255)	30
(3) Distribution of hypopharynx cancer cases according to primary involved subsite (n=104)	31
(4) Distribution of oral and oropharynx cancer cases according to primary site (n=151)	31
(5) Distribution of studied patients according to epidemiologic data	32
(6) Distribution of studied cases according to associated co morbidities: (n=255)	33
(7) Distribution of all patients according to predisposing factors (n = 255)	33
(8) Distribution of our study patients according to presenting signs and symptoms (n=255)	34
(9) Distribution of all patients according to histopathological types (n = 255)	34
(10) Distribution of hypopharynx cancer cases according to stage at diagnosis (n=104)	35
(11) Distribution of oral and oropharynx cancer cases according to stage (n=140)	36
(12) Distribution of all cases according to investigations (n=255)	37
(13) Distribution of the all patients according to type of biopsy	38
(14) Distribution of the studied cases according to whether patients received treatment or not	38
(15) Distribution of all patients according to treatment modality	39
(16) Distribution of all patients according to surgical treatment	40
(17) Distribution of the studied cases according to multiple positive LNS and positive margin	40
(18) Distribution of the studied cases according to surgery radiation time interval	41
(19) Distribution of all hypopharyngeal cancer patients according to chemotherapeutic treatment	41
(20) Distribution of all oral and oropharyngeal cancer patients according to chemotherapeutic treatment	42
(21) Distribution of hypopharyngeal cancer patients according to	43

<b>Table</b>	<b>Page</b>
	concomitant chemo-radiotherapy (CCRTx)
(22)	Distribution of Oral and Oropharynx cases according to concomitant chemo-radiotherapy 44
(23)	Distribution of Hypopharyngeal cancer patients according to radiotherapy 45
(24)	Distribution of Oral and Oropharyngeal cancer patients according to radiotherapy 46
(25)	Distribution of all patients according to development of relapse 47
(26)	Distribution of hypopharynx cancer patients according to salvage therapy 48
(27)	Distribution of oral and oropharyngeal cancer patients according to salvage therapy 49
(28)	Distribution of the studied cases according to 2nd primary 50
(29)	Distribution of the studied cases according to palliation 50
(30)	Distribution of the studied cases according to Condition at last visit 51
(31)	Overall survival estimates of hypopharyngeal cancer patients in relation to patients' age 54
(32)	Disease free survival estimates of hypopharyngeal cancer patients in relation to patients' age 54
(33)	Overall survival estimates of oral and oropharyngeal cancer patients in relation to patients' age 55
(34)	Disease free survival estimates of oral and oropharyngeal cancer patients in relation to patients' age 55
(35)	OS estimates in relation to degree of tumor differentiation in hypopharyngeal cancer patients 56
(36)	DFS estimates in relation to degree of tumor differentiation in hypopharyngeal cancer patients 56
(37)	OS estimates in relation to degree of tumor differentiation in oral and oropharyngeal cancer patients 57
(38)	DFS estimates in relation to degree of tumor differentiation in oral and oropharyngeal cancer patients 57

## LIST OF FIGURES

<b>Figure</b>		<b>page</b>
<b>(1.A)</b>	lateral view of the oral cavity	<b>4</b>
<b>(1.B)</b>	Kaplan-Meier overall survival estimates for oral, oropharyngeal, and Hypopharyngeal cancer patients	<b>52</b>
<b>(2)</b>	Kaplan-Meier disease free survival estimates for oral, oropharyngeal, and Hypopharyngeal cancer patients	<b>53</b>
<b>(3)</b>	Kaplan-Meier Stage stratified overall survival estimates for oral, oropharyngeal, and Hypopharyngeal cancer patients	<b>59</b>
<b>(4)</b>	Kaplan-Meier Stage stratified disease free survival estimates for oral, oropharyngeal, and Hypopharyngeal cancer patients	<b>60</b>
<b>(5)</b>	Kaplan Meier OS estimates in relation to type of treatment administered to hypopharyngeal cancer patients	<b>61</b>
<b>(6)</b>	Kaplan Meier DFS estimates in relation to type of treatment administered to hypopharyngeal cancer patients	<b>62</b>
<b>(7)</b>	OS estimates in relation to type of applied treatment for oral and oropharyngeal cancer patients using Kaplan Meier method	<b>63</b>
<b>(8)</b>	DFS estimates in relation to type of applied treatment for oral and oropharyngeal cancer patients using Kaplan Meier method	<b>64</b>