

LIST OF TABLES

Table		Page
(1)	Common sources of noise.	2
(2)	Speed of sound in different media.	27
(3)	Noise levels due to various sources (representative values).	33
(4)	Average normal noise levels for several acoustical environments.	37
(5)	Groups exposed to collection noise of different sources.	58
(6)	Groups exposed to predominant frequency of different sources.	58
(7)	Effect of noise exposure on the blood glucose level (mg/dl).	80
(8)	Effect of noise exposure on the WBCs counts (cells/ μ l).	82
(9)	Changes on the SOD level (U/ mg protein) after noise exposure.	84
(10)	Effect of noise exposure on the MDA level (nmol/mg protein).	86
(11)	Effect of noise exposure on the serotonin level (μ gm/ml).	88
(12)	Effect of noise exposure on the Recognition Index.	90
(13)	Parameters measured after exposure to factory noise.	94
(14)	Parameters measured after exposure to factory noise.	100
(15)	Parameters measured after exposure to furnace noise.	106
(16)	Changed in brain and small intestine as result of noise exposure ranged from mild to moderate.	112

LIST OF FIGURES

Figure		Page
(17)	Transverse waves.	19
(18)	Longitudinal waves.	19
(19)	Vibration of air particles about its equilibrium position.	20
(20)	Areas of compression (A) and rarefaction (B) produced by vibrating tuning fork.	21
(21)	(a, b) Sound waves traveling through a medium change localized air particle density.	22
(22)	Typical sine wave.	23
(23)	Two vibratory patterns that differ in frequency the panel on top is higher in frequency (higher pitch) than the panel on bottom.	24
(24)	Two vibratory patterns that differ in amplitude. The panel on top is higher in amplitude than the panel on bottom.	25
(25)	Relation between frequency and wavelength.	25
(26)	a. Vibratory patterns that differ in phase.	26
	b. The system for representation phase.	26
(27)	Two kinds of interference (a) constructive interference (b) destructive interference.	27
(28)	Time and frequency domain representations of sinusoid.	28
(29)	Time and frequency domain representations of complex periodic signals	29
(30)	Time and frequency domain representations of continuous aperiodic sounds.	29
(31)	Time domain representations and amplitude spectra for transients aperiodic sounds.	30
(32)	Spectrum envelope.	31
(33)	Approximate sound intensity levels for assortment of familiar sound sources.	33
(34)	Fletcher-Munson curves (audibility curve).	34
(35)	Frequency bands for acoustic waves.	35
(36)	Block diagram showing how a signal $x(n)$ would be converted to the cepstral domain.	36
(37)	The wave shape of noise.	37
(38)	Spectrum of white noise.	37
(39)	Spectrum of a type of colored noise.	38

Figure		Page
(40)	Schematic diagram of the auditory system.	40
(41)	Functional diagram of the human ear.	41
(42)	The stages of processing that lead to hearing.	42
(43)	Effects of oxidative stress on human.	43
(44)	Chemical structure of MDA.	44
(45)	Typical neurons.	47
(46)	Chemical structure of serotonin.	48
(47)	Synthesis of serotonin (5-hydroxytryptamine).	48
(48)	Synthesis of serotonin	49
(49)	Metabolism of serotonin.	49
(50)	Sector of polyurethane foam used as a sound insulator.	53
(51)	Sound level meter.	53
(52)	Sound level meter mode of action.	54
(53)	Spectrum is of the word 'Audacity' spoken aloud.	56
(54)	Factory sound.	57
(55)	Traffic sound.	57
(56)	Furnace sound.	57
(57)	Exposure system (Anechoic chamber).	59
(58)	Standard curve of serotonin.	61
(59)	Standard curve of SOD.	63
(60)	Standard curve of MDA.	64
(61)	Standard curve of total protein.	65
(62)	Standard curve of glucose.	66
(63)	The open field arena.	68
(64)	a. Factory noise amplitude-time plot in unexpanded mode. (The axis is in a logarithmic scale)	70
	b. Expanded band of the selected portion of the recorded factory noise.	70
(65)	Factory noise frequency-intensity. The X-axis represents the sound frequency in Hz, while the Y-axis represents the sound intensity in (dB SPL)	71
(66)	Factory noise signal cepstrum analysis	71
(67)	The standard autocorrelation of the recorded factory noise signal	72

Figure	Page
(68) a. Traffic noise amplitude-time plot in unexpanded mode. (The axis is in a logarithmic scale).	73
b. Expanded band of the selected portion of the recoded traffic noise.	73
(69) Factory noise frequency-intensity diagram.	74
(70) Traffic noise signal cepstrum analysis.	74
(71) The standard autocorrelation of the recorded traffic signal noise.	75
(72) a. Furnace noise amplitude-time plot in unexpanded mode. (The axis is in a logarithmic scale).	76
b. Expanded band of the selected portion of the recoded furnace noise.	76
(73) Furnace noise frequency-intensity diagram.	77
(74) The furnace noise signal cepstrum analysis.	77
(75) The standard autocorrelation of the recorded traffic signal noise.	78
(76) Effect of noise exposure on the blood glucose level (mg/dl).	81
(77) Effect of noise exposure on the WBCs counts (cells/ μ l).	83
(78) Effect of noise exposure on the SOD level (U/ mg protein).	85
(79) Effect of noise exposure on the MDA level (nmol/mg protein).	87
(80) Effect of noise exposure on the serotonin level (μ gm/ml).	89
(81) Effect of noise exposure on the Recognition Index.	91
(82) Mean values of blood glucose level (mg/dl) related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group(G0).	95
(83) Mean of WBCs counts (cells/ μ l) related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group (G0).	95
(84) Mean values of SOD (U/mg protein) related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group (G0).	96
(85) Mean values of MDA (nmol/mg protein) related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group (G0).	96
(86) Mean values serotonin (μ g/ml) related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group (G0).	97
(87) Mean values of RI related to different factory groups (GIa, GIaf, GIc, GIcf) in comparsion to control group (G0).	97
(88) Mean values of blood glucose (mg/dl) related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparsion to control (G0).	101
(89) Mean counts of WBCs (cells/ μ l) related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparsion to control (G0).	101
(90) Mean values of SOD (U/mg protein) related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparsion to control (G0).	102

Figure		Page
(91)	Mean values of MDA (nmol/mg protein) related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparison to control (G0).	102
(92)	Mean values of serotonin (µg/ml) related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparison to control (G0).	103
(93)	Mean values of RI related to different traffic groups (GIIa, GIIaf, GIIc, GIIcf) in comparison to control group (G0).	103
(94)	Mean values of blood glucose level (mg/dl) related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	107
(95)	Mean values of WBCs (cells/µl) counts related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	107
(96)	Mean values of SOD (U/mg protein) related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	108
(97)	Mean values of MDA (nmol/mg protein) related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	108
(98)	Mean values of serotonin related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	109
(99)	Mean values of RI related to different furnace groups (GIIIa, GIIIaf, GIIIc, GIIIcf) in comparison to control (G0).	109
(100)	Correlation between recognition index and glucose.	110
(101)	Correlation between recognition index and SOD.	110
(102)	Correlation between recognition index and MDA.	111
(103)	Photomicrograph of brain of normal mouse revealed normal tissue with clear purkingi fiber without synaptic lose. (H &E, 40X).	113
(104)	Photomicrograph of small intestine of normal mouse revealed normal villi, normal crypt, without any signs of inflammation. (H &E, 40X).	113
(105)	Photomicrograph of brain of mouse exposed to acute factory noise (100 dB, 12 hrs) (GIa) showing some chromatolysis change, vaculation and mild astrocytosis. (H &E, 40X).	114
(106)	Photomicrograph of small intestine of mouse exposed to acute factory noise (100 dB, 12 hrs) (GIa) showing moderate inflammatory changes, mild edema in the mucosa and musculosa. (H &E, 40X).	114
(107)	Photomicrograph of brain of mouse exposed to acute traffic noise (100 dB, 12 hrs) (GIIa) showing mild to moderate vaculation, moderate astrocytosis with condensed chromatin. (H &E, 40X).	115
(108)	Photomicrograph of small intestine of mouse exposed to acute traffic noise (100 dB, 12 hrs) (GIIa) showing mild to moderate inflammatory cells, note the edema between muscle bundle. (H &E, 40X).	115

Figure		Page
(109)	Photomicrograph of brain of mouse exposed to acute furance noise (100 dB, 12 hrs) (GIII _a) showing mild to mooderate vaculation, astrocytosis, with condensed chromatin. (H &E, 40X).	116
(110)	Photomicrograph of small intestine of mouse exposed to acute furance noise (100 dB, 12 hrs) (GIII _a) showing normal villi with mild distortion, mild inflammatory changes, separation of muscle fiber by edema. (H &E, 40X).	116
(111)	Photomicrograph of brain of mouse exposed to acute factory predominant frequency (100 dB, 12 hrs) (GI _{af}) showing mild vaculation, mild astrocytosis. (H &E, 40X).	117
(112)	Photomicrograph of small intestine of mouse exposed to acute factory predominant frequency (100 dB, 12 hrs) (GI _{af}) showing normal intestinal villi, normal crypt with mild inflamatory changes, mild edema. (H &E, 40X).	117
(113)	Photomicrograph of brain of mouse exposed to acute traffic predominant frequency (100 dB, 12 hrs) (GII _{af}) showing modulate vaculation, moderate astrocytosis. (H &E, 40X).	118
(114)	Photomicrograph of small intestine of mouse exposed to acute traffic predominant frequency (100 dB, 12 hrs) (GII _{af}) showing mild distortion in intestinal villi, with mild to moderate inflamatory changes, mild edema. (H &E, 40X).	118
(115)	Photomicrograph of brain of mouse exposed to acute furance predominant frequency (100 dB, 12 hrs) (GIII _{af}) showing mild vaculation, mild astrocytosis. (H &E, 40X).	119
(116)	Photomicrograph of small intestine of mouse exposed to acute furance predominant frequency (100 dB, 12 hrs) (GIII _{af}) showing mild distortion in intestinal villi, with mild inflamatory changes, mild edema. (H &E, 40X).	119
(117)	Photomicrograph of brain of mouse exposed to chronic factory noise (100 dB, 8 hrs/21d) (GI _c) showing mild vaculation, moderate astrocytosis. (H &E, 40X).	120
(118)	Photomicrograph of small intestine of mouse exposed to chronic traffic noise (100 dB, 8 hrs/21 d) (GI _c) showing mild distortion in intestinal villi, with mild to moderate inflamatory changes, mild edema. (H &E, 40X).	120
(119)	Photomicrograph of brain of mouse exposed to chronic traffic noise (100 dB, 8 hrs/21d) (GII _c) showing modulate vaculation, moderate astrocytosis, and condensed chromatin. (H &E, 40X).	121
(120)	Photomicrograph of small intestine of mouse exposed to chronic traffic noise (100 dB, 8 hrs/21 d) (GII _c) showing mild distortion in intestinal villi, with moderate inflamatory changes. (H &E, 40X).	121

Figure		Page
(121)	Photomicrograph of brain of mouse exposed to chronic furance noise (100 dB, 8 hrs/21d) (GIII _c) showing modulate vaculation, moderate astrocytosis. (H &E, 40X).	122
(122)	Photomicrograph of small intestine of mouse exposed to chronic traffic noise (100 dB, 8 hrs/21 d) (GIII _c) showing mild distortion in intestinal villi, with moderate inflamatory changes, moderate edema. (H &E, 40X).	122
(123)	Photomicrograph of brain of mouse exposed to chronic factory predominant frequency (100 dB, 8 hrs/21d) (GI _{cf}) showing modulate vaculation, moderate astrocytosis. (H &E, 40X).	123
(124)	Photomicrograph of small intestine of mouse exposed to chronic factory predominant frequency (100 dB, 8 hrs/21 d) (GI _{cf}) showing mild distortion in intestinal villi, with moderate inflamatory changes, and mild edema. (H &E, 40X).	123
(125)	Photomicrograph of brin of mouse exposed to chronic traffic predominant frequency (100 dB, 8 hrs/21d) (GII _{cf}) showing mild to modulate vaculation, mild astrocytosis. (H &E, 40X).	124
(126)	Photomicrograph of brain of mouse exposed to chronic traffic predominant frequency (100 dB, 8 hrs/21d) (GII _{cf}) showing mild to modulate vaculation, mild astrocytosis. (H &E, 40X).	124
(127)	Photomicrograph of brainof mouse exposed to chronic furance predominant frequency (100 dB, 8 hrs/21d) (GIII _{cf}) showing modulate vaculation, mild astrocytosis. (H &E, 40X).	125
(128)	Photomicrograph of small intestine of mouse exposed to chronic furance predominant frequency (100 dB, 8 hrs/21 d) (GIII _{cf}) showing moderate distortion in intestinal villi, with moderate inflamatory changes, and mild edema. (H &E, 40X).	125

LIST OF ABBREVIATIONS

5-HT	: 5-hydroxytryptamine (serotonin).
ABR	: Auditory brainstem response.
ACH	: Acetylcholine.
BP	: Blood pressure.
c	: Velocity of sound.
CAT	: Catalase.
CNs	: Central nervous system
d	: day.
DA	: Dopamine.
dB	: Decibel.
DBP	: Diastolic blood pressure.
EDTA	: Ethylene diamine tetra acetic acid
f	: Frequency.
f₀	: Fundamental frequency
FSH	: Follicle Stimulating hormone
FT	: Fourier Transform.
g, gm	: Gram.
GI	: Gastrointestinal.
GMA	: Gastromyocardial activity.
GPx	: Glutathione peroxidase.
GSH	: Oxidized glutathione.
GSSG	: Reduced glutathione.
HFN	: High frequency noise.
HR	: Heart rate.
hr	: hour.
HTN	: Hypertension.
Hz	: Hertz.
I	: Intensity.
I₀	: Reference intensity.
ICU	: Intensive care unit.
IFT	: Inverse fourier transform
Ig	: Immunoglobulin.

K-	: kilo-
L	: Length
l	: Liter
LFN	: Low frequency noise.
LH	: Luteinizing hormone
LL	: Loudness level
LPO	: Lipid peroxidation.
M	: Molar.
Max	: Maximum.
MDA	: Malondialdehyde.
min	: minute.
Min.	: Minimum.
MSC	: Mesenchymal stem cells.
μ	: micro-
n-	: nano-
NE	: Norepinephrine.
NIHL	: Noise induced hearing lose.
NO	: Nitric oxide.
NOR	: Novel Object Recognition.
NT	: Neurotransmitter.
OPT	: O- phthaldialhyde
P	: Pressure.
p	: probability of error
PA	: Public address amplifier.
Pa	: Pascale
RBCs	: Red blood Cells
RI	: Recognition index.
RMS	: Root mean square.
RNS	: Reactive nitrogen species.
ROS	: Reactive oxygen species.
rpm	: Round per minute
RS	: Reactive species.

RTA	: Real Time Audio.
sec	: Second.
SBP	: Systolic blood pressure.
SD	: Standard Deviation.
SDS	: Sodium dodecyl Sulphate
SLM	: Sound Level Meter
SOD	: Superoxide dismutase.
SPL	: Sound pressure level.
T	: Period.
t	: Time.
TBA	: Thiobarbituric acid
TMP	: Tetra methoxy propane
t₀	: Fundamental period
W	: Watt
WBCs	: White blood cells.
Wk	: Week.
WHO	: World Health Organization
Y	: Year.
λ	: Wavelength.