

## CONCLUSIONS

- Road traffic accidents are the most common cause of blunt abdominal trauma.
- Blunt abdominal trauma commonly affects the male children especially above 5 years.
- In our hospital, the role of ultrasound in the assessment of blunt abdominal trauma continued to grow in the past decade. Also, U/S have become the screening modality of choice in blunt abdominal trauma as it can help to determine the need for surgical intervention within minutes of a patient's arrival. U/S has replaced diagnostic peritoneal lavage and has redefined.
- The role of CT in the immediate evaluation of the trauma patient which may be performed if urgent laparotomy is not required.
- In patients who have minor injuries and are otherwise asymptomatic, can be managed conservatively and safely discharged with no need for CT.
- At the same time, they need a close interaction with the surgical team to reduce the risk of missed injury.
- Physicians must rely on various diagnostic tests to arrive at a diagnosis of blunt abdominal injury. Currently, the methods used in the assessment of blunt abdominal trauma include medical history and physical examination, FAST and CT.

## RECOMMENDATIONS

- It is recommended that all hospitals receiving patients with blunt abdominal trauma should have reliable clinical guidelines to allow emergency physicians to screen all patients for possible abdominal affection for early management.
- As FAST is a good bed side rapid screening test with good sensitivity and specificity, it should be used for detection of IAIs and CT should be preserved when further data are needed as grading for organ injury and in cases where FAST is not helpful.
- Recommended clinical guidelines to avoid unnecessary abdominal CT utilization in the evaluation for intra-abdominal injury following blunt torso trauma.
- Abdominal CT scan is not necessary in the initial evaluation of children with a reliable physical examination and negative
- Non operative treatment should be initially considered for the management of children with BAT. Emergency laparotomy is indicated mainly for the children who exhibit signs necessitating surgical intervention such as ongoing hemodynamic instability, pneumoperitoneum, peritonitis or diaphragmatic rupture.

## REFERENCES

1. Hamidi MI, Aldaoud KM, Qtaish I. Role of computed tomography in blunt abdominal trauma. *Sultan Qaboos Univ Med J* 2007; 7(3): 41-6.
2. Saxena M, Grewal H. Pediatric Abdominal Trauma. [Internet]. [Updated 2013 July 17, cited 2014 November 22]. Available from: <http://emedicine.medscape.com/article/1984811-overview>.
3. Wegner S, Colletti JE, Van Wie D. Pediatric blunt abdominal trauma. *Pediatr Clin North Am* 2006; 53: 243-56.
4. Holmes JF, Sokolove PE, Brant WE, Palchak MJ, Vance CW, Owings JT, et al. Identification of children with intra-abdominal injuries after blunt trauma. *Ann Emerg Med* 2002; 39: 500-9.
5. Jansen JO, Yule SR, Loudon MA. Investigation of blunt abdominal trauma. *BMJ* 2008; 336: 938-42.
6. Gaines BA, Ford HR. Abdominal and pelvic trauma in children. *Crit Care Med* 2002; 30: s416-23.
7. Gary R. Fleisher, Stephen Ludwig, Fred M. Henretig, et al. Abdominal Trauma. In: Gray Fleisher (editor). *Textbook of Pediatric Emergency Medicine*. New York (NY): Ovid; 2013. P.1453.
8. Centers for Disease Control. National Center for Injury Prevention and Control. Fatal Injury mapping. [Internet]. [Updated 2014 Feb, Cited 2014 November 21]. Available from: <http://www.cdc.gov/ncipc/osp/charts.htm>
9. Potoka DA, Saladino RA. Blunt abdominal trauma in the pediatric patient. *Clin Pediatr Emerg Med* 2005; 6: 23-31.
10. Sivit CJ. Imaging children with abdominal trauma. *AJR Am J Roentgenol* 2009; 192: 1179-89.
11. Robert Reardon. Ultrasound in trauma. [Internet]. [Updated 2008 May, Cited 2014 October 16]. Available from: <http://www.sonoguide.com/FAST.html>
12. Schmidt B, Schimpl G, Hollwarth ME. Blunt liver trauma in children. *Pediatr Surg Int* 2004; 20: 846-50.
13. Stylianos S. Compliance with evidence-based guidelines in children with isolated spleen or liver injury: a prospective study. *J Pediatr Surg* 2002; 37: 453-6.
14. Clark P, Letts M. Trauma to the thoracic and lumbar spine in the adolescent. *Can J Surg* 2001; 44: 337-45.
15. Robben S. Diagnostic Imaging in Child Abuse Non Accidental Trauma. *Am J Roentgenol Kleinman* 2006; 155: 703.

16. Legome EL, Geibel J, Keim SM, Salomone JP, Udeani J, Dunn E, et al. Blunt Abdominal Trauma. [Internet]. [Updated 2014 Aug 19, Cited 2014 Dec 22]. Available from: <http://www.emedicine.medscape.com/article/1980980-overview>
17. Sivit CJ. Imaging children with abdominal trauma. *AJR Am J Roentgenol* 2009; 192: 1179-89.
18. Sivit CJ. Pediatric abdominal trauma imaging: Imaging choices and appropriateness. *MedScape* 2013; 42: 8-13
19. Stylianos S. Outcomes from pediatric solid organ injury: role of standardized care guidelines. *Curr Opin Pediatr* 2005; 17: 402-6.
20. Visrutaratna P, Na-Chiangmai. Computed tomography of blunt abdominal trauma in children. *Singapore Med J* 2008; 49: 352-8.
21. Scott BH, Bjerke JS, Kaplan LJ, Talavera F, Sheridan RL, Zamboni P, et al. Splenic Rupture. [Internet]. [Updated 2014 Nov 24, Cited 2014 Dec 1]. Available from: <http://www.emedicine.com/med/topic2792.htm>.
22. Abdominal Trauma. [Internet]. [Updated 2013 Nov 5, Cited 2014 Jul 28]. Available from: <http://www.trauma.org/index.php>
23. Mark B, Robert P, Thomas J. *The Merck Manual of Diagnosis and Therapy*. 18<sup>th</sup> ed. New Jersey: Merck Research Laboratories, 2006. P. 157.
24. Kawashima A, Sandler CM, Corl FM, West OC, Tamm EP, Fishman EK, et al. Imaging of renal trauma: a comprehensive review. *RadioGraphics* 2001; 21: 557-74.
25. Noel A., Armenakas, et al. *The One-Page Merck Manual of Health*. [Updated 2007, Cited 2014 Aug 12]. Available from: <http://www.merckmanuals.com/professional/index.html>
26. Ruzinkó V, Willner P, Oláh A. Pancreatic injury from blunt abdominal trauma in childhood. *Acta Chir Belg* 2005; 105: 283-6.
27. Montesano G, Zanella L, Favetta U, Del Bono P, Voccia L, Rossi FS. Rupture of the pancreatic isthmus due to blunt abdominal trauma. *Chir Ital* 2009; 61: 123-6.
28. Cotton BA, Beckert BW, Smith MK, Burd RS. The utility of clinical and laboratory data for predicting intraabdominal injury among children. *J Trauma* 2004; 56: 1068-74.
29. Davies DA, Ein SH, Pearl R, Langer JC, Traubici J, Mikrogianakis A, et al. What is the significance of contrast “blush” in pediatric blunt splenic trauma? *J Pediatr Surg* 2010; 45: 916-20.
30. Vaccaro JP, Brody JM. CT cystography in the evaluation of major bladder trauma. *RadioGraphics* 2000; 20: 1373-81.
31. Committee on Trauma. *Advanced Trauma Life Support (ATLS)*. 8<sup>th</sup> ed. Chicago (Ill): American College of Surgeons, 2008. P. 478.

32. Committee on. Advanced pediatric Trauma Life Support (APLS). recognition of critically ill child. 4<sup>th</sup> ed. European resuscitation council. 2011. P. 135-40.
33. Ellis DY, Harris T, Zideman D. Cricoid pressure in emergency department rapid sequence tracheal intubations: a risk-benefit analysis. *Ann Emerg Med* 2007; 50: 653-65.
34. MCNETT M. A Review of the Predictive Ability of Glasgow Coma Scale Scores in Head-Injured Patients. *J Neurosc Nurs* 2007; 39: 68-75.
35. Bouillon B, Kanz KG, Lackner CK, Mutschler W, Sturm J. The importance of Advanced Trauma Life Support (ATLS) in the emergency room. *Unfallchirurg* 2004; 107: 844–50. [Article in German]
36. Carmont MR. The advanced trauma life support course: a history of its development and review of related literature. *Postgrad Med J* 2005; 81: 87–91
37. Shirley P. Cricoid pressure for emergency airway management. *Emerg Med Australas* 2006; 18: 99.
38. Mattu A, Goyal D, Jeffrey BW, Broder J, Michael D, Deblieux P, et al. Emergency medicine: avoiding the pitfalls and improving the outcomes. Malden, Mass: Blackwell Pub. BMJ Books, 2007; 60.
39. Holmes JF, Offerman SR, Chang CH, Randel BE, Hahn DD, Frankovsky MJ, et al. Performance of helical computed tomography without oral contrast for the detection of gastrointestinal injuries. *Ann Emerg Med* 2004; 43: 120-8.
40. Christiano JG, Tummers M, Kennedy A. Clinical significance of isolated intraperitoneal fluid on computed tomography in pediatric blunt abdominal trauma. *J Pediatr Surg* 2009; 44: 1242-8.
41. Udeani J, Salomone JA, Keim SM. Blunt Abdominal Trauma Clinical Presentation. [Internet]. [Updated 2013 Oct 21, Cited 2014 Sep 12]. Available from: <http://emedicine.medscape.com/article/1980980-clinical#a0256>
42. Hankin AD, Baren JM. Should the digital rectal examination be a part of the trauma secondary survey? *Ann Emerg Med* 2009; 53: 208-12.
43. Potoka DA, Saladino RA. Blunt Abdominal Trauma in the Pediatric Patient. *Clin Ped Emerg Med* 2005; 6: 23-31.
44. Fuller MJ. Applied radiology. [Internet]. [Updated 2010 Jul 10, Cited 2014 May 1]. Available from: <http://www.wikiradiography.com>
45. Rose JS. Emergency medicine clinics of North America. Ultrasound in blunt abdominal trauma 2004; 22: 58-599
46. Baumann B, McCans K, Stahmer S, Leonard MB, Shults J, Holmes WC. Volumetric bladder ultrasound performed by trained nurses increases catheterization success in pediatric patients. *AJEM* 2008; 26: 18-23.

47. Boulanger BR, Kearney PA, Brenneman FD, Tsuei B, Ochoa J. Utilization of FAST (focused assessment with sonography for trauma) in 1999: results of a survey of North American trauma centers. *Am Surg* 2000; 66: 1049-55.
48. James CR, Royse AG. Ultrasound in trauma. *Best Pract Res Clin Anaesthesiol* 2009; 23: 343-62.
49. Walsh N. Ultrasound is effective screen in pediatric trauma. *American College of Emergency Physicians. ACEP News.* 2007.
50. Rose JS, Bair AE, Mandavia D, Kinser DJ. The UHP ultrasound protocol: a novel ultrasound approach to the empiric evaluation of the undifferentiated hypotensive patient. *Am J Emerg Med* 2001; 19: 299-302.
51. Hecht C, Wilkins J. Physics and image artifacts. In: Ma OJ, Mateer JR, Blaivas M (eds). *Emergency Ultrasound.* New York, NY: McGraw-Hill Companies, Inc, 2008. 49-63.
52. Razi NB, Giathini D. Focused assessment with sonography for trauma. *Ultrasound Clin* 2008; 4: 23-31.
53. Rippy J, Royse A. Ultrasound in trauma. *Best Pract Res Clin Anaesthesiol* 2009; 23: 343-62.
54. Gracias VH1, Frankel H, Gupta R, Reilly PM, Gracias F, Klein W, et al. The role of positive examinations in training for the focused assessment sonogram in trauma (FAST) examination. *Am Surg* 2002; 68: 1008-11.
55. Gracias VH1, Frankel HL, Gupta R, Malczynski J, Gandhi R, Collazzo L, et al. Defining the learning curve for the Focused Abdominal Sonogram for Trauma (FAST) examination: implications for credentialing. *Am Surg* 2001; 67: 364-8.
56. Potoka DA, Schall LC, Gardner MJ, Stafford PW, Peitzman AB, Ford HR. Impact of pediatric trauma centers on mortality in a statewide system. *J Trauma* 2000; 49: 2372-45.
57. Hom J. The risk of intra-abdominal injuries in pediatric patients with stable blunt abdominal trauma and negative abdominal CT. *Acad Emerg Med* 2010; 17: 469-75.
58. Livingston H, Lavery RF, Passannate MR, Skurnick JH, Fabian TC, Fry DE, et al. Admission or observation is not necessary after a negative abdominal computed tomographic scan in patients with suspected blunt abdominal trauma. Results of a prospective multi-institutional trial. *J Trauma* 2008; 44: 273-80.
59. Nastanski F, Cohen A, Lush SP, DiStante A, Theuer CP. The role of oral contrast administration immediately prior to the computed tomographic evaluation of the blunt trauma victim. *Injury* 2001; 32: 545-9.
60. Sivit CJ. Imaging children with abdominal trauma. *AJR Am J Roentgenol* 2009; 192(5): 1179-89.

61. Radhiana H, Azian A. Computed Tomography (CT) Imaging of Injuries from Blunt Abdominal Trauma. *Malays J Med Sci* 2010; 17: 29–39.
62. Stylianos S. Outcomes from pediatric solid organ injury: role of standardized care guidelines. *Curr Opin Pediatr* 2005; 17: 402-6.
63. Mettler FA, Wiest PW, Locken JA, Kelsey CA. CT scanning: patterns of use and dose. *J Radiol Prot* 2000; 20: 353-9.
64. Brenner DJ, Hall EJ. Computed tomography: an increasing source of radiation exposure. *N Engl J Med* 2007; 357: 2277-84.
65. Shing CH, Hung CC. Pneumoperitoneum on supine abdominal X-ray: case report and review of the radiological signs. *Hong Kong J Emerg Med* 2005; 12: 46-9.
66. Hsu JM, Joseph AP, Tarlinton LJ, Macken L, Blome S. The accuracy of focused assessment with sonography in trauma (FAST) in blunt trauma patients, experience of an Australian major trauma service. *Injury* 2007; 38: 71-5.
67. American College of Surgeons Committee on Trauma. Pediatric Trauma. In: *Advanced Trauma Life Support for Doctors*. 9<sup>th</sup> ed. Chicago, IL: American College of Surgeons, 2012. P. 260-3.
68. Committee on. Advanced pediatric Trauma Life Support (APLS). recognition of critically ill child. 3<sup>rd</sup> ed. European resuscitation council. 2010. P. 10-6.
69. Leslie E, Geoffrey J, James M. Statistical analysis. In: *Interpretation and uses of medical statistics*. 4<sup>th</sup> ed. Oxford: Scientific Publications (pub). 1991. P.411-6.
70. Kirkpatrick LA, Feeney BC. A simple guide to IBM SPSS statistics for version 20.0. Student ed. Belmont, Calif.: Wadsworth, Cengage Learning, 2013. P.115.
71. Orak M, Ustündağ M, Güloğlu C, Gökdemir MT, Erdoğan MO, Al B. Clinical importance of ultrasonographic pelvic fluid in pediatric patients with blunt abdominal trauma. *Ulus Travma Acil Cerrahi Derg* 2010; 16: 155-9.
72. Menaker J, Blumberg S, Wisner DH, Dayan PS, Tunik M, Garcia M, et al. Intraabdominal Injury Study Group of the Pediatric Emergency Care Applied Research Network (PECARN). Use of the focused assessment with sonography for trauma (FAST) examination and its impact on abdominal computed tomography use in hemodynamically stable children with blunt torso trauma. *J Trauma Acute Care Surg* 2014; 77: 427-32.
73. Faruque AV, Qazi SH, Khan MA, Akhtar W, Majeed A. Focused abdominal sonography for trauma (FAST) in blunt paediatric abdominal trauma. *J Pak Med Assoc* 2013; 63: 361-4.
74. Karam O, Sanchez O, Wildhaber DE, La Scala GC. National Survey on Abdominal Trauma Practices *European Journal Pediatric Surgeon*. *Eur J Pediatr Surg* 2010; 20: 334–8.

75. Van Schuppen J, Olthof DC, Wilde JC, Beenen LF, Van Rijn RR, Goslings JC. Diagnostic accuracy of a step-up imaging strategy in pediatric patients with blunt abdominal trauma. *Eur J Radiol* 2014; 83: 206-11.
76. Ameh EA, Chirdan LB, Nmadu PT. Blunt abdominal trauma in children: epidemiology, management, and management problems in a developing country. *Pediatr Surg Int* 2000; 16: 505-9.
77. Mir M, Bali B, Mir R. Etiological And demographic profile of pediatric abdominopelvic in kashmir. *Internet J Surg* 2012; 28: 150-2.
78. Holmes JF, Mao A, Awasthi S, McGahan JP, Wisner DH, Kuppermann N. Validation of a prediction rule for the identification of children with intra-abdominal injuries after blunt torso trauma. *Ann Emerg Med* 2009; 54: 528–33.
79. Browning JG, Wilkinson AG, Beattie T. Imaging paediatric blunt abdominal trauma in the emergency department: ultrasound versus computed tomography. *Emerg Med J* 2008; 25: 645-8.
80. Schonfeld D, Lee LK. Blunt abdominal trauma in children. *Curr Opin Pediatr* 2012; 24: 314-8.
81. Letton RW Jr, Worrell V, Tuggle DW, American Pediatric Surgical Association Committee on Trauma Blunt Intestinal Injury Study Group. Delay in diagnosis and treatment of blunt intestinal perforation does not adversely affect prognosis in the pediatric trauma patient. *J Trauma* 2010; 68: 790-5.
82. Patel JC, Tepas JJ 3rd. The efficacy of focused abdominal sonography for trauma (FAST) as a screening tool in the assessment of injured children. *J Pediatr Surg* 1999; 34: 44-7.
83. Cotton BA, Beckert BW, Smith MK, Burd RS. The Utility of Clinical and Laboratory Data for Predicting Intraabdominal Injury Among Children. *J Trauma* 2004; 56: 1068–75.
84. Sokolove PE, Kuppermann N, Holmes JF. Association between the “seat belt sign” and intra-abdominal injury in children with blunt torso trauma. *Acad Emerg Med* 2005; 12: 808-13.
85. Holmes JF, Sokolove PE, Brant WE, Palchak MJ, Vance CW, Owings JT, et al. Identification of children with intra-abdominal injuries after blunt trauma. *Ann Emerg Med* 2002; 39: 500-9.
86. Chandler CF, Lane JS, Waxman KS. Seatbelt sign following blunt trauma is associated with increased incidence of abdominal injury. *Am Surg* 1997; 63: 885-8.
87. Holmes JF, Sokolove PE, Brant WE, Palchak MJ, Vance CW, Owings JT, et al. Identification of children with intra-abdominal injuries after blunt trauma. *Ann Emerg Med* 2002; 39: 500-9.

88. Holmes JF, Lillis K, Monroe D, Borgianni D, Kerrey BT, Mahajan P, et al., Pediatric Emergency Care Applied Research Network (PECARN). Identifying children at very low risk of clinically important blunt abdominal injuries. *Ann Emerg Med* 2013; 62: 107-16.e2.
89. Karam O, La Scala G, Le Coultre C, Chardot C. Liver function tests in children with blunt abdominal traumas. *Eur J Pediatr Surg* 2007; 17: 313–6.
90. Ritchie AH, Williscroft DM. Elevated liver enzymes as a predictor of liver injury in stable blunt abdominal trauma patients: case report and systematic review of the literature. *Can J Rural Med* 2006; 11: 283-7.
91. Antonio Muñiz, Adam Vella, Paula Whiteman et al. Evaluation and management of pediatric abdominal trauma. *Pediatr Emerg Med Pract* 2008; 5: 7-12.
92. Bahner D, Blaivas M, Cohen HL, Fox JC, Hoffenberg S, Kendall J, et al. AIUM practice guideline for the performance of the focused assessment with sonography for trauma (FAST) examination. *J Ultrasound Med* 2008; 27: 313–8.
93. Dean AJ, Breyer MJ, Ku BS, Mills AM, Pines JM. Emergency ultrasound usage among recent emergency medicine residency graduates of a convenience sample of 14 residencies. *J Emerg Med* 2010; 38: 214–20.
94. Scaife ER, Fenton SJ, Hansen KW, Metzger RR. Use of focused abdominal sonography for trauma at pediatric and adult trauma centers: a survey. *J Pediatr Surg* 2009; 44: 1746–9.
95. Bixby SD, Callahan MJ, Taylor GA. Imaging in pediatric blunt abdominal trauma. *Semin Roentgenol* 2008; 43: 72–82.
96. Eppich WJ, Zonfrillo MR. Emergency department evaluation and management of blunt abdominal trauma in children. *Curr Opin Pediatr* 2007; 19: 265-9.
97. Mehmet Selim Nural, Türker Yardan, Hakan Güven. Et al. Diagnostic value of ultrasonography in the evaluation of blunt abdominal trauma. *Diagn Interv Radiol* 2005; 11:41-44.
98. Stengel D, Bauwens K, Sehouli J, Porzsolt F, Rademacher G, Mutze S, et al. Systematic review and meta-analysis of emergency ultrasonography for blunt abdominal trauma. *Br J Surg* 2001; 88: 901–12.
99. Coley BD, Mutabagani KH, Martin LC, Zumberge N, Cooney DR, Caniano DA, et al. Focused abdominal sonography for trauma (FAST) in children with blunt abdominal trauma. *J Trauma* 2000; 48: 902-6.
100. Holmes JF, Brant WE, Bond WF, Sokolove PE, Kuppermann N. Emergency department ultrasonography in the evaluation of hypotensive and normotensive children with blunt abdominal trauma. *J Pediatr Surg* 2001; 36: 968–73
101. Soudack M, Epelman M, Maor R. Experience with focused abdominal sonography for trauma (FAST) in 313 pediatric patients. *J Clin Ultrasound* 2004; 32: 53–61.

102. Soundappan SV, Holland AJ, Cass DT, Lam A. Diagnostic accuracy of surgeonperformed focused abdominal sonography (FAST) in blunt paediatric trauma. *Injury* 2005; 36: 970–5.
103. Holmes JF, Gladman A, Chang CH. Performance of abdominal ultrasonography in pediatric blunt trauma patients: a meta-analysis. *J Pediatr Surg* 2007; 42: 1588–94.
104. Sivit CJ. Abdominal trauma imaging: imaging choices and appropriateness. *Pediatr Radiol* 2009; 39: 158–60.
105. Scaife ER, Rollins MD. Managing radiation risk in the evaluation of the & pediatric trauma patient. *Semin Pediatr Surg* 2010; 19: 252–6.
106. Brenner DJ, Hall EJ. Computed tomography: an increasing source of radiation exposure. *N Engl J Med* 2007; 357: 2277–84.
107. Cook SH, Fielding JR, Phillips JD. Repeat abdominal computed tomography & scans after pediatric blunt abdominal trauma: missed injuries, extra costs, and unnecessary radiation exposure. *J Pediatr Surg* 2010; 45: 2019–24.
108. Brenner D, Elliston C, Hall E, Berdon W. Estimated risks of radiation-induced fatal cancer from pediatric CT. *AJR Am J Roentgenol* 2001; 176: 289-96.
109. Mueller DL, Hatab M, Al-Senan R, Cohn SM, Corneille MG, Dent DL, et al. Pediatric radiation exposure during the initial evaluation for blunt trauma. *J Trauma* 2011; 70: 724-31
110. Frush DP, Frush KS. The ALARA concept in pediatric imaging: building bridges between radiology and emergency medicine: consensus conference in imaging safety and quality for children in the emergency setting. *Pediatr Radiol* 2008; 38(Suppl 4): S629-32.
111. Weishaupt D, Grozaj AM, Willmann JK, Roos JE, Hilfiker PR, Marincek B. Traumatic injuries: imaging of abdominal and pelvic injuries. *Eur Radiol* 2002; 12: 1295–311.
112. Holmes J. The risk of intra-abdominal injuries in pediatric patients with stable blunt abdominal trauma and negative abdominal computed tomography. *Acad Emerg Med* 2010; 17: 469–75.
113. Streck CJ Jr, Jewett BM, Wahlquist AH, Gutierrez PS, Russell WS. Evaluation for intra-abdominal injury in children after blunt torso trauma: can we reduce unnecessary abdominal computed tomography by utilizing a clinical prediction model? *J Trauma Acute Care Surg* 2012; 73: 371-6.
114. Hershkovitz Y, Zoarets I, Stepansky A, Kozler E, Shapira Z, Klin B, et al. Computed tomography is not justified in every pediatric blunt trauma patient with a suspicious mechanism of injury. *Am J Emerg Med* 2014; 32: 697-9.
115. Fenton SJ, Hansen KW, Meyers RL, Vargo DJ, White KS, Firth SD, et al. CT scan and the pediatric trauma patient – are we overdoing it? *J Pediatr Surg* 2004; 39: 1877–81.

116. Brenner D, Elliston C, Hall E, Berdon W. Estimated risks of radiation-induced fatal cancer from pediatric CT. *AJR Am J Roentgenol* 2001; 176: 289-96.
117. Melniker LA, Leibner E, McKenney MG, Lopez P, Briggs WM, Mancuso CA. Randomized controlled clinical trial of point-of-care, limited ultrasonography for trauma in the emergency department: the first sonography outcomes assessment program trial. *Ann Emerg Med* 2006; 48: 227-35.
118. Fox JC, Boysen M, Gharahbaghian L, Cusick S, Ahmed SS, Anderson CL, et al. Test characteristics of focused assessment of sonography for trauma for clinically significant abdominal free fluid in pediatric blunt abdominal trauma. *Acad Emerg Med* 2011; 18: 477-82.
119. Taş F, Ceran C, Atalar MH, Bulut S, Selbeş B, Işık AO. The efficacy of ultrasonography in hemodynamically stable children with blunt abdominal trauma: a prospective comparison with computed tomography. *Eur J Radiol* 2004; 51: 91-6.
120. Emery KH, McAneney CM, Racadio JM, Johnson ND, Evora DK, Garcia VF. Absent peritoneal fluid on screening trauma ultrasonography in children: a prospective comparison with computed tomography. *J Pediatr Surg* 2001; 36: 565-9.
121. Terry BM, Blehar D, Gaspari R, Maydell A, Bezuidenhout F, Andronikou S. FAST as a predictor of clinical outcome in blunt abdominal trauma: original article. *SA Journal of Radiology* 2011; 15: 108-10.
122. Retzlaff T, Hirsch W, Till H, Rolle U. Is sonography reliable for the diagnosis of pediatric blunt abdominal trauma? *J Pediatr Surg* 2010; 45: 912-5.
123. Paris C, Brindamour M, Ouimet A, St-Vil D. Predictive indicators for bowel injury in pediatric patients who present with a positive seat belt sign after motor vehicle collision. *J Pediatr Surg* 2010; 45: 921-4.

obeykandl.com

## الملخص العربي

تعتبر الحوادث هي السبب الرئيسي للمرض والوفيات في مرحلة الطفولة. البطن هو الموقع الثاني الأكثر شيوعاً للإصابة. الآلية الأكثر شيوعاً لإصابة في البطن هي حوادث السيارات.

ولها الأولوية ويجب التعامل مع أي مريض وفقاً لتوجيهات الإنعاش المتقدم للإصابات واستقرار الحالة العامة للمريض لسرعة ودقة التشخيص واتخاذ القرارات التي تعتمد على نتائج الفحص الكليني والفحوصات المعملية ونتائج الأشعات التي يحتاجها المريض.

ان الموجات الصوتية علي البطن في اصاباتها الرضية الآن تحل محل التشخيص عن طريق الغسيل البريتوني. لأنه وسيلة تشخيصية سريعة ويمكن أن يتم عمله أثناء عمليق الإنعاش والفحص المبدئي.

الأشعة المقطعية هي الطريقة الأمثل لفحص مريض إصابة البطن الرضية إذا كانت الحالة العامة للمريض مستقرة. وقد يكون هناك حاجة لحقن الصبغة سواء عن طريق الحقن الوريدي أو عن طريق الفم أو الحاجة إلي التخدير. ولا يعتمد قرار التدخل الجراحي عليه بشكل كبير في المقام الأول إلي اتخاذ قرارات مثل مدة الاستشفاء، وشدة الرعاية، وطول تقييد النشاط.

وقد أجريت هذه الدراسة المستقبلية على جميع المرضى الذين يعانون من إصابة رضية في البطن الذين تم قبولهم في المستشفى الجامعي الرئيسي بالإسكندرية من خلال غرفة الطوارئ في الفترة من الأول من مايو ٢٠١٣ حتي الثلاثون من أكتوبر ٢٠١٣.

الهدف من هذه الدراسة هو تحديد الجدوى وتقييم دقة الموجات الصوتية والأشعة المقطعية، والكشف عن الإصابات داخل البطن في الأطفال المصابين بإصابات البطن الرضية.

وخلال فترة الدراسة قد شملت ١٥٠ مريضا تحت سن ١٨ عاما تعرضوا لإصابات في البطن. ١٠٩ ذكور، ٤١ إناث مع متوسط العمر ٩ سنوات (مجموعة ٣ أيام - ١٧ سنة). وكان ٥٤٪ من الحوادث المرورية على الطرق، وذكرت ٣٩.٣٪ من سقوط ارتفاع. (٤٨.٧٪ من الأطفال لم يعانون من نزيف داخل البطن، وحالة واحدة فقط عانت من نزيف داخلي بسيط جدا).

وقد ساعد الفحص بالموجات الصوتية في اتخاذ قرار معالجة المريض حيث تم معالجة ٥٠ مريضا عانوا من نزيف داخلي بعلاج تحفظيا رغم وجود إصابات بالأحشاء الداخلية لهم وخروج ٦٩ مريض إلي المنزل بعد الفحص الأولي و ٥٠ حالة تمت معالجاتها جراحيا وتوفيت ٦ حالات لإصابة شديدة بالمخ. وقد كان فاصل الثقة حوالي ٩٥٪

خضع ٣٨ طفلا للأشعة المقطعية حيث كان الطحال العضو الأكثر إصابة يليه الكبد. ومن خلال مقارنة الموجات الصوتية بالأشعة المقطعية من حيث الوقت حيث كان متوسط الوقت للموجات الصوتية ١١.٢٩ دقيقة مقابل ٥٤.٧٢ دقيقة للأشعة المقطعية. وقد أظهرت الموجات الصوتية حساسية بلغت ٩٥.٢٪ وخصوصية ٩٥.١٢٪ والقيمة التنبؤية الايجابية ١٠٠٪ والقيمة التنبؤية السلبية ١٠٠٪.

وترجع نتائجنا الموجات الصوتية كوسيلة تشخيصية سريعة حساسة آمن ومتنقلة لفرز الأطفال الذين يعانون من إصابات رضية في البطن واللجوء إلي الأشعة المقطعية كمقياس متقدم في حالات مختارة لتقييم متقدم لدرجة الإصابة حيث أنه جهاز مكلف ويستهلك الوقت ويتطلب الكثير من التسهيلات التي قد لا تتوفر بقسم الطوارئ المزدهم.

دور الموجات فوق الصوتية والاشعة المقطعية فى مجال الاصابات الرضية للبطن فى الاطفال  
بقسم الطوارئ

رسالة علمية

مقدمة إلى كلية الطب- جامعة الإسكندرية  
إستيفاء للدراسات المقررة للحصول على درجة

الماجستير

فى

طب الطوارئ

مقدمة من

سعاد السيد مصطفى محمود حسين

بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٨

٢٠١٥

دور الموجات فوق الصوتية والاشعة المقطعية فى مجال الاصابات الرضية للبطن فى الاطفال  
بقسم الطوارئ

مقدمة من

سعاد السيد مصطفى محمود حسين  
بكالوريوس الطب والجراحة - جامعة الإسكندرية، ٢٠٠٨

للحصول على درجة

الماجستير

فى

طب الطوارئ

موافقون

.....

.....

.....

.....

لجنة المناقشة والحكم على الرسالة

أ.د. / عادل محمد رزق  
أستاذ الأشعة التشخيصية  
كلية الطب  
جامعة الإسكندرية

أ.د. / صابر محمد وهيب  
أستاذ الجراحة  
قسم جراحة الأطفال  
كلية الطب  
جامعة الإسكندرية

أ.د. / هشام على بدوى  
أستاذ الأشعة التشخيصية  
معهد البحوث الطبية  
جامعة الإسكندرية

أ.د. / جلال محمد أبو النجاه  
أستاذ الجراحة العامة  
قسم جراحة الأورام  
كلية الطب  
جامعة الإسكندرية

## لجنة الإشراف

أ. د. / حبشى عبد الباسط الحمادى  
أستاذ الجراحة العامة  
كلية الطب  
جامعة الإسكندرية

أ. د. / عادل محمد رزق  
أستاذ الأشعة التشخيصية  
كلية الطب  
جامعة الإسكندرية

أ. د. / صابر محمد وهيب  
أستاذ الجراحة، قسم جراحة الأطفال  
كلية الطب  
جامعة الإسكندرية