

## LIST OF FIGURES

Figure		Page
(1)	Normal femoral attachment of the ACL on proton density MR imaging. ..	2
(2)	Normal ACL bundle anatomy on oblique coronal plane. ....	3
(3)	Normal PCL on sagittal proton density MR imaging in extension and flexion. ....	4
(4)	Superior view of the menisci .....	5
(5)	Sagittal gradient echo image of the ligament of Humphrey located anterior to the PCL.....	6
(6)	Sagittal PD-weighted fat-saturated image of the superior and inferior popliteomeniscal fascicles.....	6
(7)	GRE coronal image demonstrating a discoid lateral meniscus.....	8
(8)	T2-weighted fat-saturated sagittal image demonstrating a flounce like appearance to the posterior horn and body of the medial meniscus .....	8
(9)	Sagittal intermediate-weighted fat-saturated fast SE images.....	10
(10)	Axial short inversion time inversion recovery image obtained in the knee accurately depicts a focal full-thickness chondral defect.....	10
(11)	Sagittal 2D fast SE images obtained with various techniques in the same knee as fig (2) .....	11
(12)	Axial 2D T2-weighted fast SE image provides excellent depiction of a focal region of deep cartilaginous delamination .....	12
(13)	Axial 2D T2-weighted non-fat suppressed fast SE image provides excellent contrast between cartilage surfaces and synovial fluid, but the cartilage is poorly depicted with diffuse low signal intensity.....	12
(14)	Coronal 3D intermediate-weighted fast SE source image obtained with isotropic spatial resolution shows no cartilage abnormalities.....	13
(15)	Coronal 3D SPGR image shows intact highsignal-intensity cartilaginous surfaces in the medial and lateral tibiofemoral compartments .....	14
(16)	Coronal T2-weighted fat-suppressed image shows large areas of complete loss of cartilage in the medial tibiofemoral compartment with associated edematous degenerative lesions in subchondral bone marrow .....	15
(17)	Sagittal 3D DEFT image demonstrates a cartilage fissure at the medial tibial plateau. The defect is outlined by the high signal intensity of synovial fluid. ....	16

<b>Figure</b>	<b>Page</b>
(18) Sagittal T2-W FS fast SE image and water excitation DESS image accurately depict a full-thickness cartilaginous defect at the patella. ....	17
(19) Sagittal 3D water excitation bSSFP source image shows a focal partial-thickness cartilaginous defect at the medial femoral condyle. ....	18
(20) Sagittal source and coronal reformatted 3D VIPR-SSFP images exhibit a high cartilage SNR and high contrast among cartilage, synovial fluid, and other joint structures. ....	19
(21) Partial-thickness tear of the ACL on sagittal fat-suppressed proton density MR imaging .....	20
(22) Partial-thickness ACL tear on corresponding axial T2-weighted MR imaging in same patient (Fig 20). ....	21
(23) Full-thickness tear of the ACL on sagittal fat suppressed T2-weighted MR imaging. ....	21
(24) Near full-thickness tear of the proximal ACL on fat-suppressed proton density MR imaging. ....	21
(25) Bone contusions caused by partial-thickness ACL tear.....	22
(26) Deep sulcus sign on sagittal fat-suppressed T2-weighted MR imaging .....	23
(27) Segond fracture. ....	23
(28) High-grade partial-thickness ACL tear.....	23
(29) Anteriorly subluxed tibia with respect to the posterior aspect of the femoral condyle. ....	24
(30) A hooked, or J-shaped, PCL .....	24
(31) Chronic full-thickness ACL tear on sagittal fat suppressed proton density MR imaging .....	25
(32) Partial-thickness PCL tear.....	26
(33) Full-thickness PCL tear and posterior subluxation of tibia with respect to femur .....	26
(34) Chronic PCL tear on sagittal fat-suppressed T2-weighted MR imaging.....	27
(35) Coronal FSE T2-weighted fat-saturated image demonstrating medial extrusion of the body of the medial meniscus.....	28
(36) Sagittal GRE image demonstrating a horizontal tear (arrow) of the posterior horn of the medial meniscus .....	30

<b>Figure</b>	<b>Page</b>
(37) Horizontal tear.....	30
(38) GRE image demonstrating a peripheral vertical longitudinal tear of the posterior horn of the medial meniscus .....	31
(39) FSE T2-weighted fat-saturated image demonstrating a radial tear of the posterior horn of the medial meniscus near the posterior meniscal root attachment .....	31
(40) Vertical parrot-beak tear .....	32
(41) FSE proton density-weighted fat-saturated sequence demonstrating a bucket-handle tear of the medial meniscus .....	33
(42) FSE T2-weighted fat-saturated images demonstrating bucket-handle tears of both menisci, with fragments from both meniscal bodies flipped into the intercondylar notch. ....	33
(43) FSE proton density-weighted fat-saturated image demonstrating a tear of the lateral meniscus, with the fragment of the body flipped adjacent to the anterior horn .....	34
(44) Proton density-weighted fat-saturated images demonstrating a flap of meniscal tissue .....	34
(45) GRE image demonstrating truncation of the body of the medial meniscus (truncated triangle sign), with the flap of meniscal tissue flipped under the meniscus into the inferior recess medially .....	35
(46) FSE proton density-weighted fat-saturated image demonstrating a tear of the posterior medial meniscal root .....	38
(47) FSE proton density-weighted fat-saturated image demonstrating fluid signal and widening between the medial edge of the body of the medial meniscus and the MCL, consistent with meniscocapsular separation .....	39
(48) GRE sagittal image demonstrating a perimeniscal cyst adjacent to the posterior horn of the medial meniscus, with the tear extending to the cyst. ....	40
(49) 40-year-old woman with acute knee injury.....	40
(50) Diagrams of the knee in extension (a) and flexion (b) show anterior, isometric, and over-the-top positions of the femoral tunnel site in ACL reconstruction. (Adapted, with permission, from reference 11.) .....	42
(51) Diagram shows correct positioning of the tunnels in the coronal plane. In the left knee, the femoral tunnel should open superiorly at the 1–2-o’clock position. ....	43

<b>Figure</b>	<b>Page</b>
(52) Plain x-ray showing anterior position of the tibial tunnel and degenerative osteoarthropathy of the knee joint.....	53
(53) Plain x-ray showing irregularity of the antero-inferior surface of the patella with related faint soft tissue calcification.....	53
(54) Plain x-ray normal position of the tibial and femoral tunnels. ....	55
(55) Plain x-ray showing dislodged femoral tunnel screw .....	55
(56) 25 year old male patient complaining of knee pain 1.5 years following ACL reconstruction. ....	57
(57) 31 year old male patient complaining of knee pain and limitation of movement 2 years following ACL reconstruction. ....	58
(58) Different cases showing abnormal tunnel position .....	59
(59) 24 year old male patient complaining of pain and giving way 5 years post ACL reconstruction. ....	60
(60) 31 year old male patient complaining of knee pain 2 years following ACL reconstruction. ....	61
(61) 25 year old male patient complaining of knee pain 3 years following ACL reconstruction. ....	62
(62) 38 year old male patient complaining of knee pain and locking 6 months following ACL reconstruction .....	63
(63) 31 year old male patient complaining of knee pain 2 years following ACL reconstruction.....	64
(64) 28 year old male patient complaining of knee pain, locking and giving away 1 year following ACL reconstruction.....	65
(65) 33 year old male complaining of pain and giving way 2 years after ACL reconstruction.....	67
(66) 31 year old male patient complaining of knee pain 1 year following ACL reconstruction.....	68
(67) 24 year old male patient complaining of pain and giving way 5 years post ACL reconstruction.....	69
(68) 29 year old female patient complaining of knee pain 6 months following fixation of avulsed ACL tibial attachment.....	70
(69) 64 year old male patient complaining of knee pain 2 years following partial medial meniscectomy .....	72

<b>Figure</b>	<b>Page</b>
(70) 45 year old male patient complaining of knee pain 1 year following partial medial meniscectomy .....	<b>73</b>
(71) 52 year old female patient complaining of knee pain 2 years following medial meniscus repair.....	<b>74</b>
(72) 52 year old female patient complaining of knee pain 2 years following medial meniscus repair. ....	<b>76</b>
(73) 50 year old male patient with partial medial meniscectomy complaining of knee pain following recent knee twisting .....	<b>77</b>
(74) 65 year old male patient complaining of knee pain with previous partial medial meniscectomy long time ago.....	<b>78</b>
(75) 45 year old male patient complaining of knee pain 1 year following partial medial meniscectomy .....	<b>79</b>
(76) 38 year old male patient complaining of knee pain 1 year following partial lateral meniscectomy .....	<b>80</b>
(77) 45 year old female patient complaining of knee pain 1 year following medial meniscus repair.....	<b>81</b>
(78) 49 year old female patient with previous partial lateral meniscectomy complaining of knee pain. ....	<b>82</b>
(79) Different patients with incidental finding of Baker’s cyst following meniscal surgery. ....	<b>83</b>
(80) 45 year old male patient complaining of pain 3 years post Mosaicplasty ...	<b>85</b>
(81) 32 year old male patient complaining of pain 1 year post Mosaicplasty .....	<b>86</b>
(82) 45 year old male patient complaining of pain 3 years post Mosaicplasty ...	<b>87</b>

## LIST OF ABBREVIATIONS

<b>2D</b>	Two dimensional
<b>3D</b>	Spin echo
<b>3D GRE</b>	Three dimensional gradient echo
<b>ACL</b>	Anterior cruciate ligament
<b>AMB</b>	Anteromedial bundle
<b>bSSFP</b>	Balanced steady state free precession
<b>CNR</b>	Contrast to noise ratio
<b>DEFT</b>	Driven equilibrium Fourier transform
<b>DESS</b>	Dual echo steady state
<b>FEMR</b>	Fluctuating equilibrium MR
<b>FFE</b>	Fast field echo
<b>FIESTA</b>	Fast imaging employing steady-state acquisition
<b>FISP</b>	True fast imaging with steady-state precession
<b>FLASH</b>	Fast low angle shot
<b>FSE</b>	Fast spin-echo
<b>GRE</b>	Gradient echo
<b>IDEAL</b>	Iterative decomposition of water and fat with echo asymmetry and list-squares estimation
<b>MCL</b>	Medial collateral ligament
<b>MR</b>	Magnetic resonance
<b>PCL</b>	Posterior cruciate ligament
<b>PLB</b>	Posterolateral bundle
<b>SNR</b>	Signal to noise ratio
<b>SPGR</b>	Spoiled gradient recalled echo
<b>STIR</b>	Short time inversion recovery
<b>TE</b>	Echo time
<b>TR</b>	Repetition time
<b>VIPR</b>	Vastly undersampled isotropic projection reconstruction