

## CHAPTER VIII

### SUMMARY AND CONCLUSIONS

The study area represents the central part of the northern Eastern Desert, occupying the area between the northern scarps of the El Galala El Bhariya plateau to the southern scarp of Gabal Ataqa to the north. It is bounded by latitudes  $29^{\circ} 30' - 30^{\circ} 00' N$  and longitudes  $32^{\circ} 00' - 32^{\circ} 30' E$ . The main wadis in the study area downstream into the coastal plain and the Gulf of Suez, these are from north to south, Wadi Hagoul, Wadi Beda, Wadi Akheider and Wadi Ghoweiba.

The western side of Gulf of Suez is one of the proposed promising sites for anew urbanization programs industrial national projects. As the scope of the study is to build up a geoenvironmental database for the area in a Geographic Information System (GIS) and to give the suitable recommendations to the development strategy of the area under investigation with respect to all available data.

It also aims to study the benefits of the integration between remote sensing and geophysical information in correlation between surface and subsurface structure to delineate the active faults. Geophysical study was carried out for ground water exploration.

In the study area different rock units were identified. They range in age from Carboniferous to Quaternary. Eocene, Oligocene and Miocene are the most widespread rock units in this area.

Carboniferous rocks consist of alternating thick beds of sandstone; crossed-bedded sandstone and dark green shale.

Jurassic unit consists of varicolored, cross-bedded sandstones with mudstone and siltstone interbeds.

The Cretaceous succession in the study area is classified into three rock units, which are (from base to top) the Malha, Galala, and chalky limestone unit. The Eocene rocks are the nummulitic limestones which form the main parts of the Gabal Ataqa and Galal as well as the faulted blocks of Akheider- Rammlyia and Um Zeita-Kahallya. The Eocene succession is subdivided into from base to top; the upper part of the Esna Shale Formation, the Farafra Formation, the Thebes Formation, the Muweilih Formation, the Mokattam Formation, the Observatory Formation, the Qurn Formation, the Wadi Garawi Formation and the Wadi Hof Formation. The Oligocene rocks are differentiated into two units; the lower unit is varicolored, unstratified sands, gravels, and sedimentary quartzites; the upper unit crop out in the central part of the study area and consists of basalt sheets of Gabal Ahmer Formation.

The Miocene succession exposed in the Sadat area that lays 30 km to the south west of Suez city, is subdivided as follows from base to top: Sadat Formation (Early Miocene), Hommath Formation (Middle Miocene) and Hagul Formation (Late Miocene). Recent deposits are represented by sands, gravels, clays, sabkha and sand accumulation.

Landsat Enhance Thematic Mapper (ETM+) data have been used to study geological and lineament analysis using the ERDAS imagines version 9.1. Two scenes (Landsat ETM+ data) covering the investigated area have been geometrically corrected and radiometrically balanced.

Digital image processing of Landsat (ETM+) for the study area generated several products ranging from single band images, false color composite images (7,4,2 in RGB) respectively, principal component analysis (PC1, PC2, PC3, in R,B,G), and ratio images (5/7, 5/4, 3/1), (5/7, 3/1, 4/3) and (5/7, 5/1, 4) in (RGB).

False color composite images (bands 7, 4, 2) should fair to good contrast between the Tertiary basalts and the sedimentary rocks. The near infrared bands 7 and 3 indicate high reflectance in the iron zones and also indicate low reflectance in band 1. Typically, the more iron rich rocks are slightly brownish. Thus the Tertiary basalts are indicated by dark brown color, the limestone beds are whitish brownish, the Wadi deposits are whitish to whitish blue color. The man made areas are dark blue color.

The false color composite Landsat ETM+ images (7, 4, 2 in RGB) are suitable for regional tectonic structure and provide an excellent base map in which rock units are easily discriminated and geologic structures (mainly faults) are highly inferred. Also they show brightness and fair drainage pattern.

In the present study following ratio images were used: (5/7, 3/1, 4/3), (5/7, 5/4, 3/1) and (5/7, 5/1, 4) in (R, G, and B) respectively. Ratio images (5/1, 5/7, 4) were used for lithological discrimination of different rock types.

The Cretaceous rocks are indicated by dark green color, the Eocene sediments are indicated by green to whitish green color the Oligocene rocks are indicated by orange color, the Pliocene sediments are indicated by whitish green color, the Miocene sediments are indicated by whitish blue color and Quaternary old terraces are indicated by yellowish color. The man made areas are purple color.

The Landsat ETM+ false color composite image bands 7, 4, 2 in RGB and the band ratios Landsat ETM+ images with the guidance of the geological map of EGSMA, scale of 1:100,000, to redraw the boundaries of the different rock units producing a new geological map of the study area.

The magnetic data are calculated and convert the total intensity magnetic data into reduced to the magnetic pole by aero-service company and the values of the reduced to the pole data (RTP) are replotted on a base maps and contoured with contour interval 10 gammas (nano tesla).

The description of the reduced to the magnetic pole map shows the high magnetic anomaly occupying the northeastern, northwestern and eastern parts of the study area, while the low magnetic anomaly occupying the southwestern part. Also, there are several high and low magnetic anomalies, which directed N-E trend, which are respectively for the common structural trend in the study area. The map reveals different magnetic gradients at the eastern and southern parts of NE- SW trend and at the northeastern part of NW-SE trend.

The reduced to the magnetic pole map are separated into residual and regional anomalies. The separation was carried out at wave number 0.0000287 (1/km) on the magnetic map reduced to the pole (Geosoft programs, 1994). Two maps are produced representing the two separated pattern features (regional and residual) of the magnetic map reduced to the pole.

The Mag Mode interactive software (Geosoft programs, 1994) used to determine the depth source parameters such as depth, half width and magnetic susceptibilities on twenty selected profiles distributed through out the consider area.

The results of the depth determination are used for the depth map construction using Oasis Montaj software. The depth map of the upper surface for the basement complex reveals that the depth of the basement surface is deeper at the southern part where the depth reach to 3500 m, also the depth of the basement surface is deeper at the northwestern part where the depth reaches to 3200 m. The central part of the study area is occupying by shallow depths of the basement surface where the depth is 1200 m.

In the present study, the GM-sys program (1999) was used to make the magnetic modeling along five magnetic profiles crossing the magnetic anomalies and covering the study area, P1, P2, P3, P4 and P5 are W-E trend using magnetic susceptibility of 0.00229 cgs units for the basement rocks.

The qualitative interpretation for the total intensity magnetic map show many high and low magnetic anomalies which can be resulting from up thrown for the basement blocks according to the fault elements which can dissect the study area. These fault elements have trends N-S, NE-SW, NW-SE and E-W directions.

In the present study, geoelectrical methods are carried out by Schlumberger configuration to detect the subsurface geologic section by constructing the geoelectric cross sections and also, to determine the thicknesses of the different lithologic layers through the subsurface section. Added, constructing the contour maps of the top surface of the different layers, isopah maps for the different subsurface layers

Seventeen vertical electrical soundings are measured in the area enclosed between Lat. 29° 38' - 29° 45' N and Long. 32° 12' - 32° 21' E. The well known Schlumberger configuration of electrode separation ranging from  $AB / 2 = 1.5$  m to  $AB / 2 = 1500$  m used for measuring 17 VES stations. The aim of the

quantitative interpretation of the vertical electrical soundings is to determine the thicknesses and true resistivity values of the successive strata below the different stations.

The quantitative interpretation of the vertical electrical soundings is carried out by using IPI2WIN program. The final results used for construction four geoelectrical cross-sections which exhibits the different geological units represented in the study area. The isopach maps are constructed to indicate the variation of the thicknesses of the first, second, third, fourth and fifth units through the studied area. The second layer represent the fresh water aquifer in the study area for Upper Miocene age the lithology of this layer consists of limestone and sandstone, its thickness rang from 5 to 65m. Also the iso-resistivity maps of the six main layers show considerable variation in the resistivity values ranged between 1 Ohm.m in the Fifth geoelectrical unit to 8700 Ohm.m in the frist geoelectrical unit.

The study area is mainly controlled by faults and limited folds. After tracing all lineaments from the landsat image, their orientations and lengths were determined and measured, then analyzed by preparing histogram diagram for analysis, interpretation and comparison with geologic map and geophysical trend patterns. The results indicate that the f1, f2 at the southern part and f3 at the north western of the study area which detected from geological and magnetic map have the same extension which indicates that these faults are active faults (risk area). Also the shape of the shore line of the western side of the Gulf of Suez controlled by the subsurface faults. The fault system has trends NE that affected on the major part and NNW trend which affected on the small part of the shore line in the studied area.

Data analysis using GIS environment includes two processes (overlapping, and intersecting,). In this study the intersect processes used to be correlated between the surface faults from geological map and subsurface faults from magnetic map. The analysis includes the analysis of drainage basins, shaded hill, slope, and digital elevation model DEM. Considering the risks of the common natural disasters like earthquakes which is represented by the epicenter that is detected in the area, fault zones, active fault and flooding; a risk map of the studied area is constructed. The overlying of the risk map over the classification map resulted in the production of land suitability map of the studied area.

GIS analysis has been applied in three chapters in this thesis, and the result is generating a land suitability map of the area depending upon the database which has been available. This land suitability map shows that area is of great importance in sustainable development. It has a fantastic shore, well whether and archeological places for both internal and external tourism. The mountains and the wadies are a suitable for safari tourism.

**Recommendation:**

- I recommend for drilling the four boreholes in the study area for extraction the fresh water for drinking, domestic and agriculture.
- Avoided any constructions and engineering purposes in the risk areas because it consisting of tracks of the flood plains and active faults.
- To avoid the destructive impacts of the flooding in the studied area and to get benefits of the flooding water as it is urgently needed dams in the upstream as well at the mouths of each basin have to be constructed

## References

- Abdallah, M. A. (1993):** Structural geology of the area between El Galala El-Baharyia and Gabal Akheider. Ph.D. Thesis, Fac. Sci., Ain Shams Univ., Cairo 199p.
- Abdallah, A. M. and Abd El-Hady, F. M. (1966):** Geology of Sadat area, Gulf of Suez. U. A. R. J. Geol., 10(1): 1-24.
- Abdallah, A.M. and Adindani, A. (1963):** Stratigraphy of Upper Paleozoic rocks, western side of the Gulf of Suez. - Mem. Geol. Surv., 25: 18 P.
- Abdl El Shafy, E. and Abd El Moneim, M. (1992):** Lithostratigraphy of the Miocene rocks in the area between Gabal Ataqa and the Northern Galala, Eastern Desert, Egypt: Bull. Fac. Sci., Zagazig Univ. , 14: 290- 302.
- Abdl Latif, T. A., Galal, G. H., Yousef, A. M. A. (1977):** Structural and Lithological impacts on the groundwater occurrences along Wadi El Naqr – Wadi Bada area. Proc. 15<sup>th</sup> Annu. Meet Egy. Geoph. Soc (EGS), Mar 25-26, 1997, p. 107- 124.
- Abdl-Rahman, M.A. and EL-Etr, H.A. (1979):** The structural pattern of the northern part of the Eastern Desert of Egypt.- Apollo-test Project (ASTP), Summary Science Report, V. II, Earth observations and photography, F. El Baz and D. M. Warner editors, NASA SP-412, Washington, D.C.: 87-96.
- Abou El-Anwar, E. A. and El-Gohary, A. M.(2003):** Petrographical and Geochemical characteristics of Sadat Formation and its Depositional Environment, Sadat Area, West Gulf of Suez, Egypt. J. Sediment. Soc. Egypt, 11: 77-94.
- Abu El-Enain, F. M., Ali, M. M., and Ismail, A.S., (1995):** Petrography, Geochemistry and depositional history of the Eocene rocks in the area between Northern El Galala and Gabal Ataqa, Western Gulf of Suez, Egypt. Annals Geol. Surv. Egypt. Vol. xx , pp. 551-576.

- Abou El-Saoud, W. (2006):** Geoenvironmental studies on new Cairo area and its vicinity, Egypt, using remote sensing data and geographic information systems M. Sc. Thesis Fac. Sci. Al-Azhar univ., 326p.
- Abrams, M. J., Conel, J. E., and Lang, H. R. (1984):** The joint. NASA/ Geostal test case project (USA. AAPG Bookstore) Vol. 1 part 9.
- Aero-Service (1984):** Final operational report of airborne magnetic, radiation survey in the Eastern Desert, Egypt for the Egyptian General Petroleum Corporation. Aero-Service, Houston, Texas, 1984, Six Volumes.
- Akaad and Abdallah (1971):** Contribution to geology of Gebel Ataqa area Ann. Geol. Surv. Egypt, V. 1 p. 21 – 42.
- Bandel, K. and Kuss, J. (1987):** Depositional environment of the Pre- rift sediments, Galala heights (Gulf of Suez, Egypt) - Berliner geowiss. Abh, 78 (A): 1-48.
- Baranov, V. (1957):** A new method for interpretation of aeromagnetic maps. Pseudo-gravimetric anomalies; Geophys. Vol. 22, No. 2, p. 259 – 383.
- Baranov, V. (1975):** Potential fields and their transformation in applied Geophysics. Geoexploration Monographs Series 1- 6, Gebruder, Borntraeger, Berlin – Stuttgart.
- Baranov, V. and Naudy, A. (1964):** Numerical calculation of the formula of reduction to the magnetic pole. Geophys. Vol.29, P. 67 – 79.
- Battacharya, B. K. (1965):** Two dimensional harmonic analysis as tool for magnetic interpretation Geophys. Vol. 30, No. 5, p. 829 – 857.
- Battacharya, B. K. (1967):** Some general properties of potential field in pace and Frequency domains. A review, Geoexploration 5, P. 127-143.

- Battacharya, P.K. and Patra, H.P. (1968):** Direct current geoelectric sounding (principles and interpretation), El-Sevier publishing company, Amsterdam, The Netherlands.
- Burger, H. R. (1992):** Exploration Geophysics of the shallow subsurface, Test book, prentice Hall PTR, Englewood cliffs – New Jersey 07632.
- Cherif, O. H. (1966):** Geology of Sadat area, south-west of Suez. M. Sc. Thesis, Fac. Sci., Ain Shams Univ. Cairo.
- Cherif, O. H. and Yheia, M. A. (1977):** Stratigraphy of the area between Wadi Gemal and Wadi Hommath, Gulf of Suez, Egypt. Egypt J. Geol. 21: 185-203.
- Cowen, D. (1988):** GIS versus CAD versus DBMs, what are the differences. Photogram. Eng. And Rem. Sen., Vol. 54, No. 11 Pp. 1551 – 1555.
- Crippen, E. (1989):** Selection of Landsat TM band and band-Ratio combinations to maximize lithologic information in colour composite displays-Proceed. 7th. Conf Remote Sens. Expl. Geol., Calgary. Alberta, Vol 2, pp 917-921. Ann Arbor, MI
- Darwish, M., Abdlallah, A. and Hassanain, I.M. (1984):** Contributions to the petrography and diagenesis of the Jurassic exposures, Gulf of Suez, Egypt. Geol. Surv. Egypt, 14: 173-192.
- Drury, S. A. (1993):** Image interpretation in geology. 2<sup>nd</sup> ed., 283 p. London (Chapman and hall).
- Egyptian Geological Survey and Mining Authority (EGSMA) (1999):** Geotechnical studies and groundwater exploration for Northwestern part of Gulf of Suez., Internal report, No. 43/99, 101p.
- EL Attaar, A. A. (2003):** Early middle Miocene Echinoids from Sadat Formation, Sadat area, south Gebel Ataqqa NW Gulf of Suez Egypt. Egypt. J pale. Cairo 2003, V. 1, p. 209 – 241.

- EL Azabi, M. H. (2000):** Sedimentary facies, paleoenvironments and facies development of Miocene sequences in the Sadat-Wadi Hagul stretch NW Gulf of Suez and their time correlatives in the adjacent basins Egypt: Fifth Inter. Conf. Geol. of the Arab World, Cairo Univ., p. 1135 – 1174.
- Elbaz, F. and Riad, S. (2002):** Seismic hazard and Seismotectonic of Egypt, Cairo UNESCO office.
- El Gaby, S., List, F., Tehrani, R., (1988):** Geology, evolution and metallogenesis of the Pan-African belt in Egypt. The Pan- African belt of Northeast Africa and adjacent areas. Fried.Vieweg and Sohn, Braunschweig, Wiesbaden. 17-63.
- EL-Ghamery, M.N. (1985):** Geologic development of the Cairo-Suez area. M.Sc. Thesis, Fac. Sci., Al-Azhar Univ.: 188 p.
- EL Heiny, I. (1982):** Neogene stratigraphy of Egypt. Newsl Stratiger. 11: 41-54.
- El-Sawy, K.E. (2005):** Structural and environmental studies on the sedimentary cover along the Red Sea coastal plain between 27°30' and 28° N. Using remote sensing and GIS techniques. Eastern Desert, Egypt. Ph.D Thesis, Al-Azhar Univ. 199 p.
- El Shazly, E. M. (1966):** Structural development of Egypt 14<sup>th</sup> Annual meeting, Geol. Soc. Egypt: 31-38(Abstr.).
- Farag, I.A.M. and Ismail, M.M. (1955):** On the structure of the Wadi Hof area (north-east of Helwan). - Bull. Inst.Desert Egypt, 5(1): 179-192
- Farag, I. A. M. and Ismail, M. M. (1959):** A contribution to the structure of the area east of Helwan. - Egypt Jour. Geol., 3(1): 71-86.
- Foley, E.J. (1941):** Geological survey of the Cairo-Suez district and adjoining areas. Standard Oil Co. of Egypt S.A., Unpublished report, No. 15, 45 p.

- Garfunkel, Z. and Baratov, Y. (1977):** The Tectonics of the Suez rift: Geol. Surv. Israel Bull. No. ZI. Pp. 1-44.
- Geosoft program (1994):** "Geosoft mapping and processing system" geosoft Inc., suit 500, Richmond st. west Toronto, on Canda N5UIV6.
- Ghazala, H. H. and Ibrahim, E. H.(2000):** Aeromagnetic and gravity Study on Zafrana area and the rifting on the Gulf of Suez. 5<sup>th</sup> Inter. Conf. Geol. of the Arab world, Cairo Univ. p. 1051-1064.
- GM - SYS program (1999):** Gravity and Magnetic Modelling, version 3.6, Northwest Geophysical Association, Inc. (NGA), Corvallis, Oregon, U.S.A. 97339.
- Gupta, A. P. (1991):** Remote sensing geology. 356 pp., Berlin Heidelberg (Springer).
- Halsey, G.H. and Gardner,W.C. (1975):** Tectonic analysis of Egypt using ERTS-1 Satellite Data. A lecture delivered at General Petroleum Compony of Egypt, Cairo (1975).
- Hassaan , S. M. , Abdeen , M.M. , El Kazzaaz, Y.A., Yehia , M.A and Attia , G.A. (2007) :** Characterization of the Oligocene sands and gravels , Wadi Ghoweiba , northest Gulf of Suez , Egypt , using spectral signature and principle component analysis of terra Aster Image Egypt J. remote sensing and space sci ., V. 11, pp 73-92.
- Hassaan, S. M. (2008):** Studying geological structures of Ayn-Sokhna area, north Eastern Desert, Egypt, by optimum utilization of data fusion techniques of some satellite image. Ph.D. Thesis, Fac. Sci. Helwan University, 207p.
- Hewaidy, A. A, Azab, M. M. and Farouk, S. (2003):** Ammonite biostratigraphy of the Upper Cretaceous succession in the area west of Wadi Araba, north Eastern Desert, Egypt. Jour. Paleontol., Vol. 3, P 331-359.

## Reference

---

- Hume, W. R. (1921):** Relations of the northern Red Sea and associated Gulf areas to the rift theory. Proceedings of the Geological Society of London, 77: p96- - 101.
- Hunt, G.R. (1980):** Electromagnetic radiation: The communication link in remoter sensing. Chapter 2 in Remote Sensing in geology, Siegal, B. S. and A. R. Gillespie (eds.), p. 5-45. New York: Wiley.
- IPI2 – D Program (1993):** Programs set for ID VES data interpretation, dep. of geophysics, Geological Faculty, Mosco uni. Russia, 119899.
- Ismail (1953):** Stratigraphy and structure of the Wadi Hof area (northeast of Helwan) M. Sc. Thesis, Fac. Sci.,Cairo Univ.
- Ismail, M. M. and Farag, I. A. M. (1960):** Contribution to the stratigraphy of the area east of Helwan (Egypt) Bull. Inst. Desert Egypt (1957), 7(1): 95-134.
- Iskander F. (1943):** Geological survey of the Gharaq El Sultani sheet no. 68/54. Standard oil co. Egypt S.A., Reorot no. 57: 29p.
- Keller,E.A. (2002):** Introduction to environmental geology Prentice Hall, Upper Saddle River, New Jersey, 564 p.
- Kiefer, R. W. and Lillesand, T. M. (1987):** Remote Sensing and image interpretation. 612p. John Wiley and Sons, New York.
- Koefoed, O. (1960):** A generalized Cagniard graph for interpretation of geoelectric sounding data. Geophysical prospecting V. 8, m 3, pp 459- 469.
- Landgrebe. D. A. (1978):** The quantitative approach. Concept and rational. In: Swain, P. H. & Davis, S. M., (eds.). Remote sensing: The quantitative approach, p. 1-20, New York, N Y (Me Graw Hill).
- Lillesand, T. M., and Kiefer, R. W. (1994):** Remote Sensing and image interpretation. 3th ed., 750 pp., New York.

- List, F. K. (1992a):** Basic physics of remote sensing. In P. Bankwitz and F. K. List (eds.), - Proceed, of the 3rd United Nations Int. Training Course on Remote Sens Appl. to Geol. Sci., Potsdam and Berlin -Berliner Geowiss. Abh., D, 5. pp. 21-35, Berlin.
- List, F. K. (1993):** Fundamentals of digital image processing for geologic applications. In F. K. List and P. Bankwitz (eds.), - Proceed of the 4th United Nations Int. Training Course on Remote Sens. Appl. To Geol. Sci., Poisdam and Berlin. -Berliner Geowiss. Abh., D, 5, pp 7-29, Berlin,
- Liu Jian Guo; and McM. Moore, J. (1990):** Hue image RGB colour composition A simple technique to suppress shadow and enhance spectral signature. -Int. J. Remote Sens., Vol 11, No, 8, pp. 1521-1530, London (Taylor and Francis).
- Markis. J., Rihm, R., and Allam, A. (1988):** Some geophysical aspects of the evolution and structure of the crust in Egypt, in the Pan African Belt of Northeast Africa and Adjacent areas. Ed. By S. El Gaby and R. O. Greiling, 345-369, Friedr. Vieweg, Braunschweig, Fedral Republic of Germany.
- Meshref, W. M. (1971):** The application of magnetic trend analysis of the interpretation of Egyptian tectonics. Abstracts of papers presented at the 9<sup>th</sup> Annual Metting of the Geological Society of Egypt, 19-21.
- Meshref, W. M. (1990):** Tectonic framework; The Geology of Egypt (ChapterVII), Ed. Rushdi Said by A. A. Balkema, Rotterdam.
- Meshref, W. M., Abdel Baki, S. H., Abdel Hady, H. M. and Solman, S. M. (1980):** Magnetic trend analysis in northern part of the Arabian-Nubian shield and tectonic implications. Annals of the Geol, Survey of Egypt, X, 939-953.
- Meshref, W. M. and El Sheikh, M. M. (1973):** Magnetic tectonic trend analysis in northern Egypt. Egypt Jour. Geol., 17(2), 179-184.

- Meneisy, M.Y. (1990):** *Vulcanicity.*- In Said, R. (Ed.), *The Geology of Egypt*, Rotterdam, Balkema: 157-172.
- Meneisy, M.Y. and Abdel All, A.Y. (1984):** *Geochronology of Phanerozoic volcanic rocks in Egypt.* - *Ain Shams Sci. Bull.*, 25, 24B.
- Meneisy, M.Y. and Kreuzer, H. (1974):** *Potassium-Argon ages of Egyptian basaltic rocks.*- *Geol. Jb.*, D-9: 21-31.
- Milsom, J. (1989):** *Field Geophysics* printed in Great Britain by Butler and Tanner Ltd, Frome and London.
- Moustafa, A. R., (1996):** *Internal structure and deformation of an accommodation zone in the northern part of the Suez rift.* *J. Sturct. Geol.*, Vol. 18, pp. 93-107.
- Moustafa, A.R. and Abdallah, A.M.A. (1991):** *Structural setting of the central part of the Cairo-Suez district.*-*Mid. East Res. Cent.*, Ain Shams Univ., *Earth Sci. Ser.*, V. 5, pp. 133-145.
- Moustafa, A. R., Yehia, M.A., and Abdel Tawab, S., (1985):** *Structural setting of the area east of Cairo, Maadi and Helwan.* *Mid. East Res. Cent.*, Ain Shams Univ., *Se. Res. Ser.*, Vol. 5, pp. 40-64
- Moustafa, M. E. and Zakir, F. A. (1996):** *New enhancement techniques for azimuthal analysis of lineaments for detecting tectonic trends in and around the Afro-Arabian Shield.* *Int. J. Remote Sensing*, 17 (15), 2923-2943.
- Nakhla, A.M. (1982):** *A geophysical approach of the study of Basement tectonics in northern Egypt (with special emphasis on the Cairo-Suez district).*- *M.Sc. Thesis*, Fac. Sci., Ain Shams.: 213 p.
- National sub committee (1974):** *Miocene rock stratigraphy of Egypt.* *J. Geol.*, V 18, pp. 1 – 69.
- Nettleton, L. L. (1976):** *Gravity and magnetic in oil prospecting,*

## Reference

---

McGraw-Hill Book Company Inc., New York, 322p.

**Neev, D. (1975):** Tectonic evolution of Middle East and the levantine basin (easternmost Mediterranean). *Geology* 3 683-686.

**Rabie S. I.; and Ammar A. A. (1990):** Pattern of the main tectonic trends from remote geophysics, geological structures and satellite imagery, Central Eastern Desert, Egypt. *Int. J. Remote sensing*. vol.11, No. 4, pp 669-673

**Reinhard Kirsch (2006):** Groundwater geophys. A tool for hydrogeology, Springer, pp 489.

**Richards, J. A. (1995):** Remote Sensing Digital Image analysis, an introduction, 2nd edition. 3rd printing, (Springer-Verlag). Berlin Heidelberg Pp. 340

**Robson, D. A. (1971):** The structure of the Gulf of Suez (Clysmic) rift, with special reference to the eastern side. *Journal of the Geological Society of London*, 127: p 247 – 276.

**Rowan, L. C., and Vincent, R. K. (1971):** Discrimination of iron zones using visible and near infrared spectral analysis. *Geol. Soc. America*, V. 3, no. 7, p. 691.

**Sabins, F. F. (1987):** Remote Sensing Principals and interpretation. San Francisco. (W. H. Freeman and Company) 426p.

**Sabins, F. F. (1997):** Remote Sensing Principles and Interpretations. W.H. Freeham and Co., New York, 494p.

**Sadek, H. (1926):** The geography and geology of the district between Gebel Ataqa and El-Galala El-Bahariya (Gulf of Suez). - *Geol. Surv. Egypt*, Paper no. 40, 120 p.

**Sadek, H. (1959):** The Miocene in Gulf of Suez region (Egypt). - *Geol. Surv. Egypt*: 118 p.

**Safei EL-Din, A. (1988):** Geological and Hydrogeological studies on the area between Gebel Ataqa and Northern Galala, Egypt.-

## Reference

---

- Ph.D. Thesis, Fac. Sci., Zagazig Univ.: 271 p.
- Sheffield, C. (1985):** Selecting band combination from multispectral data. Photogramm. Eng. Remote Sens., Bathesda, Maryland, Vol. 6, pp 681-687,
- Said, R. (1960):** Planktonic foraminifera from the Thebes Formation, Luxor, Egypt.- Micropaleont., 6: 277-286.
- Said, R. (1962):** The geology of Egypt - Amsterdam, Elsevier Pub. Co.: 377 p
- Said, R. (1990):** Cenozoic. - In Said, R. (ed.), The Geology of Egypt, Rotterdam, Balkema: 451-486.
- Shukri, N.M. and Akmal, M.G. (1953):** The geology of Gebel El-Nasuri and Gebel El-Anqabiya district. - Bull. Soc. Geograph. Egypte, 26: 243-276.
- Sigaev, N.A. (1959):** The main tectonic features of Egypt.-Egypt. Geol. Surv., Paper No. 39, 26 p.
- Sjogern, B. (1984):** Shallow refraction seismics Text book 1<sup>st</sup> published 1984 by Chapman and Hall Ltd, 11 New Fetter lane, London EC 48 EE.
- Steen, G. (1982):** radiometric age dating and tectonic significance of some Gulf of Suez igneous rocks.- 6th Egypt Gen. Petrol. Corp. Expl. Seminar, Cairo: 19 p.
- Strougo, A. (1985a):** Eocene stratigraphy of the eastern Greater Cairo (Gebel Mokattam-Helwan) area.- Mid. East Res. Cent., Ain Shams Univ., Sc. Res. Ser., 5: 1-39.
- Strougo, A. (1985b):** Eocene stratigraphy of the Giza Pyramids plateau.Mid. East Res. Cent., Ain Shams Univ., Sc. Res. Ser., 5: 79-99.
- Strougo, A. and Boukhary, M.A. (1987):** The Middle Eocene-Upper Eocene boundary in Egypt: Present state of the problem. - Revue De Micropaleontologie, 30(2): 122-127.

- Strougo, A., Faris, M., Tammam, M. and Abul – Nasr, R. (2006):** Miocene microfossils from the type-locality of the Hommath Formation (Sadat area, Suez-Ain Sukhna road) and their biostratigraphic implications. Mid. East Res, Cent., Ain Shams Univ., Earth Sci. Ser., V 20, p. 77-96.
- Tealeb, A. (1979):** Predominant tectonic trends in north Egypt. Bull. No. 189, Helwan Institute of Astronomy and Geophysics, 1-22.
- Telford, W. M., Geldart, L. P., Sheriff, R. E. and Key, D. A. (1976):** Applied Geophysics, Cambridge university press, Cambridge.
- Youssef, M.I. and Abdel-Rahman, M.A. (1978):** Structural map by remote sensing of the area between Gebel Ataqa and the northern Galala plateau, Gulf of Suez region, Egypt.- 10<sup>th</sup> Arab Petroleum Conference, Tripoli, Libya, 135(C-3): 8 p.
- Youssef, M.I., Bassione, M.A. and Cherif, O.H. (1971):** Some stratigraphic and tectonic aspects of the Miocene in the northeastern part of the Eastern Desert. - Bull. Inst. Desert Egypt, LII: 119-155.

دراسات جيوبئية على المنطقة الواقعة غرب خليج السويس باستخدام  
البيانات الجيولوجية و الجيوفيزيقية و الإستشعار من البعد

رسالة مقدمة من

شكرى عبد السميع العزب سليمان

للحصول على درجة الدكتوراه فى فلسفة العلوم

كلية العلوم  
جامعة الأزهر

٢٠١٠

## المخلص العربي

يقدم البحث دراسات جيوبئية باستخدام بيانات الأقمار الصناعية و المعلومات الجيوفيزيائية لمنطقة غرب خليج السويس والمحصورة بين المنحدر الجنوبي من جبل عتاقة شمالا وحتى المنحدر الشمالي من هضبة الجلالة البحرية جنوبا في الجزء الشمالي الشرقي من الصحراء الشرقية. وتمتد المنطقة بين خطي طول ٣٢° - ٣٢٣٠° شرقا وحتى خط عرض ٢٩° - ٢٩٣٠° شمالا.

و تهدف الدراسة إلى تحديد الطبقات الحاملة للمياه الجوفية خاصة في وادي غوبية حيث أن منطقة غرب خليج السويس من المناطق التي تلقى اهتماما من الحكومة خاصة في مجال التعمير و إنشاء المشاريع الكبرى. معرفة التراكيب السطحية و التحتسطحية و عمل خريطة موضح عليها التراكيب الجيولوجية.

أيضا بناء قاعدة بيانات جيوبئية لمنطقة الدراسة على أسس نظم المعلومات الجغرافية ووضع توصيات محددة ومناسبة لإستراتيجية التنمية المستدامة بناء على البيانات المتاحة.

باستخدام الأبحاث السابقة و الدراسة الحقلية أمكن التعرف على الوحدات الصخرية المختلفة والتي تتراوح أعمارها من العصر الكربوني وحتى العصر الرباعي والوحدات الصخرية الأكثر انتشارا تتبع عصور الايوسين و الأوليجوسين والميوسين.

باستخدام المكونات الأساسية للصورة في القنوات (٤، ١، ٢) في الأحداثيات الألوان (أحمر- أخضر- أزرق) على التوالي أمكن التفرقة و الفصل بين الصخور المختلفة الموجودة في منطقة الدراسة.

باستخدام الصور الناتجة من العلاقة النسبية (٧/٥ - ١/٥ - ٤) و العلاقات النسبية (٧/١ - ٢/٤ - ١/٣) في إحدائيات الألوان (أحمر- أخضر- أزرق) أمكن فصل صخور الأوليجوسين (بنفسجي اللون) و صخور البليوسين (أصفر فاتح اللون) و صخور الإيوسين (أخضر اللون) و صخور العصر الرباعي الحديث ( أزرق فاتح اللون).

أما بالنسبة لخريطة المركبة الكلية للمغناطيسية الأرضية فقد تم عمل تصحيح لها بالنسبة للقطب المغناطيسي الشمالي لتلاشى التشوهات الناتجة عن بعد المنطقة عن القطب الشمالي و إرجاع القياسات المغناطيسية كما لو قيست عند القطب المغناطيسي. تم استخدام تقنية القدرة الطيفية مغناطيسيا للوصول إلى مستوى أعماق المصادر المسببة للشاذات المغناطيسية. وباستخدام برنامج (جيوسفت ١٩٩٤) تم فصل المجال المغناطيسي الكلي ورسم خريطتين تمثل الشاذات الضحلة والعميقة. وقد تم التفسير الكمي للمركبة الكلية للمغناطيسية الأرضية المصححة باستخدام برنامج (جيوسفت) ١٩٩٤ لتحديد عمق صخور القاعدة من سطح الأرض وتم إنشاء خريطة كنتورية تبين عمق هذا السطح وقد أوضحت هذه الخريطة وجود سطح صخور القاعدة على أعماق ضحلة في الجزء الأوسط من منطقة الدراسة على عمق (١٢٠٠ متر) ويزداد العمق في الجزء الشمالي الغربي و الجزء الجنوبي ليصل إلى (٣٥٠٠ متر).

تم عمل ١٧ جسة كهربية رأسية في المنطقة المحصورة بين خطى طول ١٢°٣٢ - ٣٢,٢١ شرقا وحتى خط عرض ٢٩ ٢٨° - ٢٩,٤٥ شمالا. باستخدام طريقة توزيع شلمبرجير .

أدى التفسير الكمي للجسات الكهربية الرأسية باستخدام برنامج (IPI٢WIN) إلى إنشاء أربع قطاعات جيوكهربية. وقد أظهرت دراسة هذه القطاعات الجيوكهربية إلى وجود ستة طبقات وهي حصى ورمل وحجر جيرى وحجر رملي يحتوى على المياه الجوفية ويتراوح سمك هذه الطبقة من ٤,٥ متر إلى ٦٧ متر وتتبع عصر الميوسين الأعلى. وحجر جيرى ورمل متداخل مع الطين وحجر جيرى وحجر جيرى طينى وطين ورمال وحجر جيرى. كما تم تشييد خرائط توضح سمك ومقاومة هذه الطبقات.

تمت المقارنة بين التراكيب الجيولوجية الموجودة على الخريطة الجيولوجية و تلك التى تم استقراؤها باستخدام وسائل الإستشعار من بعد ثم مقارنة الملامح التركيبية السطحية بالتراكيب التحتسطحية للحصول على شكل واضح و كامل للوضع الجيولوجى بالمنطقة و قد أوضحت تلك الدراسة أن معظم الملامح التركيبية السطحية تمثل إمتداد لبعض التراكيب التحت سطحية بالمنطقة ( صدوع نشطة ).

- و أخيرا تم الإستفادة من قدرات نظم المعلومات الجغرافية ( برنامج ٩,١ Arc GIS ) لتجميع كل البيانات لإنتاج الخرائط الرقمية وتصحيحها جغرافيا في صورة طبقات معلوماتية أمكن مضاهاتها وإيجاد العلاقات بينها على النحو التالي :
- ١- إخراج نموذج الارتفاعات الرقمية لمنطقة الدراسة والتي من خلالها تم الحصول على خريطة نسب الميول لصخور المنطقة بالإضافة لخريطة اتجاهات الميول والتي تساعد في التعرف على أسلوب ونمط جريان المياه السطحية.
  - ٢- عمل خريطة للوضع التركيبي و استنتاج خريطة للصدوع النشطة و خريطة مخاطر السيول و المخاطر الزلزالية.
  - ٣- إنشاء خريطة توضح أنسب الأماكن المقترحة لحفر آبار المياه .
  - ٤- - إنتاج خريطة استخدام الأراضي التي توضح أماكن التنمية المختلفة.