

CHAPTER 6

REFERENCES

6. REFERENCES

- Abbott, W.S. (1925).** A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.*, 18: 265-267.
- Abdel-Mallick, A. Y.; M. A. A. Abdel-Rahman and G. R. A. Hamam (2003).** Survey of entomopathogenic fungi naturally infecting cereal aphid (Homoptera Aphididae) in Southern Egypt. (Abst.). 9th European meeting of the IOBC/WPRS working Group, Insect pathogens and Entomoparasitic nematodes in cooperation with cost actions 842 and 850 "Entomophothorales & Biocontrol Symbiosis" "Growing research and development, Schlos Salzau, Germany, 23-29 May, 2003 .
- Abd-Rabou, S.; N. Ahmed and H. Badary (2012).** Scale insects (Hemiptera: Coccoidea) infesting apple, apricot, pear trees and their abundant parasitoids in Egypt. *Acad. J. Biol. Sci.*, 5(1): 25-29.
- Abid, H.; M. Yi Tina; Yu. He; L. Ruan and S. Ahmed (2010).** *In vitro* and *in vivo* culturing impacts on the virulence characteristics of serially passed entomopathogenic fungi. *J. Food, Agric. Environ.*, 8 (3& 4): 4 8 1 - 4 8 7.
- Adames, M.; M. Fernández; G. Peña and V. Hernández (2011).** Effects of passages through a suitable host of the fungus, *Metarhizium anisopliae*, on the virulence of acaricide-susceptible and resistant strains of the tick, *Rhipicephalus microplus*. *J. Insect Sci.*, 21: 1-13.
- Ajaykumar, P. and K. R. Kanaujia (2005).** Effect of different grain media on sporulation, germination and virulence of *Beauveria bassiana* against *Spodoptera litura* larvae. *J. Biol. Control*, 19(2): 129-133.
- Ali, A.; C. G. Rojas and M. Torre (2005).** Isolation of dipicolinic acid as an insecticidal toxin from *Paecilomyces fumosoroseus*. *J. Appl. Microbiol. Biotechnol.*, 68: 542– 547.
- Altre, J. A.; J. D. Vandenberg and F. A. Cantone (1999).** Pathogenicity of *Paecilomyces fumosoroseus* isolates to diamondback moth, *Plutella xylostella*: correlation with spore size, germination speed, and attachment to cuticle. *J. Invert. Pathol.*, 73: 332-338.
- Alves, R. T.; P. R. Bateman; J. Gunn; C. Prior and S. R. Leather (2002).** Effects of different formulation of viability and medium term storage of *Metarhizium anisopliae* conidia. *Neotropical Entomol.*, 31 (1): 91-99.

- Amala, U.; T. Jiji and A. Naseema (2012).** Mass multiplication of entomopathogenic fungus, *Paecilomyces lilacinus* (Thom) Samson with solid substrates. J. Biopest., 5(2): 168-170.
- Anitha, K. R. (2007).** Seasonal incidence and management of sucking pest of okra. M. Sc.(Agric) Thesis, Univ. Agric. Sci., Dharwad, Karnataka, India.
- Anonymous (2005).** Efficient input management. The Hindu survey, Indian Agriculture. pp 47-49.
- Ansari, M. A. and T. M. Butt (2011).** Effects of successive subculturing on stability, virulence, conidial yield, germination and shelf-life of entomopathogenic fungi. J. Appl. Microbiol., 110: 1460-1469.
- Asghar, M. (2013).** Virulence of *Beauveria bassiana* and *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) passaged through artificial media and an insect host *Uvarovistia zebra* (Orthoptera: Tettigoniidae), Int. J. Agric Crop Sci., 16: 1147-1152.
- Ayyasamy, R. and P. Baskaran (2006).** Evaluation of various synthetic and grain media for radial growth, sporulation and infectivity of the entomopathogenic fungi *Paecilomyces farinosus*. J. Ecobiol., 19 (5) : 229-233.
- Banu, J. G. and N. Gopalakrishnan (2012).** Development of formulations of a native entomopathogenic fungus, *Lecanicillium lecanii* and testing virulence against mealybug, *Paracoccus marginatus* infesting cotton. Indian J. Plant Prote. 40 (3): 182- 186
- Barranco-Florido, J. E.; R. Alatorre-Rosas; M. Gutierrez-Rojas; G. Viniegra-Gonzalez and G. Saucedo-Castaneda (2002).** Criteria for the selection of strains of entomopathogenic fungi *Verticillium lecanii* for solid state cultivation. Enzyme Microb. Technol., 30:910-915.
- Bekheit, H. K. M. (2005).** Biological control tools and their use in insect magement in Egypt. Egypt. J. Agric. Res., 83(3):1249-1288.
- Bhagat, R. M.; R. B. L. Gupta and C. P. S. Yadava (2003).** Field efficacy of two entomopathogenic fungal formulations against whitegrubs in Himachal Pradesh. Indian J. Entomol., 65 (1): 76-81.
- Bhanu, P. B.; S. Puri and P. K. Singh (2012).** Mass production of entomopathogenic fungi using agricultural products. J. Life Sci., 7(2): 229-232.
- Bhattacharyya, A.; A. C. Samal and S. Kar (2004).** Entomophagous fungus in pest management . News Letter, 5: 2-9

- Boucias, D. G. and J. C. Pendland (1998).** Principles of insect pathology. Kiuwe Academic Publishers, 537 p.
- Brownbridge, M.; A. Adamowicz; M. Skinner and B. L. Parker (1999).** Prevalence of fungal entomopathogens in the life cycle of pear thrips, *Taeniothrips inconsequens* (Thysanoptera : Thripidae) in Vermont sugar maple forests. Biol. Control, 16(1): 54-59.
- Brownbridge, M.; S. Costa and S. T. Jaronski (2001).** Effects of *in vitro* passage of *Beauveria bassiana* on virulence of *Bemisia argentifolii*. J. Invert. Pathol., 77: 280- 283.
- Burges, H. D. (1981).** Microbial Control of Pests and Plant Diseases :1970-1980. Academic Press, London. 949 pp.
- Burgess, H. D. and N.W. Hussey (1971).** Microbial control of pests and plant Diseases, 1970-1980, Academic press, London.
- Burges H. D. and N.W. Hussey (1981).** Microbial Control of Insect Pests and Mite, Academic Press, London, pp. 161-167.
- Butt, T. M; M. Goettel (2000).** Bioassays of entomopathogenic fungi. In: Bioassays of entomopathogenic microbes and nematodes. Navon, A. and K. R. S. Ascher (editors). pp. 141-195. CAB International.
- Butt, T. M.; C. Wang; F. A. Shah and R. Hall (2006).** Degeneration of entomogenous fungi. In: An ecological and societal approach to biological control. Eilenberg, J. H. M. T. Hokkanen (editors). pp. 213–226. Springer
- Carruthers, R. I. and R. S. Soper (1987).** Fungal diseases. In: Epiizootiology of Insect Diseases. Fuxa, J. R. and Tanaday (eds) Wiley. Interscience, New York, pp. 357-416.
- Chandrashekharaiyah, G.; B. Gowda; A. Kumar and A. K. Chakravarthy (2013).** Bio-efficacy of *Verticillium lecanii* against white fly on greengram Insect Environ., 19(2):83-85.
- Cooper, R. and A.W. Sweeney (1986).** Laboratory studies on the recycling potential of the mosquito pathogenic fungus *Culicinomyces clavisporus*. J. Invert. Pathol., 48: 152–158.
- Dai, Y.; H. Hiromori and M. Hatsukade (2006).** Virulence of the entomopathogenic fungus *Beauveria brongniartii* to several life stages of the yellowish elongate chafer *Heptophylla picea* Motschulsky (Coleoptera: Scarabaeidae). Appl. Entomol. Zool., 41 (2): 287–293 .
- Daust, A. R.; G. M. Ward and W. D. Roberts (1983).** Effect of formulation on the viability of *Metarhizium anisopliae* conidia. J. Invert. Pathol., 41:151-160.

- Dent, D. (1991).** Insect Pest Management C.A.B International. ISBB No-85198-666-8) 604 pages.
- Derakhshan, A.; R. J. Rabindra; B. Ramanujam and M. Rahimi (2008).** Evaluation of different media and methods of cultivation on the production and viability of entomopathogenic fungi, *Verticillium lecanii* (Zimm.) Viegas. Pakistan J. Biol. Sci., 11: 1506-1509.
- Derakhshan, A.; R. J. Rabindra and B. Ramanujam (2008).** Effect of storage conditions of formulations on viability of *Verticillium lecanii* (Zimm.) Viegas and its virulence to *Brevicoryne brassicae* (L.). Pakistan J. Biol.Sci., 8 (2): 498-501.
- De-Souza, L.; N. C. De-Oliveira; C. F. Wilcken and A. B. Filho (2004).** Pathogenicity of *Verticillium lecanii* to pine aphid. Rev. Arvore, 28 (5): 765-770.
- Dreistadt, S. H.; J. K. Clark and M. L. Flint (1994).** Pests of landscape trees and shrubs: An integrated pest management guide. Oakland: Univ. Calif. Div. Agric. Nat. Res., 3359.
- Duncan, D. B. (1955).** Multiple range and multiple F tests. Biometrics, 11:1-44.
- Easwaramoorthy, S and S. Jayaraj (1978).** Effectiveness of the white halo fungus, *Cephalosporium lecanii*, against field populations of coffee green bug, *Coccus viridis*. J. Invert. Pathol., 32:88-96.
- Easwaramoorthy, S.; J. Srikanth; G. Santhalakshmi and N. Geetha (2003).** Mass culture and formulation of three entomogenous fungi, with special reference to *Beauveria brongniartii* (Sacc.) patch, against *Holotrichia serrata* F. (Coleoptera : Scarabaeidae). Proceedings of the 64th Annual Convention of the Sugar Technologists Association of India, Cochin, Kerala, 17th-19th August, 2002.
- Ekesi, S.; N. K. Maniania; I. Onu and B. Lohr (1998).** Pathogenicity of entomopathogenic fungi (Hyphomycetes) to the legume flower thrips, *Megalurothrips sjostedtoi* (Trybom) (Thysanoptera : Thripidae). J. Appl. Entomol., 122 (9-10): 629-634.
- Eliana, T. I.; G. V. Pereira; D. T. Miyagui; M. H. P. Pinotti and P. M. O. J. Neves (2007).** Production of extracellular protease by a Brazilian strain of *Beauveria bassiana* reactivated on coffee berry borer *Hypothenemus hampei*. Brazilian Arch. Biol. Technol. J., 50(2):217-223.
- El-Serafi, H. A.; A. A. Ghanim; A. H. El-Heneidy and M. K. El-Sherbenie (2004).** Ecological studies on certain insects infesting guava orchards and their predatory insects at Mansoura District. Egypt. J. Biological Pest Control, 14: (1) 77-85.

- El-Sinary, N. H. and S. A. Rizk (2007).** Entomopathogenic fungus, *Beauveria bassiana* (Bals.) and gamma irradiation efficiency against the greater wax moth, *Galleria melonella* (L.). American-Eurasian J. Sci. Res., 2 (1): 13-18.
- Esfandiari, M.; M. S. Mossadegh and R. Eslamizadeh (2007).** Biology of the cottony cushion scale, *Icerya purchasi* Mask. (Hom. : Margarodidae) in laboratory and its seasonal fluctuations in citrus orchards of Northern Khuzestan. J. Sci. Tech. Agric. Nat. Res., 10 (4B): 393-403.
- Evans, H. C. (1974).** Natural control of arthropods, with special reference to ants (Formicidae), by fungi in The Tropical High Forest of Ghana. J. Appl. Eco. , 11:37-49.
- Evans, H. C. (1999).** Biological control of weed and insect pests using fungal pathogen, with particular reference to Srilanka. Bio control News and Information, 20 (2): 63 -68.
- Ezzat, Y. M. and S. M. A. Nada (1986).** List of super family Coccoidea as known to exist in Egypt. Bull. Lab. Entomol. Agric. 43: 85-90.
- Ezzati-tabrizi, R.; T. Reza, P. Hamid-reza (2009).** Effect of formulating of *Beauveria bassiana* conidia on their viability and pathogenicity to the onion thrips, *Thrips tabaci* Lind. (Thysanoptera: Thripidae) J. Plant Protec. Res., 49 (1):97-104.
- Fadayivata, S.; F.Moravvej and J. Karimi (2014).** Pathogenicity of the fungus *Lecanicillium longisporum* against *Sipha maydis* and *Metopolophium dirhodum* in laboratory conditions, J. of Plant Protec. Res., 54: (1): 67-73.
- Fargues, J. F. and P. H. Robert (1983).** Effect of passaging through scarabaeid hosts on the virulence and host specificity of two strains of the entomopathogenic hyphomycete *Metarhizium anisopliae*. Can. J. Microbiol., 29: 575 – 583.
- Farooq A. S.; C. S. Wang and T. M. Butt (2005).** Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. FEMS Microbiol. Letters, 251: 259–266.
- Farsi, M. J.; H. Askari; K. T. Jahromi and A. K. Pakdel, (2005).** Effect of important ecological factors on blastospore production of *Verticillium lecanii* DAOM 198499 in liquid medium. Iranian J. Agric. Sci., 36: 109-119.
- Ferron, P. (1977).** Influence of relative humidity on the development of fungal infection caused by *Beauveria bassiana* (Fungi imperfecti, Moniliales), in imagines of *Acanthoscelides obtectus* (Coleoptera: Bruchidae). Entomophega, 22: 393-396.

- Ferron, P. (1985).** Fungal control. In: Comprehensive Insect Physiology, Biochemistry and Pharmacology, G.A. Kerkut and L.I. Gilbert [eds.], pp 313-316., Academic, New York.
- Finney, D. N. (1971)** Probit analysis. 3rd Ed, Cambridge Univ. Press London pp: 318.
- Forschler, B. T. and G. L. Nordin (1989).** Impact of *Beauveria bassiana* on the cotton wood borer *Plectrodera sclator* (Coleoptera : Cerambycidae), in a commercial cotton wood nursery. J. Entomol. Sci., 24: 186-190.
- Fransen, J. J.; K. Winkelman and J. C. Van Lenteren (1987).** The differential mortality at various life stages of the greenhouse whitefly, *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae), by infection with the fungus *Aschersonia aleyrodis*(Deuteromycotina: Coelomycetes). J. Invert. Pathol., 50: 158-165.
- Fumio, I.; K. Yaginuma; N. Kobayashi; K. Mishiro and T. Sato (2001).** Screening of entomopathogenic fungi against the brown-winged green bug, *Plautia stali* Scott (Hemiptera: Pentatomidae) Appl. Entomol. Zool., 36 (4): 495–500 .
- Fumio, I.; M. Toyama and T. Sato (2003).** Pathogenicity of *Metarhizium anisopliae* to the chestnut weevil larvae under laboratory and field conditions. Appl. Entomol. Zool., 38 (4): 461–465.
- Fuxa, J. R. (1987).** Ecological considerations for the use of entomopathogens in IPM. Ann. Rev. Entomol., 32: 225-251.
- Ganga Visalakshy, P. N.; A. Manoj kumar and A. Krishnamoorthy (2004).** Evaluation of entomopathogenic fungi *Verticillium lecanii* against *Scirtothrips dorsalis*. Project Directorate of Biological Control, Bangalore, Annu. Progr Repo : 2004-2005, pp. 125-126.
- Garcia, M. F. (2006).** Phytosanitary status of citrus in Spain: Insects and mites. Informatore Fitopatologico, 56(1): 28-31.
- Gerlach, J. (2010).** Margarodidae (Hemiptera: Insecta) of the Seychelles islands. *Phelsuma* 18: 70-73.
- Gindin, G.; N.U.Geschtovt; B. Raccah and I. Barash (2000).** Pathogenicity of *Verticillium lecanii* to different developmental stages of the silverleaf whitefly, *Bemisia argentifolii*. Phytoparasitica, 28 (3): 1-11.
- Godonou, I.; K. R. Green; K. A. Oduro; C. J. Lomer and K. Afreh-Nuaman (2000)** . Field evaluation of selected formulations of *Beauveria bassiana* for the management of the banana weevil

(*Cosmopollites sordidus*) on plantain (*Musa* spp.). International Symposium on Biological Control Agents in Crop and Animal Protection, Swansea, U. K. 24-28 August 1999. Bio control Sci. & Technol., 10 (6): 779-788.

Goettel, M. S., G. D. Inglis and S. P. Wraight (2000). Fungi *In: field manual technique in invertebrate pathology*, L.A. lacey and H.K.kaya (eds.) Kluwer Academic publisher , NetherLands ., pp.255-282 .

Gopalkrishnan, C. and K. S. Mohan (2000). A simple and cost effective *in vitro* method for mass production of conidia of *Nomuraea rileyi*. Pest Mgt. Hort. Ecosyst., 6: 36-39.

Green, E. E. (1907). Notes on the Coccidae collected by the Percy Sladen Trust expedition to the Indian Ocean: supplemented by a collection received from Mr R. Dupont, Director of Agriculture, Seychelles. Trans. Linn. Soc. London, 2 (12): 197-207.

Guedes-Frazzon, A. P.; I. Vaz Junior; A. Masuda; A. Schrank and M. Henning (2000). *In vitro* assessment of *Metarhizium anisopliae* isolates to control the cattle tick *Boophilus microplus*. Vet. Parasitol., 94: 117-125.

Guillon, M. (1997). Mass-production of biopesticides: Scale up and quality assurance. British Crop Production Council Symposium, 68. 1997, Farnham. Insecticides: novelty or necessity: proceedings. Farnham: The British Crop Protection Council., p.151 - 162.

Gullan, P.J. and M. Kosztarab (1997). Adaptations in scale insects. Ann. Rev. Entomol., 42: 23–50.

Hajek, A. E.; R. A. Humber and M. H. Griggs (1990). Decline in virulence of *Entomophaga maimaiga* (Zygomycetes: Entomophthorales) with repeated *in vitro* Subculture. J. Invert. Pathol. 56: 91-97.

Hall, R. A. and B. Papierok (1982). Fungi as biological control agent of arthropods of agricultural and medical importance. Parasitol., 84: 205-240.

Hall, R. A. (1982). Deuteromycetes : Virulence and bioassay design. *In: Invertebrate pathology and microbial control*. Proceedings III International Colloquium Invertebrate Pathology, XV Annual Medium Soc. of Invertebrate Pathology, Brighton, U.K. pp.191-196

Hamam, G. H. (2003). Studies of entomopathogenic fungi of cereal aphids infecting wheat plants in Assiut, Egypt. M.Sc. Thesis, Assiut Univ. Egypt..

Hassan, N. A. and S. G. Radwan (2008). Population dynamics of *Icerya seychellarum*(Homoptera: Margaroididae) and *Rodalia cardinalis*

(Coleoptera : Coccinellidae) on persimmon (*Diospyros kaki*) at Qalubiya Governorate. Egypt. J. Agric. Res., 86(3):1015-1027.

Hatting, J. L., R. A. Humber, T. J. Poprawski and R. M. Miller (1999). A survey of fungal pathogens of aphids from South Africa, with special reference to cereal aphids. Biol. Control, 16 (1): 1–12.

Henderson, C. F. and E. W. Tilton (1955). Tests with acaricides against the brown wheat mite, J. Econ. Entomol., 48:157-161.

Hedgecock, D.; P. M. Moore; P. M. Higgins and C. Prior (1995). Influence of moisture content on on temperature tolerance and storage of *Metarhizium flavoviride* conidia in oil formulation. Biocontrol Sci.Technol., 5(3):371-377.

Hidalgo, E.; D. Moore and G. N. Lepatourel (1998). The effect of different formulations of *Beauveria bassiana* on *Sitophilus zeamais* in stored maize. J. Stored Prod. Res., 34 (2-3) : 171-179.

Hincapie, R. H.; A. Ospina; Y. Bustillo and A. Saldarriaga (1990). Evaluacion del entomopatogeno *Verticillium lecanii* en el control del afido *Mysus persicae* encrisantemos. Revista Colombiana de Entomologia, 16(2): 21-27.

Hsia, I. C. C.; M.T. Islam; Y. Ibrahim; T. Y. Howand and D. Omar (2014). Evaluation of conidial viability of entomopathogenic fungi as influenced by temperature and additive. Int. J. Agric. Biol., 16: 146–152.

Ibrahim, L.; T. M. Butt and P. Jenkinson (2002). Effect of artificial culture media on germination, growth, virulence and surface properties of the entomopathogenic hyphomycete *Metarhizium anisopiae*. Mycol. Res., 106: 705-715.

Ignoffo, C. M.; D. L. Hostetter; P. P. Sikorowski; G. Sutter and W. M. Brooks (1977). Inactivation of representative species on entomopathogenic virus, a bacterium, fungus and protozoa by the ultraviolet light source. Environ Entomol., 6: 411-415.

Ignoffo, C. M.; A. H. McIntosh; C. Garcia; M. Kroha and J. M. Johnson (1982). Effectiveness of successive *in vitro* and *in vivo* passages on the virulence of the entomopathogenic fungus, *Nomuraea rileyi*. Entomophaga, 27: 371-378.

Inglis, G. D.; T. M. Goettel and B. Strasser (2001). Use of hyphomycetous fungi for managing insect pest. In: Fungi as biocontrol agentes. Butt, T.M.; Jackson, C.; Magan, N. (Ed.). Wallingford: CAB International, pp 23-69.

Jana, S. (2009). Influence of different storage conditions on vitality and virulence of *Beauveria bassiana* spores, J.Agrobiol., 26 (2): 75-81.

- Jaronski, S.T. (1986).** Commercial development of deuteromycetous fungi of arthropods: a criterial appraisal. *In: Fundamental and applied aspects of invertebrate pathology.* Samson, R.A., J.M. Vlask, R. Peters, (Ed.). Wageningen: Society of Invertebrate Pathology: 653-656.
- Jayaraj, S. (1989).** Integrated management of coffee green scale *Coccus viridis* (Homoptera: Coccidae). *J. Plantation Crops*, 16: 195-201.
- Jenkins, N. E.; G. Heviefo; J. Langewald; A. J .Cherry and C. J. Lomer (1998).** Development of mass production technology for aerial conidia for use as mycopesticides. *Biocontrol News and Information*, 19: 21-31.
- Jeong, J. K. and K. C. Kim (2008).** Selection of a highly virulent isolate of *Lecanicillium attenuatum* against cotton aphid. *J. Asia-Pacific Entomol.*, 11(1):1- 4.
- Kadir, A. A. and H. S. Barlow (1992).** Pest Management and the Environment in 2000. C.A.B. International, Malaysia, pp. 401.
- Karthikeyan, A. and V. Selvanarayanan (2011).** Studies on the suitability of certain culture media for *Beauveria bassiana* (Bals.) Vuill., and *Verticillium lecanii* (Zimm.) Viegas *J.Res. Biol.*, 7: 508-512.
- Kawakami, K. (1960).** On the change of characteristics of the silkworm muscardines through successive cultures. *Bull. Seric. Exp. Stn. Japan.* (16): 83-99.
- Khalil, S. K.; M. Kahn and M. Naeem (1990).** Studies on entomopathogenic fungi *Verticillium lecanii* (Zimm.) for the control of green pearl aphid, *Myzus persicae* Sulz. *Sarhad J.Agric.*, 6: 597-600.
- Khalil, S. K.; V. Taborsky and J. Bartos (1983).** Studies on *Verticillium lecanii* for the biological control of aphids. In 10th International Congress of Plant Protection at Brighton, England, 2: 20-25.
- Knudsen, G. R.; J. B. Johnson and D. J. Eschen (1990).** Alginate pellet formulation of a *Beauveria bassiana* (Fungi: Hyphomycetes) isolate pathogenic to cereal aphids. *J. Economic Entomol.*, 83 (6): 2225-2228.
- Krishnamurthy, P. N. M.; S. G. Eswara- Reddy (2004).** Survey on potential microbial agent against thrips. *Pest Management in Hortil. Ecosystem*, 7(2) : 117-123.

- Kulat, S. S.; N. N. Zade; L. N. Peshkar and S. S. Varhade (2002).** Influence of different culture media on the growth and sporulation of *Metarhizium anisopliae* (Metsch.) Sorokin. J. Biolo. Control. 16(2): 177-179.
- Kulkarni, N. S. (1999).** Utilisation of fungal pathogen *Nomuraea rileyi* (Farlow) Samson in the management of lepidopterous pests. Ph.D Thesis, Univ. Agric. Sci., Dharwad, India.
- Landa, Z. (1984).** Protection against glasshouse whitefly, *Trialeurodicus vaporariorum* in integrated protection programmes for glasshouse chambers. Sbornik Urtiz Zahoraduichini, 11: 215-218.
- Latgé, J. L.; R. A. Hall; R. I. Cabrera and J. C. Kerwin (1986).** Liquid fermentation of entomopathogenic fungi. In: Fundamental and applied aspects of invertebrate pathology. Samson, R.A., J.M. Vlask, R. Peters (Ed.). Wageningen: Society of Invertebrate Pathology, p.603-606.
- Lawrence, A. L. and D. I. Shapiro (2003).** The potential role for microbial control of orchard insect pests in sustainable agriculture Food, Agric. Environ., 1(2): 326-331.
- Lee, J. C. and D. A. Landis (2001).** Natural enemies in field crops: A guide to biological control. Michigan State Univ. Ext. Bull., 2721.65p.
- Leena, M. D.; S. Easwaramoorthy and R. Nirmala (2003).** *In vitro* production of entomopathogenic fungi, *Paecilomyces farinosus* (Hotmskiold) and *Paecilomyces lilacinus* (Thom.) Samson using by products of sugar industry and other agro industrial by-products and wastes. Sugar Technol., 5 (4): 231-236.
- Lopes, E.; M. I. Lopes and N. M. Barros (1995).** Virulence of stored conidia of *Nomuraea rileyi* against soybean caterpillar, *Anticarsia gemmatilis*. Ciencia Rural, 25: 197-200.
- Machado, A.; A. C. Monteiro; A. M. B. de Almeida and M. E. Martins (2010).** Production technology for entomopathogenic fungus using a biphasic culture system Pesq. Agropec. Bras., Brasília, 45 (10):1157-1163.
- Makoto, A. and T. Ikegami (2005).** Susceptibility of five species of thrips to different strains of the entomopathogenic fungus, *Beauveria bassiana*. Appl. Entomol. Zool., 40 (4): 667-674.
- Mamta, T.; M. Thakur; N. Malik and S. Ganger (2011).** Mass scale cultivation of entomopathogenic fungus *Nomuraea rileyi* using agricultural products and agro wastes J. Biopestic., 4 (2): 176-179.
- Mangoud, A. A. H. (2000).** Integrated pest management of apple trees. Ph.D. Thesis, Fac. Agric. Cairo Univ., Cairo, Egypt
- Mangoud, A. A. H. and H. A. S. Abd El-Gawad (2003).** Evaluation of different integrated pest management concepts for controlling the

Egyptian fluted mealybug, *Icerya aegyptiaca* on the ornamental plants. Bull. Ent. Soc. Egypt, Econ. Ser., 29 : 137-149.

Mangoud, A. A. H.; M. A. Salem and M. A. Abd El-Aziz (2012). Effect of different compounds on *Icerya seychellarum* (Hemiptera: Monophlebidae) and *Rodalia cardinalis* (Coleoptera: Coccinellidae) on mango leaves under laboratory conditions. Egypt. Acad. J. Biolog. Sci., 5(3): 113 -119.

Mangoud, A. A. H.; M. S. Abd El-Wahid and M. A. Abd El-Aziz (2012). Screening test of different compounds against *Icerya seychellarum* and harmful effect on the beneficial predator *Rodalia cardinalis* under laboratory conditions. Egypt, J. Agric. Res., 86 (5): 1861-1875.

Marcos, R. F. and S. P. Wraight (2007). Mycoinsecticides and Mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. Biological Control, 43: 237–256.

Masuda, T. and O. Kikuchi (1992). Pathogenicity of *Verticillium lecanii* isolates to whitefly and aphids. Japanese J. Appl. Entomol. Zool., 36: 239-245.

McCoy, C. W. (1990). Entomogenous fungi as microbial pesticides, *In: New directions in biological control.* R. R. Baker and P. E. Dunn (eds.). Alan R. Liss, New York, NY, 139-159.

McCoy, C. W.; R. A. Samson and D. G. Boucias (1988). Entomogenous fungi, *In: Handbook of Natural Pesticides, Vol.5 microbial insecticides, part A,* C. M. Ignoffo and N. B. Mandava (eds.). CRC Press, Inc., Boca Raton, FL, pp. 151-236.

Meade, D. L. and D. N. Bruce (1991). The use of *Verticillium lecanii* against submarginal instars of *Bemisia tabaci*. J. Invert. Pathol., 57: 296-298.

Mehta, J.; D. J. Kiran; K. Ambika; S. Priya; K. Neha and D. Sakshi (2012). Biomass Production of entomopathogenic fungi using various agro products in Kota Region, India. Res. J. Biol. Sci., 1(4):12-16.

Mesbah, H. A.; K. S. Moursi ; A. K. Mourad; E. A. Zakzouk and R. S. Abdel-Fattah (2012). Ecological studies on the common white mealybug, *Icerya seychellarum seychellarum* (Hemiptera: Monophlebidae) associated with *Dodonia viscosa* in Alexandria, Egypt. Egypt. Acad. J. Biolog. Sci., 5(3): 27-31.

Midwest biological control news online (2014).

C.F: <http://www.entomology.wisc.edu/mbcn/kyf612.html>.

- Milner, R. J. and G. G. Lutton (1986).** Dependence of *Verticillium lecanii* on high humidities for infection and sporulation using *Myzus persicae* as host. *Environmental Entomology*, (15): 380-382.
- Miranpuri, G. S. and G. G. Khachatourians (1995).** Application of *Beauveria bassiana* and *Verticillium lecanii* against Saskatoon berry leaf aphid *Acyrtosiphon macrosiphum*. *J. Insect Sci.*, 8: 93-95.
- Mitsuaki, S. (2004).** A novel technique to inoculate conidia of entomopathogenic fungi and its application for investigation of susceptibility of the Japanese pine sawyer, *Monochamus alternatus*, to *Beauveria bassiana*. *Appl. Entomol. Zool.*, 39 (3): 485-490.
- Mitsuaki, S. (2001).** *Paecilomyces cateniannulatus* Liang, a commonly found, but an unrecorded entomogenous fungus in Japan. *Appl. Entomol. Zool.*, 36 (3): 283-288.
- Mitsuaki, S.; H. Sato and N. Maehara (2002).** Density of the entomopathogenic fungus, *Beauveria bassiana* Vuillemin (Deuteromycotina: Hyphomycetes) in forest air and soil. *Appl. Entomol. Zool.*, 37 (1): 19-26.
- Mohammad, A. H.; S. F. Moussa; A. H. Abo-Ghaila and S. A. Ahmed (2010).** Efficiency of certain insecticides on the population of the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Green) and their natural enemies under the field condition in Ismailia governorate. *Egypt. Acad. J. Biolog. Sci.*, 2 (2): 11- 17.
- Mondal, P. and A. K. Bhattacharya (2004).** Assessment of different media for mass multiplication of entomopathogenic fungus, *Beauveria bassiana* (Balsamo) Vuillemin. *Proc. Nat. Acad. Sci., India Section B, Biol. Sci.*, 74 (2): 161-169.
- Moore, D.; O. K. Douro-kpindou; N. E. Jenkins and C. J. Lomer (1996).** Effects of moisture content and temperature on storage of *Metarhizium flavoviride* conidia. *Biocontrol Sci. Technol.*, 6(1):51-61.
- Mote, U. N.; P. R. Mahajan and J. R. Kadam (2003).** Bioefficacy of *Verticillium lecanii* (Zimm.) Viegas against sucking pests of gerbera in polyhouse. *In: Green Pesticides for Insect Pest Management*. Chennai, pp. 179-183.
- Murphy, B. C.; T. Morisawa and M. P. Parrella (1998).** Insect-killing fungi. *Floriculture's IPM future Grower Talks*, 61: 60 - 68.
- Nada, M. (1990).** The scale insects (Homoptera : Coccoidea) on citrus in Egypt. *J. Pest Cont. Environ. Sci.*, 2:201-209.
- Nada, M.; S. Abd-Rabou and G. E. Hussien (1990).** Scale insects infesting mango trees in Egypt (Homoptera: Coccoidea). *Proc. ISSIS, VI, Part II: 133-134.*

- Nagaich, B. B. (1973).** *Verticillium* species pathogenic on aphids. Indian J. Pathol. Microbiol., 26: 163-165.
- Nagaraja, S. D. (2005).** Effect of formulations of *Nomuraea rileyi* (Farlow) Samson and spray equipments in the management of tobacco caterpillar in groundnut and podborer in chickpea ecosystem. M.Sc. (Agric.) Thesis, University of Agric. Sci., Dharwad, India.
- Nahar, P.; P. Yadav; M. Kulye; A. Hadapad; M. Hassani; U. Tuor; S. Keller; A. G. Chandele; B. Thomas and M. V. Deshpande (2004).** Evaluation of indigenous fungal isolates, *Metarhizium anisopliae* M 34412, *Beauveria bassiana* B3301 and *Nomuraea rileyi* N 812 for the control of *Helicoverpa armigera* (Hub.) in pigeonpea field. J. Biol. Control, 18 (1): 1-8.
- Nankinga, C. M. and D. Moore (2000).** Reduction of banana weevil populations using different formulations of entomopathogenic fungus *Beauveria bassiana*. Biocontrol Sci. Technol., 10(5): 645-657.
- Nelson T. L.; A. Low and T. R. Glare (1996).** Large scale production of New Zealand strains of *Beauveria* and *Metarhizium*. Proceedings of the 49th New Zealand Plant Protection Conf., 257-261.
- Nier, T.; F. Rive and J. C. Bermudez (1993).** First report for Mexico on the isolation of *Verticillium lecanii* from whitefly and *in vitro* pathogenicity tests on this insect. Revista Mexicana De Micologia, 7: 149-156.
- Nirmala, R.; B. Ramanujam; R. J. Rabindra and N. S. Rao (2005).** Growth parameters of some isolates of entomofungal pathogens and production of dust free spores in rice medium. J. Biol. Control, 19 (2): 121-128.
- Nirmala, R.; B. Ramanujam; R. J. Rabindra and N. S. Rao (2006).** Effect of entomofungal pathogen on mortality of three aphid species. J. Biol. Control, 20(1): 89-94.
- Oetting, R. (2004).** Insect and arthropod pest identification and management. A hand-out for Southeast Greenhouse Conference workshops 2000-2004. UGA/CAES/Griffin campus. Available on-line at [http://mrec.ifas.ufl.edu/lso/Manual/insect-textspecies only.htm](http://mrec.ifas.ufl.edu/lso/Manual/insect-textspecies%20only.htm) accessed Aug. 2008 (verified 27 May 2009).
- Olga, M.; M. Kilian; E. C. Oerke and H. Wilhelmdehne (2002).** Oils for increased efficacy of *Metarhizium anisopliae* to control whiteflies, Biocontrol Sci. Technol., 12: 337- 348.
- Orozco Santo, S. M.; J. Farias-Larios; J. Lopez-Perez and N. R. Ramirezvazquez (2000).** Use of *Beauveria bassiana* for the control

of *Bemisia argentifolli* in melon. Maneho-Integrado-de-plagas, 56: 45-51.

Osborne, L. S.; F. L. Pettitt; Z. Landa and K. A. Hoelmer (1994). Biological control of pests attacking crops grown in protected culture: The Florida experience. *In: Pest management in the subtropics: biological control: a Florida perspective*, D. Rosen, F.D. Bennett and J.L. Capinera (eds.). Vol. 1. pp. 327-342. Intercept Limited, Andover, UK.

Ota, M.; A. Ozawa and H. Kobayashi (1999). Efficacy of *Beauveria bassiana* preparation against whiteflies on tomato. Annual Report of the Kanto-Tosan- Plant Protection Soc., 46: 103-112.

Pena, J. E.; R. M. Baranowski and R. E. Litz (1987). Life history, behavior and natural enemies of *Philephedra tuberculosa* (Homoptera: Coccidae). Florida Entomologist, 70(4):421-427.

Pereira, R. M. and D. W. Roberts (1991). Alginate and corn starch mycelial formulations of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae*. J.Econ. Entomol. 84(6): 1657-1661.

Peter, W. I. and M. S. Tigano (2006). Identification and taxonomy of some entomopathogenic *Paecilomyces* spp. (Ascomycota) isolates using rDNA-ITS Sequences Genetics and Molecular Biology, 29(1): 132-136.

Prior, C.; P. Jollnds and G. Le Patourel (1988). Infectivity of oil and water formulation of *B. bassiana* to the cocoa weevil pest J. Invert. Pathol., 52:62-72.

Quesada-Moraga, E. and A. Vey (2003). Intra-specific variation in virulence and *in vitro* production of macromolecular toxins active against locust among *Beauveria bassiana* strains and effects of *in vivo* and *in vitro* Passage on these factors .Biocontrol, Sci. Technol., 13: 323-340.

Quinden, R. J. (1984). The use of *Verticillium lecanii* an entomopathogenic fungus to control of glasshouse whitefly. Proceedings Brighton, England, : 9-12

Rachappa, V. (2003). Occurrence of entomopathogenic fungi and utilization of *Metarhizium anisopliae* in the management of selected crop pests in Northern Karnataka, Ph.D Thesis. Univ. Agric. Sci., Dharwad, India.

Ramarethinam, S.; S. Marimuthu; W. V. Murugesan and S. Lognathan (2000). Evaluation of *Paecilomyces fumosoroseus*, on entomopathogenic fungus for controlling red spider mite,

Oligonychus coffeae (Nietner) (Acarina: Tetranychidae), infesting tea in India. *Pestology*, 24 (9): 1-5.

Ramarethinam, S.; S. Marimuthu and V. Murugesan (2005). Entomopathogenic fungi *Verticillium lecanii* – an overview. *Pestology*, 29 (12): 9-28.

Ramarethinam, S.; S. Marimuthu; S. Loganathan and N. V. Murgesan (2002a). Potentials of entomopathogenic fungal based commercial formulations on some important pests of selected vegetable crops in India. *Pestology*, 26(7): 17-21.

Ramarethinam, S.; S. Marimuthu; W. V. Murugesan and S. Loganathan (2002b). Evaluation of *B. bassiana* against diamond back moth, *P. xylostella* L. (Lepidoptera: Xponomenidae). *J. Entomol. Res.*, 25(4) : 267-271.

Ramarethinam, S.; S. Marimuthu; N. V. Murgesan and S. R. Vishwanathan (2001). Studies on the shelf life and container content compatibility of a commercial *B. bassiana* formulation – Bio-Power®. *Pestology*, 25 (91): 5-12.

Ramegowda, G. K. (2005). Aerobiology, Epizootiology and Utilization of *Nomuraea rileyi* (Farlow) Samson. Ph. D. Thesis, University of Agric. Sci., Dharwad, India.

Ravensberg, W. J. (2010). The development of microbial pest control products for control of arthropods: a critical evaluation and a roadmap to success. Ph.D Thesis Wageningen Univ., Netherlands.

Raymond, J. S. L. S. and B. Shams-Pirzadehi (2000). Lack of host specialization in *Aspergillus flavus* *Appl. Environ. Microbiol.*, 66(1): 320 –324.

Reda F. A. B.; S. F. Mousa; L. S. Hamouda; R. M. Badawy and S. A. Atteia (2012). Scale insects infesting guava trees and control measure of *Pulvinaria psidii* (Hemiptera: Coccidae) by using the alternative insecticides. *Egypt. Acad. J. Biolog. Sci.*, 5(3): 89 -106

Rezk, M. A. A. (2009). Biological control of certain scale insects and mealybugs infesting certain fruit trees. M.SC. Thesis, Alexandria, Univ., Egypt.

Roberts, D.W. (1981). Toxins of entomopathogenic Fungi. *In: Microbial control of pests and plant Diseases 1970-1980.* Burgess, H.D. (ed.). Academic press, London pp 441- 84.

Romback, M. C. (1989). Production of *Beauveria bassina* conidia in submerged culture. *Entomophaga*, 5: 45-52.

Rousson, S.; M. Rainbault and B.K. Lonsane (1983). Zymotics a large scale fermenter design and evaluation. *Appl. Biochem. Biotechnol.*, 42: 161-167 .

- Sahayaraj, K. and S. Karthick (2008).** Mass production of entomopathogenic fungi using agricultural products and by products, *Afr. J. Biotechnol.*, 7 (12): 1907-1910.
- Salman, A. M. A. and M. M. S. Bakry (2012).** Relationship between the rate of infestation with the mealybug, *Icerya Seychellarum* (Westwood) (Margarodidae: Homoptera) and the yield loss of seedy balady mango trees at Luxor governorate. *World Rural Observations*, 4(4): 50-56.
- Santoro, P. H.; P. M. Neves; R. Z. Silvada; S. Akimi and J. Zorzetti (2005).** Produção de esporos de *Beauveria bassiana* (Bals.) Vuill. num processo bifásico utilizando diferentes meios líquidos. *Semina Ciências Agrárias*, 26: 313-320.
- Schaerffenberg, B. (1964).** Biological and environmental conditions for the development of mycoses caused by *Beauveria* and *Metarhizium*. *Insect. Pathol.*, 6: 8-20.
- Scully, L. R. and M. J. Bidochka (2005).** Serial passage of the opportunistic pathogen *Aspergillus flavus* through an insect host yields decreased saprobic capacity. *Can. J. Microbiol.*, 51:185-189.
- Shah, F. A.; C. S. Wang and T. M. Butt (2005).** Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. *FEMS Microbiol. Lett.*, 51: 259 -266.
- Sharma, S.; R. B. L. Gupta and C. P. S. Yadav (1999).** Mass multiplication and formulation of entomopathogenic fungi and their efficacy against white grub. *J. Mycol. Plant Pathol.*, 29 (3): 299-305.
- Sood, A. K.; P. K. Mehta; N. P. Kashyap and R. Lal (2001).** Bioefficacy of *Beauveria bassiana* against diamond back moth, *Plutella xylostella* L. (Lepidoptera : Yponomenidae). *J. Entomol. Res.*, 25 (4): 267-271.
- Sprenkel, R. K. and W. M. Brooks (1977).** Winter survival of the entomogenous fungus *Nomuraea. rileyi* in North Carolina. *J. Invert. Pathol.*, 29: 262-266.
- Sreeramkumar, P.; L. Singh and H. Tabassum (2005).** Potential use of polyethylene glycol in the mass production of non-synnematos and synnematos strains of *Hirsutella thompsonii* Fisher in submerged culture. *J. Biol. Control*, 19 (2) : 105-113.
- Starnes, R. L; C. L. Liu and P. G. Marrone (1993).** History, use and future of microbial insecticides. *American Entomol.*, 39 (2): 83-91.
- Stathers, T. E.; D. Moore and C. Prior (1993).** The effect of different temperatures on the viability of *Metarhizium flavoviride* conidia stored in vegetable and mineral oils. *J. Invert. pathol.*, 62(2):111-115.

Steel, G. D. and J. H. Torrie (1981). Principles and Procedures of Statistics (2nd edition). McGraw-Hill Book Company. Inc. N. Y. 633pp.

Steenberg, T. and R. A. Humber (1999). Entomopathogenic potential of *Verticillium* and *Acremonium* species (Deuteromycotina: Hyphomycetes). J. Invert. Pathol., 73: 309-314.

Steinhaus, E. A. (1949). Principles of Insect Pathology- New York ; McGraw – Hill.

Steinkraus, D. C.; C. J. Geden and D. A. Rutz (1991). Susceptibility of lesser meal worm (Coleoptera: Tenebrionidae) to *Beauveria bassiana* (Moniliales: Moniliaceae): Effects of host stage, substrate, formulation and host passage. J. Med. Entomol., 28: 314-321.

Stimac, L. J.; M. R. Pereira; B. S. Alves and A. L. Wood (1993). Mortality in laboratory colonies of *Solenopsis invicta* (Hymenoptera : Formicidae) treated with *Beauveria bassiana* (Deuteromycetes). J. Econ. Entomol., 86 (1):1083-1087.

Suklova, T. I. (1989). The biological method in the greenhouse. Rashchita Rastenii, 2: 37-38.

Susumu, S. and M. Yamaji (2003). Effect of density of the termite, *Reticulitermes speratus* Kolbe (Isoptera: Rhinotermitidae), on the susceptibilities to *Metarhizium anisopliae*. Appl. Entomol. Zool., 38 (1): 125–130.

Tamizharasi, V.; J. Srikanth and G. Santhalakshmi (2005). Molasses based medium requires no nitrogen supplement for culturing three entomopathogenic fungi. J. Biol. Control, 19(2): 135-140.

Tawfik, M. H. and Z. K. Mohammad (2001). Ecological studies of two scale insects (Hemiptera, Coccoidea) on *Morus alba* in Egypt. Bollettino di Zoologia Agraria e di Bachicoltura, 33 (3): 267-273.

Tounou, K.; H. M. Agboka; K. J. poehling; G. Z. Langewald and C. Borgemeister (2003). Evaluation of the entomopathogenic fungi *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* (Deuteromycotina:Hyphomycetes) for control of the green leafhopper *Empoasca decipiens* (Homoptera: Cicadellidae) and potential side effects on the egg parasitoid *Anagrus atomus* (Hymenoptera: Mymaridae). J. Biocont. Sci. Technol., 13(8): 715-728.

Tsutomu, S. (2005). Preliminary experiments to control the silverleaf whitefly with electrostatic spraying of a mycoinsecticide. Appl. Entomol. Zool., 40 (2): 289–292.

- Tsutomu, S. and K. Sugiyama (2005).** Pathogenicity of three Japanese strains of entomopathogenic fungi against the silverleaf whitefly, *Bemisia argentifolii*. *Appl. Entomol. Zool.*, 40 (1): 169 –172.
- Unruh C. M. and P. J. Gullan (2008).** Identification guide to species in the scale insect *tribe Iceryini* (Coccoidea: Monophlebidae). *Zootaxa*, 1803: 1-106.
- Van Emden, H. F. (1996).** *Pest Control*. Cambridge press 117 pp.
- Van- Hanh, V.; S. Hong and K. Kim (2008).** Production of aerial conidia of *Lecanicillium lecanii* 41185 by Solid-State. *Mycobiol.*, 36 (3): 183-189.
- Vandenberg, J. D.; A. M. Shelton; W. T. Wilsey and M. Rmos (1998).** Assessment to *B. bassiana* sprays for control of diamond back moth (Lepidoptera: Plutellidae) on crucifers. *J.Econ.Entomol.*, 91(3): 624-630.
- Vandenberg, J. D. and F. A. Cantone (2004).** Effect of serial transfer of three strains of *Paecilomyces fumosoroseus* on growth *in vitro*, virulence and host specificity. *J. Invert. Pathol.*, 85: 40–45.
- Vimaladevi, P. S. (1994).** Conidia production of the entomopathogenic fungus *N. rileyi* and its evaluation for control of *Spodoptera litura* on *Ricinus communis*. *J. Invert. Pathol.*, 63: 145-150.
- Vyas, R. V.; D. N. Yadav and R. J. Patel (1991).** Mass production of entomopathogenic fungus, *Beauveria brongniartii* on solid substrates. *Indian J. Expl. Biol.*, 29: 795-797.
- Wasti, S. S. and G. C. Hartman (1975).** Experimental parasitization of larvae of the gypsy moth, *Porthetria dispar* (L) with the entomogenous fungus, *Beauveria bassiana* (Balsamo) Vuill. *Parasitol.*, 70: 341-346.
- Watson, G.W. and J. Kubiriba (2005).** Identification of mealybugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. *African Entomol.*, 13:35-47.
- Westwood, J. O. (1855).** The Seychelles Dorthesia. *Gardeners' Chronicle and Agricultural Gazette* 51: 836.
- Williams, D. J. and G. W. Watson (1988).** The scale insects of the tropical south pacific Region . part 2. The mealy bugs (Pseudococcidae). CAB International Walling, Oxford, UK., 260pp
- Williams, D. J. (2004).** *Mealybugs of southern Asia*. The Natural History Museum, UK.
- Wiwat, C. (2004).** Development of *Nomuraea rileyi* based biopesticide for controlling Lepidopteran larvae. Ph. D. Thesis, Mahidol University, Malaysia.

Wraight, S. P.; R. I. Carruthers; S. T. Jaronski; C. J. Bradley; C. A. Garza and S. Galani-Wraight (2000). Evaluation of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* for microbial control of the silverleaf whitefly, *Bemisia argentifolii*. *Biol. Control*, 17:203-217.

Zare, R. and W.Gams (2001). A revision of *Verticillium* sect. *Prostrata*. III. Generic classification. *Nova Hedwigia*, 72: 329- 337.

Zhang, A. W.; W. Z. Liu; X. Q. Nong; C. S. Deng; W. L. Guo and B. Jang (1992). A trial production of wettable powder of *Beauveria bassiana*. *Chinese J.Biol. Control*, 8 (3): 118-120.