

CONCLUSIONS

CONCLUSIONS

1. The most important and significant data in our study in predicting weaning outcome of COPD patients from mechanical ventilator were integrative weaning index(IWI), rapid shallow breathing index(RSBI), glasgow coma score(GCS), heart rate and presence of abnormal value of resistance. GCS had the best specificity ,positive predictive value and accuracy, while presence of abnormal resistance had the best sensitivity and negative predictive value.
2. There were other data in our study that may predict weaning of COPD patients from mechanical ventilator although they were statistically non significant as they should be studied on larger number of patients. They were presence of structural brain damage, presence of electrolyte disturbance especially hypophosphatemia, presence of taking sedation prior to weaning, presence of acid base disturbance, presence of ischaemic heart disease, presence of heart failure, presence of atrial fibrillation and BMI.
3. Haemoglobin level, systolic blood pressure, respiratory rate, tidal volume, static compliance, maximum inspiratory pressure, P0.1 were poor predictors of weaning outcome of COPD patients from mechanical ventilator in our study.
4. PaO₂, HI, PaCO₂ could not accurately predict weaning outcome of COPD patients from MV.
5. Non invasive PSV gave more than 50% chance of weaning of COPD patients from MV and it can also shorten the duration of weaning from mechanical ventilator.

**LIMITATIONS
OF THE STUDY**

LIMITATIONS OF THE STUDY

1. Lack of continuous ECG monitoring to detect the exact time of occurrence of ischaemic changes to the heart whether at the time of mechanical ventilation, during weaning process or at the time of extubation.
2. Although static compliance can be measured during weaning from mechanical ventilation, it was not an easy task to be performed during the weaning process, because the patient's inspiratory effort during the assisted breath could interfere with the inspiratory plateau pressure measurement. Also it was difficult to give the patient a muscle relaxant or strong sedative during the course of weaning. In our study we minimized this limitation by observing the digital display of the pressure-time inspiratory plateau curve thus avoiding respiratory cycles that revealed clear inspiratory efforts of the patients and sometimes by giving a short acting sedative like propofol.
3. Static compliance could not be measured directly by the ventilator in all cases so, if it could not be measured directly it was calculated using this equation: $C_{st} = VT / (\text{inspiratory plateau pressure} - \text{positive end expiratory pressure})$.
4. We tested the effect of acid base disturbance in general on weaning outcome of patients with COPD from mechanical ventilator without specifying the effect a certain type of acid base disturbance on weaning outcome.

RECOMMENDATION

RECOMMENDATION

1. There were data in our study that may predict weaning of COPD patients from mechanical ventilator although they were statistically non significant as they should be studied on larger number of patients. They were presence of structural brain damage, presence of electrolyte disturbance especially hypophosphatemia, presence of taking sedation prior to weaning, presence of acid base disturbance, presence of ischaemic heart disease, presence of heart failure, presence of atrial fibrillation and BMI. So, we recommend studying these data on larger number of patients in further studies.
2. We recommend studying the effect of certain value of abnormal resistance on weaning outcome of patients with COPD from mechanical ventilator as we have examined in our study the effect of abnormal value of resistance in general (without specifying the value) on weaning outcome of COPD patients from mechanical ventilator.
3. We recommend also studying the effect of a specific subtype of acid base disturbance (metabolic or respiratory, acidosis or alkalosis) on weaning outcome of patients with COPD from mechanical ventilator as we tried in our study to detect the effect of acid base disturbance in general on weaning outcome of patients with COPD from mechanical ventilator.
4. It is advisable to study the effect of certain type alone of heart failure whether it is left sided heart failure, right sided heart failure or diastolic dysfunction on weaning outcome of patients with COPD from mechanical ventilator.

REFERENCES

REFERENCES

1. Esteban A, Frutos F, Tobin MJ, Alia I, Solsona JF, Valverdu I, et al. A comparison of four methods of weaning patients from mechanical ventilation. Spanish Lung Failure Collaborative Group. *N Engl J Med* 1995;332(6):345-50.
2. Brochard L, Rauss A, Benito S, Conti G, Mancebo J, Rekik N, et al. Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1994;150(4):896-903.
3. Esteban A, Alia I, Gordo F, Fernandez R, Solsona JF, Vallverdu I, et al. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. The Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1997;156(2 Pt 1):459-65.
4. Ely EW, Baker AM, Dunagan DP, Burke HL, Smith AC, Kelly PT, et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med* 1996;335(25):1864-9.
5. Esteban A, Alia I, Tobin MJ, Gil A, Gordo F, Vallverdu I, et al. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1999;159(2):512-8.
6. Yang KL, Tobin MJ. A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991;324(21):1445-50.
7. Sassoon CS, Mahutte CK. Airway occlusion pressure and breathing pattern as predictors of weaning outcome. *Am Rev Respir Dis* 1993;148(4 Pt 1):860-6.
8. Jabour ER, Rabil DM, Truwit JD, Rochester DF. Evaluation of a new weaning index based on ventilatory endurance and the efficiency of gas exchange. *Am Rev Respir Dis* 1991;144(3 Pt 1):531-7.
9. Levy MM, Miyasaki A, Langston D. Work of breathing as a weaning parameter in mechanically ventilated patients. *Chest* 1995;108(4):1018-20.
10. MacIntyre NR, Cook DJ, Ely EW, Jr., Epstein SK, Fink JB, Heffner JE, et al. Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College of Critical Care Medicine. *Chest* 2001;120(6 Suppl):375s-95s.
11. Nemer SN, Barbas CS, Caldeira JB, Carias TC, Santos RG, Almeida LC, et al. A new integrative weaning index of discontinuation from mechanical ventilation. *Crit Care* 2009;13(5):R152.
12. Nucci G, Mergoni M, Polese G, Cobelli C, Rossi A. On-line monitoring of respiratory mechanics. In: Nucci G, Mergoni M, Polese G, Cobelli C, Rossi A (eds). *Mechanics of breathing*. Milan: Springer-Verlag; 2002. 327-36.

13. Holliday JE, Hyers TM. The reduction of weaning time from mechanical ventilation using tidal volume and relaxation biofeedback. *Am Rev Respir Dis* 1990;141(5 Pt 1):1214-20.
14. Truwit JD, Marini JJ. Validation of a technique to assess maximal inspiratory pressure in poorly cooperative patients. *Chest* 1992;102(4):1216-9.
15. DeHaven CB, Kirton OC, Morgan JP, Hart AM, Shatz DV, Civetta JM. Breathing measurement reduces false-negative classification of tachypneic preextubation trial failures. *Crit Care Med* 1996;24(6):976-80.
16. Kuhlen R, Hausmann S, Pappert D, Slama K, Rossaint R, Falke K. A new method for P_{0.1} measurement using standard respiratory equipment. *Intensive Care Med* 1995;21(7):554-60.
17. Girard TD, Ely EW. Protocol-driven ventilator weaning: reviewing the evidence. *Clin Chest Med* 2008;29(2):241-52.
18. Martinez A, Seymour C, Nam M. Minute ventilation recovery time: a predictor of extubation outcome. *Chest* 2003;123(4):1214-21.
19. Lee KH, Hui KP, Chan TB, Tan WC, Lim TK. Rapid shallow breathing (frequency-tidal volume ratio) did not predict extubation outcome. *Chest* 1994;105(2):540-3.
20. Epstein SK. Etiology of extubation failure and the predictive value of the rapid shallow breathing index. *Am J Respir Crit Care Med* 1995;152(2):545-9.
21. Tobin MJ, Van de Graaff W. Monitoring of lung mechanics and work of breathing. In: Tobin M (ed). *Principles and practice of mechanical ventilation*. New York: McGraw-Hill; 1994. 967–1003.
22. Jaeschke R, Guyatt GH, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? The Evidence-Based Medicine Working Group. *Jama* 1994;271(9):703-7.
23. Reyes A, Vega G, Blancas R, Morato B, Moreno JL, Torrecilla C, et al. Early vs conventional extubation after cardiac surgery with cardiopulmonary bypass. *Chest* 1997;112(1):193-201.
24. Vallverdu I, Calaf N, Subirana M, Net A, Benito S, Mancebo J. Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1998;158(6):1855-62.
25. Jones DP, Byrne P, Morgan C, Fraser I, Hyland R. Positive end-expiratory pressure vs T-piece. Extubation after mechanical ventilation. *Chest* 1991;100(6):1655-9.
26. Petrof BJ, Legare M, Goldberg P, Milic-Emili J, Gottfried SB. Continuous positive airway pressure reduces work of breathing and dyspnea during weaning from mechanical ventilation in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1990;141(2):281-9.

27. MacIntyre NR, Cheng KC, McConnell R. Applied PEEP during pressure support reduces the inspiratory threshold load of intrinsic PEEP. *Chest* 1997;111(1):188-93.
28. Sydow M, Golisch W, Buscher H, Zinserling J, Crozier TA, Burchardi H. Effect of low-level PEEP on inspiratory work of breathing in intubated patients, both with healthy lungs and with COPD. *Intensive Care Med* 1995;21(11):887-95.
29. Ranieri VM, Giuliani R, Cinnella G, Pesce C, Brienza N, Ippolito EL, et al. Physiologic effects of positive end-expiratory pressure in patients with chronic obstructive pulmonary disease during acute ventilatory failure and controlled mechanical ventilation. *Am Rev Respir Dis* 1993;147(1):5-13.
30. Cohen CA, Zigelbaum G, Gross D, Roussos C, Macklem PT. Clinical manifestations of inspiratory muscle fatigue. *Am J Med* 1982;73(3):308-16.
31. Brochard L, Harf A, Lorino H, Lemaire F. Inspiratory pressure support prevents diaphragmatic fatigue during weaning from mechanical ventilation. *Am Rev Respir Dis* 1989;139(2):513-21.
32. Laghi F, D'Alfonso N, Tobin MJ. Pattern of recovery from diaphragmatic fatigue over 24 hours. *J Appl Physiol* (1985) 1995;79(2):539-46.
33. Slutsky AS. Mechanical ventilation. American College of Chest Physicians' Consensus Conference. *Chest* 1993;104(6):1833-59.
34. Marini JJ, Rodriguez RM, Lamb V. The inspiratory workload of patient-initiated mechanical ventilation. *Am Rev Respir Dis* 1986;134(5):902-9.
35. Bonmarchand G, Chevron V, Chopin C, Jusserand D, Girault C, Moritz F, et al. Increased initial flow rate reduces inspiratory work of breathing during pressure support ventilation in patients with exacerbation of chronic obstructive pulmonary disease. *Intensive Care Med* 1996;22(11):1147-54.
36. Sassoon CS, Del Rosario N, Fei R, Rheeman CH, Gruer SE, Mahutte CK. Influence of pressure- and flow-triggered synchronous intermittent mandatory ventilation on inspiratory muscle work. *Crit Care Med* 1994;22(12):1933-41.
37. Sassoon CS, Lodia R, Rheeman CH, Kuei JH, Light RW, Mahutte CK. Inspiratory muscle work of breathing during flow-by, demand-flow, and continuous-flow systems in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1992;145(5):1219-22.
38. Tutuncu AS, Cakar N, Camci E, Esen F, Telci L, Akpir K. Comparison of pressure- and flow-triggered pressure-support ventilation on weaning parameters in patients recovering from acute respiratory failure. *Crit Care Med* 1997;25(5):756-60.
39. Le Bourdelles G, Mier L, Fiquet B, Djedaini K, Saumon G, Coste F, et al. Comparison of the effects of heat and moisture exchangers and heated humidifiers on ventilation and gas exchange during weaning trials from mechanical ventilation. *Chest* 1996;110(5):1294-8.

40. Iotti GA, Olivei MC, Palo A, Galbusera C, Veronesi R, Comelli A, et al. Unfavorable mechanical effects of heat and moisture exchangers in ventilated patients. *Intensive Care Med* 1997;23(4):399-405.
41. Nava S, Ambrosino N, Clini E, Prato M, Orlando G, Vitacca M, et al. Noninvasive mechanical ventilation in the weaning of patients with respiratory failure due to chronic obstructive pulmonary disease. A randomized, controlled trial. *Ann Intern Med* 1998;128(9):721-8.
42. Girault C, Daudenthun I, Chevron V, Tamion F, Leroy J, Bonmarchand G. Noninvasive ventilation as a systematic extubation and weaning technique in acute-on-chronic respiratory failure: a prospective, randomized controlled study. *Am J Respir Crit Care Med* 1999;160(1):86-92.
43. Jiang JS, Kao SJ, Wang SN. Effect of early application of biphasic positive airway pressure on the outcome of extubation in ventilator weaning. *Respirology* 1999;4(2):161-5.
44. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3(11):e442.
45. Lopez AD, Shibuya K, Rao C, Mathers CD, Hansell AL, Held LS, et al. Chronic obstructive pulmonary disease: current burden and future projections. *Eur Respir J* 2006;27(2):397-412.
46. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. Executive summary 2011. Available from:<http://www.goldcopd.org/>. [Accessed On: 6 Aug, 2014].
47. Tirimanna PR, van Schayck CP, den Otter JJ, van Weel C, van Herwaarden CL, van den Boom G, et al. Prevalence of asthma and COPD in general practice in 1992: has it changed since 1977? *Br J Gen Pract* 1996;46(406):277-81.
48. Halbert RJ, Natoli JL, Gano A, Badamgarav E, Buist AS, Mannino DM. Global burden of COPD: systematic review and meta-analysis. *Eur Respir J* 2006;28(3):523-32.
49. Halbert RJ, Isonaka S, George D, Iqbal A. Interpreting COPD prevalence estimates: what is the true burden of disease? *Chest* 2003;123(5):1684-92.
50. Andersson F, Borg S, Jansson SA, Jonsson AC, Ericsson A, Prutz C, et al. The costs of exacerbations in chronic obstructive pulmonary disease (COPD). *Respir Med* 2002;96(9):700-8.
51. National Heart, Lung, and Blood Institute. Morbidity and mortality chartbook on cardiovascular, lung and blood diseases. Bethesda, Maryland: US Department of Health and Human Services, Public Health Service, National Institute of Health; 2004.

52. Chapman KR, Mannino DM, Soriano JB, Vermeire PA, Buist AS, Thun MJ, et al. Epidemiology and costs of chronic obstructive pulmonary disease. *Eur Respir J* 2006;27(1):188-207.
53. Jansson SA, Andersson F, Borg S, Ericsson A, Jonsson E, Lundback B. Costs of COPD in Sweden according to disease severity. *Chest* 2002;122(6):1994-2002.
54. Hurd SS. International efforts directed at attacking the problem of COPD. *Chest* 2000;117(5 Suppl 2):336s-8s.
55. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet* 1997;349(9064):1498-504.
56. Celli BR, Halbert RJ, Nordyke RJ, Schau B. Airway obstruction in never smokers: results from the Third National Health and Nutrition Examination Survey. *Am J Med* 2005;118(12):1364-72.
57. Behrendt CE. Mild and moderate-to-severe COPD in nonsmokers: distinct demographic profiles. *Chest* 2005;128(3):1239-44.
58. Hogg JC. Pathophysiology of airflow limitation in chronic obstructive pulmonary disease. *Lancet* 2004;364(9435):709-21.
59. Saetta M, Turato G, Maestrelli P, Mapp CE, Fabbri LM. Cellular and structural bases of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;163(6):1304-9.
60. Cosio MG, Majo J, Cosio MG. Inflammation of the airways and lung parenchyma in COPD: role of T cells. *Chest* 2002;121(5 Suppl):160s-5s.
61. Wright JL, Levy RD, Churg A. Pulmonary hypertension in chronic obstructive pulmonary disease: current theories of pathogenesis and their implications for treatment. *Thorax* 2005;60(7):605-9.
62. Birring SS, Brightling CE, Bradding P, Entwisle JJ, Vara DD, Grigg J, et al. Clinical, radiologic, and induced sputum features of chronic obstructive pulmonary disease in nonsmokers: a descriptive study. *Am J Respir Crit Care Med* 2002;166(8):1078-83.
63. Barnes PJ, Shapiro SD, Pauwels RA. Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J* 2003;22(4):672-88.
64. Stockley RA. Neutrophils and the pathogenesis of COPD. *Chest* 2002;121(5 Suppl):151s-5s.
65. Barnes PJ. Alveolar macrophages as orchestrators of COPD. *Copd* 2004;1(1):59-70.
66. Barnes PJ. Mediators of chronic obstructive pulmonary disease. *Pharmacol Rev* 2004;56(4):515-48.

67. Rahman I. Oxidative stress in pathogenesis of chronic obstructive pulmonary disease: cellular and molecular mechanisms. *Cell Biochem Biophys* 2005;43(1):167-88.
68. MacNee W. Oxidative stress and lung inflammation in airways disease. *Eur J Pharmacol* 2001;429(1-3):195-207.
69. Ito K, Ito M, Elliott WM, Cosio B, Caramori G, Kon OM, et al. Decreased histone deacetylase activity in chronic obstructive pulmonary disease. *N Engl J Med* 2005;352(19):1967-76.
70. Hogg JC, Chu F, Utokaparch S, Woods R, Elliott WM, Buzatu L, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350(26):2645-53.
71. O'Donnell DE, Revill SM, Webb KA. Dynamic hyperinflation and exercise intolerance in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;164(5):770-7.
72. Burgel PR, Nadel JA. Roles of epidermal growth factor receptor activation in epithelial cell repair and mucin production in airway epithelium. *Thorax* 2004;59(11):992-6.
73. Barbera JA, Peinado VI, Santos S. Pulmonary hypertension in chronic obstructive pulmonary disease. *Eur Respir J* 2003;21(5):892-905.
74. Wouters EF, Creutzberg EC, Schols AM. Systemic effects in COPD. *Chest* 2002;121(5 Suppl):127s-30s.
75. Agusti AG, Noguera A, Sauleda J, Sala E, Pons J, Busquets X. Systemic effects of chronic obstructive pulmonary disease. *Eur Respir J* 2003;21(2):347-60.
76. Similowski T, Agusti A, MacNee W, Schonhofer B. The potential impact of anaemia of chronic disease in COPD. *Eur Respir J* 2006;27(2):390-6.
77. Gan WQ, Man SF, Senthilselvan A, Sin DD. Association between chronic obstructive pulmonary disease and systemic inflammation: a systematic review and a meta-analysis. *Thorax* 2004;59(7):574-80.
78. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. American Thoracic Society. *Am J Respir Crit Care Med* 1995;152(5 Pt 2):S77-121.
79. Siafakas NM, Vermeire P, Pride NB, Paoletti P, Gibson J, Howard P, et al. Optimal assessment and management of chronic obstructive pulmonary disease (COPD). The European Respiratory Society Task Force. *Eur Respir J* 1995;8(8):1398-420.
80. BTS guidelines for the management of chronic obstructive pulmonary disease. The COPD Guidelines Group of the Standards of Care Committee of the BTS. *Thorax* 1997;52 Suppl 5:S1-28.

81. American Thoracic Society. Chronic bronchitis, asthma, and pulmonary emphysema. *Am Rev Respir Dis* 1962;85:762-8.
82. Rennard SI. COPD: overview of definitions, epidemiology, and factors influencing its development. *Chest* 1998;113(4 Suppl):235s-41s.
83. Pierce JA, Hocott JB, Ebert RV. The collagen and elastin content of the lung in emphysema. *Ann Intern Med* 1961;55:210-22.
84. Vlahovic G, Russell ML, Mercer RR, Crapo JD. Cellular and connective tissue changes in alveolar septal walls in emphysema. *Am J Respir Crit Care Med* 1999;160(6):2086-92.
85. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: Executive summary 2006. GoldCOPD 2007. Available from: <http://www.goldcopd.org>. [Accessed On: 29 Mar, 2014].
86. Jeffery PK. Comparison of the structural and inflammatory features of COPD and asthma. Giles F. Filley Lecture. *Chest* 2000;117(5 Suppl 1):251s-60s.
87. Aoshiba K, Nagai A. Differences in airway remodeling between asthma and chronic obstructive pulmonary disease. *Clin Rev Allergy Immunol* 2004;27(1):35-43.
88. Baraldo S, Turato G, Badin C, Bazzan E, Beghe B, Zuin R, et al. Neutrophilic infiltration within the airway smooth muscle in patients with COPD. *Thorax* 2004;59(4):308-12.
89. Sutherland ER, Martin RJ. Airway inflammation in chronic obstructive pulmonary disease: comparisons with asthma. *J Allergy Clin Immunol* 2003;112(5):819-28.
90. Turato G, Zuin R, Miniati M, Baraldo S, Rea F, Beghe B, et al. Airway inflammation in severe chronic obstructive pulmonary disease: relationship with lung function and radiologic emphysema. *Am J Respir Crit Care Med* 2002;166(1):105-10.
91. Celli BR, Cote CG, Marin JM, Casanova C, Montes de Oca M, Mendez RA, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350(10):1005-12.
92. Rodriguez-Roisin R. Toward a consensus definition for COPD exacerbations. *Chest* 2000;117(5 Suppl 2):398s-401s.
93. Celli BR. Change in the BODE index reflects disease modification in COPD: lessons from lung volume reduction surgery. *Chest* 2006;129(4):835-6.
94. Puhan MA, Mador MJ, Held U, Goldstein R, Guyatt GH, Schunemann HJ. Interpretation of treatment changes in 6-minute walk distance in patients with COPD. *Eur Respir J* 2008;32(3):637-43.

95. Cote CG, Pinto-Plata VM, Marin JM, Nekach H, Dordelly LJ, Celli BR. The modified BODE index: validation with mortality in COPD. *Eur Respir J* 2008;32(5):1269-74.
96. Burge S, Wedzicha JA. COPD exacerbations: definitions and classifications. *Eur Respir J Suppl* 2003;41:46s-53s.
97. Anthonisen NR, Manfreda J, Warren CP, Hershfield ES, Harding GK, Nelson NA. Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;106(2):196-204.
98. Connors AF, Jr., Dawson NV, Thomas C, Harrell FE, Jr., Desbiens N, Fulkerson WJ, et al. Outcomes following acute exacerbation of severe chronic obstructive lung disease. The SUPPORT investigators (Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments). *Am J Respir Crit Care Med* 1996;154(4 Pt 1):959-67.
99. National Institute for Clinical Excellence (NICE). Chronic obstructive pulmonary disease. National clinical guideline on management of chronic obstructive pulmonary disease in adults in primary and secondary care. *Thorax* 2004;59(Suppl 1):1232.
100. Celli BR. Standards for the optimal management of COPD: a summary. *Chest* 1998;113(4 Suppl):283s-7s.
101. Bach PB, Brown C, Gelfand SE, McCrory DC. Management of acute exacerbations of chronic obstructive pulmonary disease: a summary and appraisal of published evidence. *Ann Intern Med* 2001;134(7):600-20.
102. Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med* 2001;163(5):1256-76.
103. Choi PP, Day A, Etchells E. Gaps in the care of patients admitted to hospital with an exacerbation of chronic obstructive pulmonary disease. *Cmaj* 2004;170(9):1409-13.
104. Saudny-Unterberger H, Martin JG, Gray-Donald K. Impact of nutritional support on functional status during an acute exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997;156(3 Pt 1):794-9.
105. Celli BR, MacNee W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 2004;23(6):932-46.
106. Stoller JK. Clinical practice. Acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 2002;346(13):988-94.
107. Tarpy SP, Celli BR. Long-term oxygen therapy. *N Engl J Med* 1995;333(11):710-4.

108. Cydulka RK, Emerman CL. Effects of combined treatment with glycopyrrolate and albuterol in acute exacerbation of chronic obstructive pulmonary disease. *Ann Emerg Med* 1995;25(4):470-3.
109. Nair S, Thomas E, Pearson SB, Henry MT. A randomized controlled trial to assess the optimal dose and effect of nebulized albuterol in acute exacerbations of COPD. *Chest* 2005;128(1):48-54.
110. O'Driscoll BR, Taylor RJ, Horsley MG, Chambers DK, Bernstein A. Nebulised salbutamol with and without ipratropium bromide in acute airflow obstruction. *Lancet* 1989;1(8652):1418-20.
111. In chronic obstructive pulmonary disease, a combination of ipratropium and albuterol is more effective than either agent alone, an 85-day multicenter trial: COMBIVENT Inhalation Aerosol Study Group. *Chest* 1994;105(5):1411-9.
112. Karpel JP. Bronchodilator responses to anticholinergic and beta-adrenergic agents in acute and stable COPD. *Chest* 1991;99(4):871-6.
113. Albert RK, Martin TR, Lewis SW. Controlled clinical trial of methylprednisolone in patients with chronic bronchitis and acute respiratory insufficiency. *Ann Intern Med* 1980;92(6):753-8.
114. Niewoehner DE, Erbland ML, Deupree RH, Collins D, Gross NJ, Light RW, et al. Effect of systemic glucocorticoids on exacerbations of chronic obstructive pulmonary disease. Department of Veterans Affairs Cooperative Study Group. *N Engl J Med* 1999;340(25):1941-7.
115. Quon BS, Gan WQ, Sin DD. Contemporary management of acute exacerbations of COPD: a systematic review and metaanalysis. *Chest* 2008;133(3):756-66.
116. Davies L, Angus RM, Calverley PM. Oral corticosteroids in patients admitted to hospital with exacerbations of chronic obstructive pulmonary disease: a prospective randomised controlled trial. *Lancet* 1999;354(9177):456-60.
117. de Jong YP, Uil SM, Grotjohan HP, Postma DS, Kerstjens HA, van den Berg JW. Oral or IV prednisolone in the treatment of COPD exacerbations: a randomized, controlled, double-blind study. *Chest* 2007;132(6):1741-7.
118. Daniels JM, Snijders D, de Graaff CS, Vlaspolder F, Jansen HM, Boersma WG. Antibiotics in addition to systemic corticosteroids for acute exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2010;181(2):150-7.
119. Stockley RA, O'Brien C, Pye A, Hill SL. Relationship of sputum color to nature and outpatient management of acute exacerbations of COPD. *Chest* 2000;117(6):1638-45.
120. Nouria S, Marghli S, Belghith M, Besbes L, Elatrous S, Abroug F. Once daily oral ofloxacin in chronic obstructive pulmonary disease exacerbation requiring mechanical ventilation: a randomised placebo-controlled trial. *Lancet* 2001;358(9298):2020-5.

121. Glynn TJ, Manley MW. How to help your patients stop smoking. A National Cancer Institute manual for physicians. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute;1990.
122. Woodhead M, Blasi F, Ewig S, Huchon G, Ieven M, Ortqvist A, et al. Guidelines for the management of adult lower respiratory tract infections. *Eur Respir J* 2005;26(6):1138-80.
123. Seemungal T, Harper-Owen R, Bhowmik A, Moric I, Sanderson G, Message S, et al. Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;164(9):1618-23.
124. Monso E, Ruiz J, Rosell A, Manterola J, Fiz J, Morera J, et al. Bacterial infection in chronic obstructive pulmonary disease. A study of stable and exacerbated outpatients using the protected specimen brush. *Am J Respir Crit Care Med* 1995;152(4 Pt 1):1316-20.
125. Pela R, Marchesani F, Agostinelli C, Staccioli D, Cecarini L, Bassotti C, et al. Airways microbial flora in COPD patients in stable clinical conditions and during exacerbations: a bronchoscopic investigation. *Monaldi Arch Chest Dis* 1998;53(3):262-7.
126. Blasi F, Damato S, Cosentini R, Tarsia P, Raccanelli R, Centanni S, et al. Chlamydia pneumoniae and chronic bronchitis: association with severity and bacterial clearance following treatment. *Thorax* 2002;57(8):672-6.
127. Seemungal TA, Wedzicha JA, MacCallum PK, Johnston SL, Lambert PA. Chlamydia pneumoniae and COPD exacerbation. *Thorax* 2002;57(12):1087-8.
128. Soler N, Torres A, Ewig S, Gonzalez J, Celis R, El-Ebiary M, et al. Bronchial microbial patterns in severe exacerbations of chronic obstructive pulmonary disease (COPD) requiring mechanical ventilation. *Am J Respir Crit Care Med* 1998;157(5 Pt 1):1498-505.
129. Fagon JY, Chastre J, Trouillet JL, Domart Y, Dombret MC, Bornet M, et al. Characterization of distal bronchial microflora during acute exacerbation of chronic bronchitis. Use of the protected specimen brush technique in 54 mechanically ventilated patients. *Am Rev Respir Dis* 1990;142(5):1004-8.
130. Miravittles M, Espinosa C, Fernandez-Laso E, Martos JA, Maldonado JA, Gallego M. Relationship between bacterial flora in sputum and functional impairment in patients with acute exacerbations of COPD. Study Group of Bacterial Infection in COPD. *Chest* 1999;116(1):40-6.
131. Eller J, Ede A, Schaberg T, Niederman MS, Mauch H, Lode H. Infective exacerbations of chronic bronchitis: relation between bacteriologic etiology and lung function. *Chest* 1998;113(6):1542-8.

132. Kolodziej MA, Jensen L, Rowe B, Sin D. Systematic review of noninvasive positive pressure ventilation in severe stable COPD. *Eur Respir J* 2007;30(2):293-306.
133. Sethi S, Evans N, Grant BJ, Murphy TF. New strains of bacteria and exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 2002;347(7):465-71.
134. Fogarty C, de Wet R, Mandell L, Chang J, Rangaraju M, Nusrat R. Five-day telithromycin once daily is as effective as 10-day clarithromycin twice daily for the treatment of acute exacerbations of chronic bronchitis and is associated with reduced health-care resource utilization. *Chest* 2005;128(4):1980-8.
135. Wilson R, Allegra L, Huchon G, Izquierdo JL, Jones P, Schaberg T, et al. Short-term and long-term outcomes of moxifloxacin compared to standard antibiotic treatment in acute exacerbations of chronic bronchitis. *Chest* 2004;125(3):953-64.
136. Wilson R, Schentag JJ, Ball P, Mandell L. A comparison of gemifloxacin and clarithromycin in acute exacerbations of chronic bronchitis and long-term clinical outcomes. *Clin Ther* 2002;24(4):639-52.
137. Snow V, Lascher S, Mottur-Pilson C. Evidence base for management of acute exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 2001;134(7):595-9.
138. Black PN, Morgan-Day A, McMillan TE, Poole PJ, Young RP. Randomised, controlled trial of N-acetylcysteine for treatment of acute exacerbations of chronic obstructive pulmonary disease [ISRCTN21676344]. *BMC Pulm Med* 2004;4:13.
139. Shannon M. Predictors of major toxicity after theophylline overdose. *Ann Intern Med* 1993;119(12):1161-7.
140. Bahls FH, Ma KK, Bird TD. Theophylline-associated seizures with "therapeutic" or low toxic serum concentrations: risk factors for serious outcome in adults. *Neurology* 1991;41(8):1309-12.
141. Mulloy E, McNicholas WT. Theophylline improves gas exchange during rest, exercise, and sleep in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1993;148(4 Pt 1):1030-6.
142. Brander PE, Salmi T. Nocturnal oxygen saturation and sleep quality in patients with advanced chronic obstructive pulmonary disease during treatment with moderate dose CR-theophylline. *Eur J Clin Pharmacol* 1992;43(2):125-9.
143. Martin RJ, Pak J. Overnight theophylline concentrations and effects on sleep and lung function in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1992;145(3):540-4.
144. Berry RB, Desa MM, Branum JP, Light RW. Effect of theophylline on sleep and sleep-disordered breathing in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1991;143(2):245-50.

145. Hirano T, Yamagata T, Gohda M, Yamagata Y, Ichikawa T, Yanagisawa S, et al. Inhibition of reactive nitrogen species production in COPD airways: comparison of inhaled corticosteroid and oral theophylline. *Thorax* 2006;61(9):761-6.
146. Barnes PJ. Theophylline for COPD. *Thorax* 2006;61(9):742-4.
147. Greenstone M, Lasserson TJ. Doxapram for ventilatory failure due to exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2003(1):Cd000223.
148. Clinical indications for noninvasive positive pressure ventilation in chronic respiratory failure due to restrictive lung disease, COPD, and nocturnal hypoventilation--a consensus conference report. *Chest* 1999;116(2):521-34.
149. Lightowler JV, Wedzicha JA, Elliott MW, Ram FS. Non-invasive positive pressure ventilation to treat respiratory failure resulting from exacerbations of chronic obstructive pulmonary disease: Cochrane systematic review and meta-analysis. *Bmj* 2003;326(7382):185.
150. Meyer TJ, Hill NS. Noninvasive positive pressure ventilation to treat respiratory failure. *Ann Intern Med* 1994;120(9):760-70.
151. Brochard L, Mancebo J, Wysocki M, Lofaso F, Conti G, Rauss A, et al. Noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 1995;333(13):817-22.
152. Kramer N, Meyer TJ, Meharg J, Cece RD, Hill NS. Randomized, prospective trial of noninvasive positive pressure ventilation in acute respiratory failure. *Am J Respir Crit Care Med* 1995;151(6):1799-806.
153. Bott J, Carroll MP, Conway JH, Keilty SE, Ward EM, Brown AM, et al. Randomised controlled trial of nasal ventilation in acute ventilatory failure due to chronic obstructive airways disease. *Lancet* 1993;341(8860):1555-7.
154. Plant PK, Owen JL, Elliott MW. Early use of non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease on general respiratory wards: a multicentre randomised controlled trial. *Lancet* 2000;355(9219):1931-5.
155. International Consensus Conferences in Intensive Care Medicine: noninvasive positive pressure ventilation in acute Respiratory failure. *Am J Respir Crit Care Med* 2001;163(1):283-91.
156. Rossi A, Gottfried SB, Zocchi L, Higgs BD, Lennox S, Calverley PM, et al. Measurement of static compliance of the total respiratory system in patients with acute respiratory failure during mechanical ventilation. The effect of intrinsic positive end-expiratory pressure. *Am Rev Respir Dis* 1985;131(5):672-7.
157. Parker CM, Voduc N, Aaron SD, Webb KA, O'Donnell DE. Physiological changes during symptom recovery from moderate exacerbations of COPD. *Eur Respir J* 2005;26(3):420-8.

158. Conti G, Antonelli M, Navalesi P, Rocco M, Bui M, Spadetta G, et al. Noninvasive vs. conventional mechanical ventilation in patients with chronic obstructive pulmonary disease after failure of medical treatment in the ward: a randomized trial. *Intensive Care Med* 2002;28(12):1701-7.
159. Esteban A, Anzueto A, Frutos F, Alia I, Brochard L, Stewart TE, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *Jama* 2002;287(3):345-55.
160. Wildman MJ, Sanderson C, Groves J, Reeves BC, Ayres J, Harrison D, et al. Implications of prognostic pessimism in patients with chronic obstructive pulmonary disease (COPD) or asthma admitted to intensive care in the UK within the COPD and asthma outcome study (CAOS): multicentre observational cohort study. *Bmj* 2007;335(7630):1132.
161. Gunen H, Hacievliyagil SS, Kosar F, Mutlu LC, Gulbas G, Pehlivan E, et al. Factors affecting survival of hospitalised patients with COPD. *Eur Respir J* 2005;26(2):234-41.
162. Purro A, Appendini L, De Gaetano A, Gudjonsdottir M, Donner CF, Rossi A. Physiologic determinants of ventilator dependence in long-term mechanically ventilated patients. *Am J Respir Crit Care Med* 2000;161(4 Pt 1):1115-23.
163. Torres A, Reyes A, Roca J, Wagner PD, Rodriguez-Roisin R. Ventilation-perfusion mismatching in chronic obstructive pulmonary disease during ventilator weaning. *Am Rev Respir Dis* 1989;140(5):1246-50.
164. Beydon L, Cinotti L, Rekik N, Radermacher P, Adnot S, Meignan M, et al. Changes in the distribution of ventilation and perfusion associated with separation from mechanical ventilation in patients with obstructive pulmonary disease. *Anesthesiology* 1991;75(5):730-8.
165. Hilbert G, Gruson D, Portel L, Gbikpi-Benissan G, Cardinaud JP. Noninvasive pressure support ventilation in COPD patients with postextubation hypercapnic respiratory insufficiency. *Eur Respir J* 1998;11(6):1349-53.
166. Celli BR. Pulmonary rehabilitation in patients with COPD. *Am J Respir Crit Care Med* 1995;152(3):861-4.
167. Heffner JE, Fahy B, Hilling L, Barbieri C. Outcomes of advance directive education of pulmonary rehabilitation patients. *Am J Respir Crit Care Med* 1997;155(3):1055-9.
168. Esteban A, Frutos-Vivar F, Ferguson ND, Arabi Y, Apezteguia C, Gonzalez M, et al. Noninvasive positive-pressure ventilation for respiratory failure after extubation. *N Engl J Med* 2004;350(24):2452-60.
169. Marini JJ, Smith TC, Lamb VJ. External work output and force generation during synchronized intermittent mechanical ventilation. Effect of machine assistance on breathing effort. *Am Rev Respir Dis* 1988;138(5):1169-79.

170. Ward ME, Corbeil C, Gibbons W, Newman S, Macklem PT. Optimization of respiratory muscle relaxation during mechanical ventilation. *Anesthesiology* 1988;69(1):29-35.
171. Jubran A, Tobin MJ. Reliability of pulse oximetry in titrating supplemental oxygen therapy in ventilator-dependent patients. *Chest* 1990;97(6):1420-5.
172. Bickler PE, Feiner JR, Severinghaus JW. Effects of skin pigmentation on pulse oximeter accuracy at low saturation. *Anesthesiology* 2005;102(4):715-9.
173. Gibney RT, Wilson RS, Pontoppidan H. Comparison of work of breathing on high gas flow and demand valve continuous positive airway pressure systems. *Chest* 1982;82(6):692-5.
174. Giuliani R, Mascia L, Recchia F, Caracciolo A, Fiore T, Ranieri VM. Patient-ventilator interaction during synchronized intermittent mandatory ventilation. Effects of flow triggering. *Am J Respir Crit Care Med* 1995;151(1):1-9.
175. Aslanian P, El Atrous S, Isabey D, Valente E, Corsi D, Harf A, et al. Effects of flow triggering on breathing effort during partial ventilatory support. *Am J Respir Crit Care Med* 1998;157(1):135-43.
176. Tobin MJ, Jubran A, Laghi F. Patient-ventilator interaction. *Am J Respir Crit Care Med* 2001;163(5):1059-63.
177. Pepe PE, Marini JJ. Occult positive end-expiratory pressure in mechanically ventilated patients with airflow obstruction: the auto-PEEP effect. *Am Rev Respir Dis* 1982;126(1):166-70.
178. Tobin MJ, Jubran A, Hines E, Jr. Pathophysiology of failure to wean from mechanical ventilation. *Schweiz Med Wochenschr* 1994;124(47):2139-45.
179. Leung P, Jubran A, Tobin MJ. Comparison of assisted ventilator modes on triggering, patient effort, and dyspnea. *Am J Respir Crit Care Med* 1997;155(6):1940-8.
180. Coussa ML, Guerin C, Eissa NT, Corbeil C, Chasse M, Braidy J, et al. Partitioning of work of breathing in mechanically ventilated COPD patients. *J Appl Physiol* (1985) 1993;75(4):1711-9.
181. Smith TC, Marini JJ. Impact of PEEP on lung mechanics and work of breathing in severe airflow obstruction. *J Appl Physiol* (1985) 1988;65(4):1488-99.
182. Tobin MJ, Lodato RF. PEEP, auto-PEEP, and waterfalls. *Chest* 1989;96(3):449-51.
183. Tobin MJ, Alex CA, Fahey PJ. Fighting the ventilator. In: Tobin MJ (ed). *Principles and practice of mechanical ventilation*. New York: McGraw Hill; 2006. 1121
184. Connors AF, Jr., McCaffree DR, Gray BA. Effect of inspiratory flow rate on gas exchange during mechanical ventilation. *Am Rev Respir Dis* 1981;124(5):537-43.

185. Laghi F, Segal J, Choe WK, Tobin MJ. Effect of imposed inflation time on respiratory frequency and hyperinflation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;163(6):1365-70.
186. Puddy A, Younes M. Effect of inspiratory flow rate on respiratory output in normal subjects. *Am Rev Respir Dis* 1992;146(3):787-9.
187. Jubran A, Van de Graaff WB, Tobin MJ. Variability of patient-ventilator interaction with pressure support ventilation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;152(1):129-36.
188. Dhand R, Duarte AG, Jubran A, Jenne JW, Fink JB, Fahey PJ, et al. Dose-response to bronchodilator delivered by metered-dose inhaler in ventilator-supported patients. *Am J Respir Crit Care Med* 1996;154(2 Pt 1):388-93.
189. Dhand R, Tobin MJ. Inhaled bronchodilator therapy in mechanically ventilated patients. *Am J Respir Crit Care Med* 1997;156(1):3-10.
190. Laubscher TP, Heinrichs W, Weiler N, Hartmann G, Brunner JX. An adaptive lung ventilation controller. *IEEE Trans Biomed Eng* 1994;41(1):51-9.
191. Mishra M, Chaudhri S, Tripathi V, Verma AK, Sampath A, Chauhan NK. Weaning of mechanically ventilated chronic obstructive pulmonary disease patients by using non-invasive positive pressure ventilation: A prospective study. *Lung India* 2014;31(2):127-33.
192. Stawicki SP. Mechanical ventilation weaning and extubation. *OPUS* 2007;1(2):13-5.
193. Boutou AK, Abatzidou F, Tryfon S, Nakou C, Pitsiou G, Argyropoulou P, et al. Diagnostic accuracy of the rapid shallow breathing index to predict a successful spontaneous breathing trial outcome in mechanically ventilated patients with chronic obstructive pulmonary disease. *Heart Lung* 2011;40(2):105-10.
194. Matic I, Danic D, Majeric-Kogler V, Jurjevic M, Mirkovic I, Mrzljak Vucinic N. Chronic obstructive pulmonary disease and weaning of difficult-to-wean patients from mechanical ventilation: randomized prospective study. *Croat Med J* 2007;48(1):51-8.
195. Conti G, Antonelli M, Arzano S, Gasparetto A. Measurement of occlusion pressures in critically ill patients. *Crit Care* 1997;1(3):89-93.
196. Moon R, Camporesi E. Respiratory monitoring. In: Miller R (ed). *Miller's anesthesia*. 6th ed. New York: Elsevier; 2005. 1466

197. Lamia B, Cuvelier A, Benichou J, Muir JF. [A multi-centre randomized controlled trial of domiciliary non-invasive ventilation vs long-term oxygen therapy in survivors of acute hypercapnic respiratory failure due to COPD. Non-invasive ventilation in obstructive lung disease (NIVOLD) study]. *Rev Mal Respir* 2012;29(9):1141-8.
198. Gluck EH. Predicting eventual success or failure to wean in patients receiving long-term mechanical ventilation. *Chest* 1996;110(4):1018-24.
199. Logan CM, Rice MK. Logan's medical and scientific abbreviations. Philadelphia: J. B. Lippincott Company; 1987.
200. World Health Organization (WHO). Global database on body mass index. Geneva: WHO; 2006.
201. Pruitt B. Weaning patients from mechanical: ventilation. *Nursing* 2006;36(9):36-42.
202. Zheng DW, Wang CZ, Liu RS, Gao F, Deng SL, Zhou P, et al. The application of improved Glasgow coma scale score of 15 as switching point for invasive noninvasive mechanical ventilation in treatment of severe respiratory failure in chronic obstructive pulmonary disease. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 2011;23(4):224-7.
203. Karanjia N, Nordquist D, Stevens R, Nyquist P. A clinical description of extubation failure in patients with primary brain injury. *Neurocrit Care* 2011;15(1):4-12.
204. Farah R, Khamisy-Farah R, Arraf Z, Jacobson L, Makhoul N. Hypophosphatemia as a prognostic value in acute exacerbation of COPD. *Clin Respir J* 2013;7(4):407-15.
205. Barrientos-Vega R, Mar Sanchez-Soria M, Morales-Garcia C, Robas-Gomez A, Cuena-Boy R, Ayensa-Rincon A. Prolonged sedation of critically ill patients with midazolam or propofol: impact on weaning and costs. *Crit Care Med* 1997;25(1):33-40.
206. Zanotti E, Rubini F, Iotti G, Braschi A, Palo A, Bruschi C, et al. Elevated static compliance of the total respiratory system: early predictor of weaning unsuccess in severed COPD patients mechanically ventilated. *Intensive Care Med* 1995;21(5):399-405.
207. Nozawa E, Kobayashi E, Matsumoto ME, Feltrim MI, Carmona MJ, Auler JJ. Assessment of factors that influence weaning from long-term mechanical ventilation after cardiac surgery. *Arq Bras Cardiol* 2003;80(3):301-10.
208. Terzano C, Di Stefano F, Conti V, Di Nicola M, Paone G, Petroianni A, et al. Mixed acid-base disorders, hydroelectrolyte imbalance and lactate production in hypercapnic respiratory failure: the role of noninvasive ventilation. *PLoS One* 2012;7(4):e35245.

209. Li-dong S, Chang-sheng G, Zi-yu Z. Explore the influence of BMI in the optimal time of weaning from sequential mechanical ventilation for severity chronic obstructive pulmonary disease. *BMC Emerg Med* 2013;13(Suppl 1):S1.
210. Frazier SK, Brom H, Widener J, Pender L, Stone KS, Moser DK. Prevalence of myocardial ischemia during mechanical ventilation and weaning and its effects on weaning success. *Heart Lung* 2006;35(6):363-73.
211. Lara TM, Hajjar LA, de Almeida JP, Fukushima JT, Barbas CS, Rodrigues AR, et al. High levels of B-type natriuretic peptide predict weaning failure from mechanical ventilation in adult patients after cardiac surgery. *Clinics (Sao Paulo)* 2013;68(1):33-8.
212. Mekontso Dessap A, Roche-Campo F, Kouatchet A, Tomicic V, Beduneau G, Sonnevile R, et al. Natriuretic peptide-driven fluid management during ventilator weaning: a randomized controlled trial. *Am J Respir Crit Care Med* 2012;186(12):1256-63.
213. Lai YC, Ruan SY, Huang CT, Kuo PH, Yu CJ. Hemoglobin levels and weaning outcome of mechanical ventilation in difficult-to-wean patients: a retrospective cohort study. *PLoS One* 2013;8(8):e73743.
214. Marcelino P, Germano N, Nunes AP, Flora L, Moleiro A, Marum S, et al. Cardiac influence on mechanical ventilation time and mortality in exacerbated chronic respiratory failure patients. The role of echocardiographic parameters. *Rev Port Pneumol* 2006;12(2):131-46.
215. Routsis C, Stanopoulos I, Zakyntinos E, Politis P, Papas V, Zervakis D, et al. Nitroglycerin can facilitate weaning of difficult-to-wean chronic obstructive pulmonary disease patients: a prospective interventional non-randomized study. *Crit Care* 2010;14(6):R204.
216. Huang CT, Tsai YJ, Lin JW, Ruan SY, Wu HD, Yu CJ. Application of heart-rate variability in patients undergoing weaning from mechanical ventilation. *Crit Care* 2014;18(1):R21.
217. Madani S, Saghafinia M, Nezhad H, Ebadi A, Ghojani A, Tavasoli A, et al. Validity of integrative weaning index of discontinuation from mechanical ventilation in Iranian ICUs. *Thrita* 2013;2(4):62-8.
218. Tobin MJ, Perez W, Guenther SM, Semmes BJ, Mador MJ, Allen SJ, et al. The pattern of breathing during successful and unsuccessful trials of weaning from mechanical ventilation. *Am Rev Respir Dis* 1986;134(6):1111-8.
219. Lima EJ. Respiratory rate as a predictor of weaning failure from mechanical ventilation. *Braz J Anesthesiol* 2013;63(1):1-6.
220. Tzanis G, Vasileiadis I, Zervakis D, Karatzanos E, Dimopoulos S, Pitsolis T, et al. Maximum inspiratory pressure, a surrogate parameter for the assessment of ICU-acquired weakness. *BMC Anesthesiol* 2011;11:14.


221. Vargas F, Boyer A, Bui HN, Salmi LR, Guenard H, Gruson D, et al. Respiratory failure in chronic obstructive pulmonary disease after extubation: value of expiratory flow limitation and airway occlusion pressure after 0.1 second (P0.1). *J Crit Care* 2008;23(4):577-84.
222. El Khoury MY, Panos RJ, Ying J, Almoosa KF. Value of the PaO₂:FiO₂ ratio and Rapid Shallow Breathing Index in predicting successful extubation in hypoxemic respiratory failure. *Heart Lung* 2010;39(6):529-36.
223. Rialp G, Raurich JM, Llompart-Pou JA, Ayestaran I, Ibanez J. Respiratory CO₂ response depends on plasma bicarbonate concentration in mechanically ventilated patients. *Med Intensiva* 2014;38(4):203-10.
224. Burns KE, Meade MO, Premji A, Adhikari NK. Noninvasive positive-pressure ventilation as a weaning strategy for intubated adults with respiratory failure. *Cochrane Database Syst Rev* 2013;12:Cd004127.

PROTOCOL

سامية من

**COMPARISON BETWEEN PREDICTORS OF
WEANING FROM MECHANICAL VENTILATOR IN COPD
PATIENTS AND THE EFFECT OF USAGE OF CPAP IN
DIFFICULT WEANING PATIENTS**

مقارنة بين قياسات التنبيب بالقطام من جهاز التهوية الميكانيكية في مرضى
الأنسداد الرئوي المزمن وتأثير استخدام الضغط الهوائى الموجب المستمر فى مرضى
القطام الصعب



Protocol of thesis submitted,
To the faculty of medicine,
Alexandria university,
In partial fulfillment of the
requirements of the degree of
master of critical care.

خطة بحث مقدمة
لكلية الطب
جامعة الاسكندرية
ايفاء جزئيا
لشروط الحصول على درجة الماجستير
في الطب الحرج.

by

من

Khaled Bahaa Eldin Mohammed Desouky,
MBBCh:Alex,
Resident doctor,
Department of critical care,
Faculty of medicine,
University of Alexandria,
2013

خالد بهاء الدين محمد دسوقي
بكالوريوس الطب والجراحة - الاسكندرية
طبيب مقيم
قسم الطب الحرج
كلية الطب
جامعة الاسكندرية
٢٠١٣



SUPERVISORS

Prof. Dr. Mahmoud Ibrahim Mahmoud

Professor of chest diseases,
Department of chest diseases,
Faculty of Medicine,
University of Alexandria.

د. محمود ابراهيم محمود

استاذ الامراض الصدرية
قسم الامراض الصدرية
كلية الطب
جامعة الاسكندرية.

Dr. Mohammed Mustafa Megahed

Assistant professor of Critical Care,
Department of Critical Care,
Faculty of Medicine,
University of Alexandria.

د. محمد مصطفى مجاهد

استاذ مساعد الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الاسكندرية.




CO- RESEARCHER

Mustafa Gamal Belal,
Sixth grade,
Faculty of Medicine,
University of Alexandria,
Telephone number:01093288871
Email: Moonhi_2001@yahoo.com.

3
الباحث المساعد

مصطفى جمال بلال
الفرقة السادسة
كلية الطب
جامعة الاسكندرية.



Page 4

INTRODUCTION

Mechanical ventilation(MV) is the defining event of intensive care unit management. Although it is a life saving intervention in many diseases, a major goal of critical care clinicians should be to liberate patients from MV as early as possible to avoid the multitude of complications and risks associated with prolonged unnecessary MV including ventilator induced lung injury, ventilator associated pneumonia, increased length of ICU and hospital stay and cost of care delivery. ^(1, 2)

Weaning decisions based only on expert clinical judgment are not always correct. Premature discontinuation places severe stress on the respiratory and cardiovascular systems, while unnecessary delays can lead to diaphragmatic atrophy so several predictors of weaning are therefore used to aid decision-making. ⁽³⁾

According to the American college of chest physicians, society of critical care medicine and the American association of respiratory care the weaning indices that have some predictive capacity are: ⁽⁴⁾

1) Tidal volume: Spontaneous tidal volumes greater than 5 ml/kg have been considered as good predictors of weaning outcome. ⁽⁵⁾

2) Maximal inspiratory pressure($P_{I_{max}}$): it is commonly used to test respiratory muscle strength. The proximal end of the endotracheal tube is occluded for 20 to 25 seconds with a one-way valve that allows the patient to exhale but not to inhale. This procedure leads to increasing inspiratory effort and $P_{I_{max}}$ measured towards the end of the occlusion period. Several early studies have shown that $P_{I_{max}} \leq -30$ cmH₂O has a high predictive value for weaning. ⁽⁶⁾

Handwritten initials: "NB" and "NB"

3) Breathing frequency (f): Tachypnea ($f \geq 30-35$ breaths/minute) is a sensitive marker of respiratory distress but can prolong intubation when used as an exclusive criterion. ⁽⁷⁾

4) $P_{0.1}/P_{I_{max}}$: The airway occlusion pressure ($P_{0.1}$) is the pressure measured at the airway opening 0.1 second after inspiring against an occluded airway. The $P_{0.1}$ is effort independent and correlates well with central respiratory drive. When combined with the maximal inspiratory pressure $P_{I_{max}}$, the $P_{0.1}/P_{I_{max}}$ ratio at a value of <0.3 has been found to be a good early predictor of weaning success. ⁽⁸⁾

5) Oxygenation: Arterial oxygen saturation (SaO_2) $\geq 90\%$ with a fraction of inspired oxygen $FiO_2 \leq 0.4$ associated with successful weaning. ⁽⁹⁾

6) Minute ventilation (V_E): $V_E < 10$ l/minute is associated with weaning success. Studies showed that V_E values $> 15-20$ l/minute are helpful in identifying patients who are unlikely to be liberated from mechanical ventilation. ⁽¹⁰⁾

7) Rapid shallow breathing index: It is the ratio of breathing frequency to tidal volume obtained during the first 1 minute of a T-piece trial. At a threshold value of ≤ 105 breaths/minute/liter, RSBI was a significantly better predictor of weaning outcomes than other 'classic' and commonly used parameters. ^{(11),(12)}

L *MSB*

→ Cw⁶

AIM OF THE STUDY

To compare between different weaning parameters in combination of the general condition of the COPD patients as predictors of weaning and to study the effect of usage of CPAP in difficult weaning patients.

Dr. N. S.

oboikeyanadl.com

→ Calc 7

PATIENTS

The study will be conducted on 100 COPD patients admitted to Alexandria main university hospital who are mechanically ventilated for more than 24 hours and will be considered candidates for weaning judged by the physician in charge.

Inclusion criteria: Mechanically ventilated for at least 24 hours and the cause of mechanical ventilation has been resolved judged by the physician in charge.

Exclusion criteria: Patients younger than 18 years, patients on vasopressors, patients with toxicological diseases.

Informed consent will be taken from every patient included in the study whenever possible or from patient's next of kin. All selected patients fulfilling the inclusion criteria will be subjected to full history (including age, sex and preexisting underlying disease), full clinical examination and Laboratory investigations will be done on admission .



METHODS

The following values will be measured in all COPD patients candidate for weaning:

1.frequency.

2.tidal volume.

3.rapid shallow breathing index.

4.maximum inspiratory pressure: will be measured on volume control ventilation while the patient fully sedated and during inspiratory pause. A value less than 25cmH₂O is considered one of predictors of weaning success.

5.P0.1: will be measured on volume control ventilation while the patient is fully sedated and during first 0.1 second of patient inspiration.

6.compliance: will be measured while patient on volume control ventilation, fully sedated and inspiratory pause for 0.5 to 1 S, $Cst = VT / (\text{inspiratory plateau pressure} - \text{positive end expiratory pressure})$.^(14,15)

7.PaO₂: will be measured using arterial blood gas analysis.

8.PaCO₂: will be measured using arterial blood gas analysis.

9.PEEP.

10.hypoxic index: will be measured by dividing PaO₂ by FiO₂.



9

11. haemoglobin level: should be more than 9 g/dl.

12. integrative weaning index (IWI): will be calculated by this equation: $IWI = Cst \times SaO_2 / RSBI$. (Cst: static compliance, SaO₂: saturation of O₂ in arterial blood, RSBI: rapid shallow breathing index). A value of more than 25 ml/cmH₂O breaths/minute/liter is a good predictor of weaning outcome.

These parameters will be considered with neurological status (structural brain damage or electrolyte disturbances), respiratory status (myopathy, neuropathy, change in compliance or resistance or V/Q mismatch), metabolic status (nutrition and acid base status) and cardiovascular status (presence of ischemic heart disease arrhythmias or heart failure). In addition vital signs of the patient will be measured (BP, temperature, respiratory rate and heart rate).

Then patient will be discontinued from MV by the physician in charge through a T-piece trial for 2 hours and if spontaneous breathing trial succeeded then success of weaning will be determined by the ability to breath normally apart from MV for 48 hours⁽¹⁶⁾. The decision to return to MV will be made by the physician in charge (who will be completely blind to the study).⁽¹⁷⁾

nb

According to the success of the weaning trial patients are classified into 2 groups:

GROUP 1: failure of weaning group.

GROUP 2: success of weaning group.

The study will also include the effect of usage of CPAP in difficult weaning patients.

Different weaning parameters (frequency, TV, RSBI, Pmax, minute ventilation, P0.1, compliance, Pmax, minute ventilation, P0.1, compliance, PaO₂, PaCO₂, PEEP, hypoxic index) in association with the general condition of the patient (neurologically, respiratory, cardiovascular and metabolic) and vital signs will be collected.

Patients will be divided into 2 groups: group 1: who succeeded in the trial of weaning (sustained ability to breath apart from mechanical ventilator for 48 hours after successful spontaneous breathing trial for 2 hours using T-piece) and group 2: who failed in the trial of weaning and all weaning parameters, general condition and vital signs of the patient will be compared to determine specificity, sensitivity, positive and negative predictive values and accuracy of each of them.

The study will also include the effect of usage of CPAP in difficult weaning patients.

WB

NT

11

RESULTS

The results will be reported, interpreted and analysed to fulfill the aim of the research.

 MBS

obeyikandani.com

→ Col¹²

DISCUSSION

The results of this study will be interpreted and compared with other studies.

to
not

obeyikandani.com

REFERENCES

1. Hall JB, Wood LDH. Liberation of the patient from mechanical ventilation. JAMA 1987, 257:1621-28.
2. Afessa B, Hogans L, Murphy R. Predicting 3-day and 7-day outcomes of weaning from mechanical ventilation. Chest 1999, 116:456-61.
3. Tobin MJ. Advances in mechanical ventilation. N Engl J Med 2001, 344:1986-96.
4. MacIntyre NR, Cook DJ, Ely Jr, Epstein SK, Fink JB, Heffner JE, Hess D, Hubmayer RD, Scheinborn DJ: Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College for Critical Care Medicine. Chest 2001 , 120:375-95.
5. Holliday JE, Hyers TM. The reduction of weaning time from mechanical ventilation using tidal volume and relaxation feedback. Am Rev Respir Dis 1990, 141:1214-20.
6. Trwit JD, Marini JJ. Validation of a technique to assess maximal inspiratory pressure in poorly cooperative patients. Chest 1992, 102:1216-19.
7. DeHaven CB, Kirton OC, Hart AM, Shatz DV, Civetta JM. Breathing measurement reduces false-negative classification of tachypneic preextubation trial failures. Crit Care Med 1996, 24:976-80.

nb

nl

8. Kuhlen R, Hausmann S, Pappert D, Slama K, Rossaint R, Falke K. A new method for P0.1 measurement using standard respiratory equipment. *Intensive Care Med* 1995, 21:554-60.
9. Girard TD, Ely WE. Protocol-driven ventilator weaning: Reviewing the evidence. *Clin Chest Med* 2008, 29:241-52.
10. Martinez A, Seymour C, Nam M. Minute ventilation recovery time: a predictor of extubation outcome. *Chest* 2003, 123:1214-21.
11. Lee KH, Hui KP, Chan TB, Tan WC, Lim TK: Rapid shallow breathing (frequency tidal volume ratio) did not predict extubation outcome. *Chest* 1994 ; 105:540-3.
12. Epstein SK: Evaluation of the rapid shallow breathing index in the clinical setting. *Am J Respir Crit Care Med* 1995 ; 152:545-9.
13. Nemer S, Carmen SV, Jefferson B, Thiago C , Ricardo G, Luiz C et al A new integrative weaning index of discontinuation from mechanical ventilation. *Critical care* 2009; 13:1176 - 93.
14. Nucci G, Mergoni M, Polese G, Cobelli C, Rossi A. On-line monitoring of respiratory mechanics. In: Brusasco V, Macklem PT, Pedotti A, eds. *Mechanics of Breathing*. Milan, Springer-Verlag, 2002: 327-36
15. Tobin MJ, Van de Graaff WB. Monitoring of lung mechanics and work of breathing. In: Tobin MJ, ed. *Principles and Practice of Mechanical Ventilation*. McGraw-Hill, New York, 1994, 967-1003.

16. Manthous CA, Schmidt GA, Hall JB. Liberation from mechanical ventilation: a decade of progress. *Chest* 1998, 114:886-901.
17. Epstein SK. Weaning parameters. *Respir Care Clin N Am* 2000; 6:253-301.
18. Vallverdu I, Calaf N, Subirana M. Clinical characteristics respiratory functional parameters, and outcomes of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1998; 158:1855-62
19. Brochard L, Rauss A, Benito S. Comparison of three methods of gradual withdrawal from ventilatory support during weaning from MV. *Am J Respir Crit Care Med*. 1994; 150(4):896-903.
20. Esteban A, Frutos F, Tobin MJ. A comparison of four methods of weaning patients from mechanical ventilation. Spanish Lung Failure Collaborative Group. *N Engl J Med*. 1995; 332(6):345-50.
21. Esteban A, Alia I, Ibanez J. Modes of mechanical ventilation and weaning: a national survey of Spanish Hospitals; the Spanish Lung failure collaborative group. *Chest* 1994; 106:1188-93.
22. Esteban A, Alia I, Gordo F. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. *Am J Respir Crit Care Med* 1997, 156:459-65.
23. Farias JA, Retta A, Alia I. A comparison of two methods to perform a breathing trial before extubation in paediatric intensive care patients. *Intensive Care Med* 2001; 27:1649-54.

nl
nls

16

24. Yang KL, Tobin MJ. A Prospective Study of Indexes Predicting the Outcome of Trials of Weaning from Mechanical Ventilation. N Engl J Med 1991; 324:1445-50.

25. Krieger BP, Isber J, Breitenbucher A, Throop G, Ershowsky P. Serial Measurements of the Rapid-Shallow-Breathing Index as a Predictor of Weaning Outcome in Elderly Medical Patients. American college of chest physician (CHEST) 1997;112:1029-34.

Handwritten signature

obitkarnadi.com

ARABIC SUMMARY

الملخص العربي

الغطام من التهوية الميكانيكية يمكن تعريفها بأنها عملية سحب دعم التنفس الصناعي فجأة أو تدريجياً. مرض الانسداد الرئوي المزمن هو مرض يمكن الوقاية منه وعلاجه مع بعض الآثار الهامة خارج الرئة التي يمكن أن تسهم في شدة المرض في عدد من المرضى. يتميز عنصرها الرئوي بالحد من تدفق الهواء الذي لا يمكن عكسه بشكل كامل. الحد من تدفق الهواء عادة ما يكون تقدماً ومصحوباً باستجابة التهابية غير طبيعية من الرئة إلى الجزيئات السامة أو الغازات .

في دراستنا قارنا بين مؤشرات الغطام المختلفة بالإضافة الي الحالة العامة لمرضى السدة الرئوية المزمنة (العصبية والايونية والجهاز التنفسي والقلب والأوعية الدموية والوضع الحمضي القاعدي ومؤشر كتلة الجسم) كمتنبآت للغطام ودرسنا تأثير استخدام التهوية بدعم الضغط غير الغازي في مرضى الغطام الصعب .

وقد أجريت الدراسة على ١٠٠ مريض من مرضى السدة الرئوية المزمنة تم قبولهم إلى المستشفى الجامعي الرئيسي بالإسكندرية الذين استوفوا معايير الاشتمال والاقصاء، و تم وضعهم علي جهاز التهوية الميكانيكية لأكثر من ٢٤ ساعة وتم اعتبارهم مرشحين للغطام من قبل الطبيب المختص. تعرض كل المرضى المختارين والذين حققوا معايير الاشتمال الي اخذ التاريخ المرضي كاملاً(بما في ذلك السن والجنس والأمراض الكامنة سابقة الوجود)، والفحص السريري الكامل والفحوصات المختبرية أجريت عند الحجز ويومياً .

تم قياس القيم التالية في جميع مرضى الانسداد الرئوي المزمن المرشحين للغطام: العلامات الحيوية(ضغط الدم الانقباضي والانبساطي، الحرارة، معدل التنفس، معدل ضربات القلب)، الحجم المدي(TV) ، مؤشر التنفس السريع الضحل(RSBI)، الضغط الشهيق الاقصى(MIP)، الضغط الشهيق في اول ١٠٠ملي ثانية(P0.1)، والامتثال، الضغط الشرياني للاكسجين(PaO2)، الضغط الشرياني لثاني اكسيد الكربون(PaCO2)، النهاية الايجابية للضغط الزفيري(PEEP)، مؤشرا نقص الاكسجين(HI)، ومؤشر الغطام التكامل(IWI) والنسبة بين الضغط الشهيق في اول ١٠٠ملي ثانية و الضغط الشهيق الاقصى(P0.1/MIP ratio).

واعترت هذه المؤشرات مع الحالة العصبية والايونية (وجود تلف هيكلي في المخ، ووجود اضطرابات بالايونات الكهربائية، مقياس جلاسكو للغيوبية (GCS)، وجود أخذ المهدنات أو المنومات قبل محاولة الغطام)، وحالة الجهاز التنفسي (وجود قيمة غير طبيعية للامتثال أو المقاومة و التدرج الأوكسجيني الشرياني الحويصلي -PAO2)) (PaO2)، والوضع الحمضي القاعدي ومؤشر كتلة الجسم (مؤشر كتلة الجسم ووجود اضطراب حمضي قاعدي) وحالة القلب والأوعية الدموية (مرض نقص تروية القلب، اضطراب بضربات القلب أو فشل عضلة القلب ومستوى الهيموجلوبين) .

وفقاً لنجاح محاولة غطام المرضى تم تصنيفهم إلى مجموعتين: المجموعة ١: مجموعة فشلت في محاولة الغطام والمجموعة ٢ نجحت في محاولة الغطام.(نجاح الغطام تقرر من القدرة على التنفس بشكل طبيعي بعيداً عن جهاز التهوية الميكانيكية لمدة ٤٨ ساعة). ثم تمت مقارنة كل هذه المؤشرات الفطامية بالإضافة إلى المؤشرات التي تعكس الحالة العصبية والايونية والجهاز التنفسي والوضع الحمضي القاعدي ومؤشر كتلة الجسم وحالة القلب والأوعية الدموية للمريض إلى جانب العلامات الحيوية للمريض وفقاً لخصوصيتها، وحساسيتها، والقيم التنبؤية الإيجابية والسلبية ودقتها للكشف عن قدراتهم كمتنبآت للغطام من جهاز التهوية الميكانيكية. شملت الدراسة أيضاً تأثير استخدام التهوية بدعم الضغط الغير الغازي في مرضى الغطام الصعب.

وجدنا أن:

١. كانت أهم البيانات وأكثرها مغزاة في دراستنا في توقع نتائج الغطام من التهوية الميكانيكية في مرضى الانسداد الرئوي المزمن هي مؤشر الغطام التكامل (IWI)، مؤشر التنفس السريع الضحل (RSBI)، ومقياس جلاسكو للغيوبية (GCS) ومعدل ضربات القلب و وجود قيمة غير طبيعية من المقاومة. كان مقياس جلاسكو للغيوبية (GCS) الأفضل في الخصوصية والقيمة التنبؤية الإيجابية والدقة، في حين أن وجود المقاومة غير طبيعية الأفضل في الحساسية والقيمة التنبؤية السلبية .

- ٢ . كانت هناك بيانات أخرى في دراستنا التي يمكنها التنبؤ بالفطام في مرضى السدة الرئوية المزمنة من جهاز التهوية الميكانيكية على الرغم من أنها كانت إحصائيا غير هامة لأنها ينبغي دراستها على عدد أكبر من المرضى . وهم وجود تلف هيكلي في المخ، ووجود اضطرابات بالأيونات الكهربائية خاصة نقص فوسفات الدم، ووجود أخذ مهدئات قبل الفطام، ووجود اضطراب حمضي قاعدي، ووجود مرض نقص تروية القلب ، ووجود فشل في عضلة القلب، ووجود الذبذبة الاذينية ومؤشر كتلة الجسم.
- ٣ . كان مستوى الهيموغلوبين و ضغط الدم الانقباضي ومعدل التنفس والحجم المدي والامتثال الثابت و الضغط الشهيقى الاقصى و الضغط الشهيقى في اول ١٠٠ملي ثانية مؤشرا فقيرا للفطام في مرضى الانسداد الرئوي المزمن من جهاز التهوية الميكانيكية في دراستنا .
- ٤ . الضغط الشرياني للاكسجين(PaO2)، الضغط الشرياني لثاني اكسيد الكربون (PaCO2)و مؤشر نقص الكسجين (HI) لا يمكنهم التنبؤ بدقة الفطام في مرضى الانسداد الرئوي المزمن من جهاز التهويم الميكانيكية.
- ٥ . التهوية بدعم الضغط أعطى أكثر من ٥٠٪ فرصة للفطام في مرضى السدة الرئوية المزمنة من جهاز التهوية الميكانيكية ويمكن ان ينقص من مدة الفطام من جهاز التهوية الميكانيكية.

الملخص العربي

لجنة الممتحنين

موافقون

أ.د/ محمود ابراهيم محمود

أستاذ الأمراض الصدرية
قسم الأمراض الصدرية
كلية الطب
جامعة الإسكندرية

.....

أ.د/ صلاح عبد الفتاح محمد اسماعيل

أستاذ التخدير والعناية المركزة الجراحية
قسم التخدير والعناية المركزة الجراحية
كلية الطب
جامعة قناة السويس

.....

أ.د/ تيسير محمد زيتون

أستاذ الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الإسكندرية

.....

أ.م.د/ محمد مصطفى مجاهد

أستاذ مساعد الطب الحرج
قسم الطب الحرج
كلية الطب
جامعة الإسكندرية

.....



جامعة الإسكندرية
كلية الطب
قسم الطب الحرج

مقارنة بين قياسات التنبؤ بالفطام من جهاز التهوية الميكانيكية في مرضى
الانسداد الرئوى المزمن وتأثير استخدام الضغط الهوائى الموجب المستمر فى مرضى
الفطام الصعب

رسالة مقدمة

لقسم الطب الحرج - كلية الطب - جامعة الإسكندرية
ضمن متطلبات درجة

الماجستير

فى

الطب الحرج

مقدمة من

خالد بهاء الدين محمد دسوقي الطويلة

بكالوريوس الطب والجراحة: الإسكندرية، ٢٠٠٨

كلية الطب

جامعة الإسكندرية

٢٠١٥