

CHAPTER 6

REFERENCES

- A. Necchi, N. E. Pogna, and S. Mapelli (1987)** “Early and late heat shock proteins in wheats and other cereal species,” *Plant Physiology*, vol. 84, no. 4, pp. 1378–1384, 1987.
- Aase, T.H., R.P. Chaudhary, and O.R. Vetaas, (2010).**Farming flexibility and food security under climatic uncertainty: Manang, Nepal Himalaya. *Area* 42:228–238.
- Abd El-Monem (2007)** Polyamines as modulators of wheat growth, metabolism and reproductive development under high temperature stress. Ph.D. Thesis, Ain Shamas Univ., Cairo, Egypt.
- Abdel-Aal ESM, Sosulski FW, Youssef MM, Shehata AAY, (1993).**Selected nutritional, physical and sensory characteristics of pan and flat breads prepared from composite flours containing fababean. *Plant Foods for Human Nutrition*, 51: 409–414.
- Abd-El-Haleem SHM, Reham MA, Mohamed SMS, Abdel-Aal ESM, Sosulski FW, Hucl P, (1998).**Origins, characteristics and potentials of ancient wheats. *Cereal Foods World*, 43: 708–715.
- Acevedo, E., P. Silva, and H. Silva. (2002).** Wheat growth and physiology. In: Curtis, B. C., S. Rajaram and H. G. Macpherson (Eds.) *Bread Wheat Improvement and Production*, FAO Plant Production and Protection Series, No. 30. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Adam, Z. and Clarke, A.K. (2002)** Cutting edge of chloroplast proteolysis. *Trends Plant Sci.* 7, 451–456
- Adam, Z. et al. (2001)** Chloroplast and mitochondrial proteases in Arabidopsis. A proposed nomenclature. *Plant Physiol.* 125, 1912–1918
- Adams ML, Lombi E, Zhao FJ, McGrath SP, (2002).** Evidence of low selenium concentrations in UK bread-making wheat grain. *Journal of the Science of Food and Agriculture*, 82: 1160–1165.
- Agarwal, M. et al. (2001)** Arabidopsis thaliana Hsp100 proteins: kith and kin. *Cell Stress Chaperones* 6, 219–224
- Agarwal, M. et al. (2003)** Molecular characterization of rice HSP101: complementation of yeast hsp104 mutation by disaggregation of protein granules and differential expression in indica and japonica rice types. *Plant Mol. Biol.* 51, 543–553
- Agarwal, M., Katiyar-Agarwal, S. and Grover, A. (2002)** Plant Hsp100 proteins: structure, function and regulation. *Plant Sci.* 163, 397–405.
- Agarwal, M., Katiyar-Agarwal, S., Sahi, C., Gallie, D.R. and Grover, A. (2001)** Arabidopsis thaliana Hsp100 proteins: kith and kin. *Cell Stress Chaperon.* 6, 219–224.
- Aghaei, K., A.A. Ehsanpour, A.H. Shah & S. Komatsu, (2009).**Proteome analysis of soybean hypocotyl and root under salt stress. *Amino Acids.*, 36: 91-98.
- Ahmed JU and Hasan MA, (2011)** Evaluation of seedling proline content of wheat genotypes in relation to heat tolerance *Bangladesh J. Bot.* 40(1): 17-22, 2011 (June)
- Ahn YJ, Claussen K, Zimmerman JL (2004)** Genotypic differences in the heat shock response and thermotolerance in four potato varieties. *Plant Science* 166, 901-911

- Ahn, Y.-J., Zimmerman and J.L., (2006).** Introduction of the carrot HSP17.7 into potato (*Solanum tuberosum* L.) enhances cellular membrane stability and tuberization in vitro. *Plant Cell Environ.* 29, 95–104.
- Akhunov, E., Nicolet, C. and Dvorak, J. (2009)** Single nucleotide polymorphism genotyping in polyploidy wheat with the Illumina GoldenGate assay. *Theor. Appl. Genet.* 119, 507–517.
- Al katib, K. and G.M. Paulsen, (1990).** Photosynthesis and productivity during high temperature stress of wheat genotypes from major world regions. *Crop Sci.*, 30: 1127 .
- Alan JB, Changrun LMS, Fergus M, Clydesdale FACN, Decker EA, (2000).** Potential of wheat-based breakfast cereals as a source of dietary antioxidants. *Journal of the American College of Nutrition*, 19(3): 308S–311S.
- Al-Khatib, K., and Paulsen, G.M. (1999).** High-temperature effects on photosynthetic analysis and chlorophyll fluorescence imaging. *Plant Physiology and*
- Allakhverdiev, S.I.; Kreslavski, V.D.; Klimov, V.V; Los, D.A. (2008);** Carpentier, R.; Mohanty, P. Heat stress: An overview of molecular responses in photosynthesis. *Photosynth. Res.* 2008, 98, 541–550.
- Allan A.C. & Fluhr R. (2007).** Ozone and Reactive Oxygen Species. *Encyclopedia of Life Sciences*, DOI: 10.1038/npg.els.0001299
- Almeselmani, M., P.S. Deshmukh, R.K. Sairam., S.R .Kushwaha and T.P.Singh, (2006).** Protective role of antioxidant enzymes under high temperature stress. *Plant Sci.*, 171: 382-388.
- Al-Whaibi (2011)** MH plant heat shock proteins: A mini review. *Journal of King Saud University* 23(2), 139-150
- Amani, I., R.A. Fisher, and M.P. Reynolds, (1996).** Canopy temperature depression association with yield of irrigated spring wheat cultivars in hot climate. *J. Agron. Crop Sci* 176:119-129.
- Anonymous, 2010** United States Department of Agriculture, Foreign Agriculture Service. www.fas.usda.gov.
- Anonymous (2002)** World Wheat Facts, Western Organization of Resources Council. <http://www.worc.org/pdfs/WorldWheatFacts.pdf>. Accessed May 2006.
- Anonymous. 2005a** The biology and ecology of bread wheat (*Triticum aestivum* L.) in Australia. Australian Government. [Online] Available: http://www.ogtr.gov.au/pdf/ir/biology_wheat.
- Arbona, V.; López-Climent, M.F.; Pérez-Clemente, R.M. 2009;** Gómez-Cadenas, A. Maintenance of a high photosynthetic performance is linked to flooding tolerance in citrus. *Environ. Exp. Bot.* 2009, 66, 135–142.
- Ashraf, M., Foolad, M.R., (2007)** Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environ. Exp. Bot.* 59, 206–216.
- Ashraf, M. and P.J.C. Harris, (2004)** Potential biochemical indicators of salinity tolerance in plants. *Plant Science*, 166: 3-16.

- Ashraf, M., Saeed, M.M., Qureshi, M.J., (1994).** Tolerance to high temperature in cotton (*Gossypium hirsutum* L.) at initial growth stages. *Environ. Exp. Bot.*
- Baga M., Fowler D. B. and Chibbar R. N. (2009)** Identification of genomic regions determining the phenological development leading to floral transition in wheat (*Triticum aestivum* L.). *J. Exp.Bot.*60, 3575–3585.
- Bailey-Serres, J.; Voesenek, L.A.C.J. (2008)** Flooding stress: Acclimations and genetic diversity. *Ann. Rev. Plant Biol.* 2008, 59, 313–339.
- Baker, N.R. (2008).** Chlorophyll Fluorescence: a probe of photosynthesis in vivo. *Annual Review of Plant Biology*, 59, 89-113, ISSN 1543-5008
- Baker, N.R., & Rosenqvist, E. (2004).** Applications of chlorophyll fluorescence can improve crop production strategies: an examination of future possibilities. *Journal of Biochemistry*, 46, 189-195, ISSN 0981- 9428 *Biophys. Acta* 1517, 270–277.
- Baker, N.R., East, T.M., & Long, S.P. (1983).** Chilling damage to photosynthesis in young *Zea mays*. II. Photochemical function of thylakoids in vivo. *Journal of Experimental Botany*,34, 189-197, ISSN 0022-0957
- Balachandran, S., Osmond, C.B., & Daley, P.F. (1994).** Diagnosis of the earliest strain-specific interactions between tobacco mosaic virus and chloroplasts of tobacco leaves *in vivo* by means of chlorophyll fluorescence imaging. *Plant Physiology*, 104, 1059-1065,
- Bale JS, Masters GJ, Hodkinson ID, et al .** Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology* 2002;8:1-16.
- Baniwal, S.K., Bharti, K., Chan, K.Y., Fauth, M., Ganguli, A., Kotak, S., Mishra, S.K., Nover, L., Port, M., Scharf, K., Tripp, L., Weber, C., Zielinski, D., von Koskull-Doering, P., (2004).** Heat stress response in plants: a complex game with chaperones and more than 20 heat stress transcription factors. *J. Biosci.* 29, 471–487.
- Barker, G.L.A. and Edwards, K.J. (2009)** A genome-wide analysis of single nucleotide polymorphism diversity in the world’s major cereal crops. *Plant Biotechnol. J.* 7, 318–325.
- Barnett NM and AW Naylor (1966)** Amino acid and protein metabolism in Bermuda grass during water stress. *Plant Physiol.* 41: 1222-1230.
- Barnett, T., Altschuler, M., McDaniel, C.N., Mascarenhas, J.P., (1980).**Heat shock induced proteins in plant cells. *Dev. Genet.* 1, 331–340.
- Bates, L.S., Waldran, R.P. and Teare, I.D. (1973)** Rapid determination of free proline for water stress studies. *Plant Soil* 39: 205-208.
- Behl, R.K., H.S. Nainawatee and K.P. Singh. 1993.** High temperature tolerance in wheat, pp. 349-355. In: *International Crop Science I.* Crop Science Society of America. 677 S. Segoe Rd., WI 53711, USA.

- Beillard E, Pallisgaard N, van der Velden VH, Bi W, Dee R, van der Schoot E, Delabesse E, Macintyre E, Gottardi E, Saglio G, Watzinger F, Lion T, van Dongen JJ, Hokland P, Gabert J (2003)** Evaluation of candidate control genes for diagnosis and residual disease detection in leukemic patients using ‘real-time’ quantitative reverse-transcriptase polymerase chain reaction (RQ-PCR) – a Europe against cancer program. *Leukemia* 17:2474–2486
- Bennett, M. D. (1972)** Nuclear DNA content and minimum generation time in herbaceous plants. *Proc. R. Soc. (Lond.) Ser. B.* 181: 109_135.
- Bernardo, A.N., Bradbury, P.J., Ma, H., Hu, S., Bowden, R.L., Buckler, E.S. and Bia, G. (2009)** Discovery and mapping of single feature polymorphisms in wheat using Affymetrix arrays. *BMC Genomics*, 10, 251.
- Berry JA and O Bjorkman (1980)** Photosynthetic response and adaptation to temperature in higher plants. *Ann. Rev. Plant Physiol.* 31: 491-543.
- Bhanu Parkash, (1997)** Influence of moisture and temperature stress in tolerant and susceptible wheat genotypes. Ph.D . Thesis, IARI., New Delhi, India.
- Bharti, K., Nover, L., (2002).** Heat stress-induced signaling. In: Scheel, D., Wasternack, C. (Eds.), *Plant Signal Transduction: Frontiers in Molecular Biology.* Oxford University Press, Oxford, UK., pp. 74–115.
- Bhoja R. Basnet, Mohamed B. Ali, Amir M.H. Ibrahim, Thomas Payne, Moussa G. Mosaad., (2011).** Evaluation of genetic bases and diversity of Egyptian wheat cultivars released during the last 50 years using coefficient of parentage. *Communications in Biometry and Crop Science* Vol. 6, No. 1, 2011, pp. 31–47
- Blum A and A Ebercon (1976)** Genotypic responses in sorghum to drought stress. III. Free proline accumulation and drought resistance. *Crop Sci.* 16: 428-431.
- Blum A., and A. Ebercon (1981)** Cell membrane as a measure of drought and heat tolerance in wheat. *Crop Sci.* 21: 43-47.
- Blum, A., Klueva, N., Nguyen, H.T., (2001).** Wheat cellular thermotolerance is related to yield under heat stress. *Euphytica* 117, 117–123.
- Boggess SF and GR Stewart (1980)** The relationship between water stress induced proline accumulation and inhibition of protein synthesis in tobacco leaves. *Plant Sci. Letter* 17: 245-252.
- Bohnert, H.J., Gong, Q., Li, P., Ma, S., (2006).** Unraveling abiotic stress tolerance mechanisms—getting genomics going. *Curr. Opin. Plant Biol.* 9, 180–188.
- Boisson, M., Mondon, K., Torney, V., Nicot, N., Laine, A.L., Bahrman, N., et al. (2005).** Partial sequences of nitrogen metabolism genes in hexaploid wheat. *Theor. Appl. Genet.* 110: 932–940. doi:10.1007/s00122-004-1913-4. PMID:15714330.
- Boyer, J.S. (1982)** Plant productivity and environment. *Science* 218:443–448.
- Braun, H.J., Atlin, G. and Payne, T. (2010)** Multi location testing as a tool to identify plant response to global climate change. In: Reynolds, C.R.P. (Ed.), *Climate Change and Crop Production.* CABI: London, UK.

- Bray EA, Bailey-Serres J, Weretilnyk E, (2000).** Responses to abiotic stresses. In: Gruitsem W, Buchannan B, Jones R. *Biochemistry and molecular biology of plants*. Rockville, MD: American Society of Plant Physiologists; 2000. p. 1158-1203.
- Bray, N.J., Buckland, P.R., Owen, M.J. and O'Donovan, M.C. (2003)** Cis-acting variation in the expression of a high proportion of genes in human brain. *Hum. Genet.* 113, 149–153.
- Briantais, J.M., Dacosta, J., Goulas, Y., Ducruet, J.M., & Moya, I. (1996).** Heat stress induces in leaves an increase of the minimum level of chlorophyll fluorescence, F₀ – A timeresolved analysis. *Photosynthesis Research*, 48, 189-196, ISSN 0166-8595
- Bridgwater, W. & Beatrice Aldrich. (1966)** *The Columbia-Viking Desk Encyclopedia*. Columbia University. p. 1959.
- Briggle, L. W. and B. C. Curtis. (1987)** Wheat worldwide. In E. G. Heyne (ed.). *Wheat and wheat improvement*, 2nd ed. *Agronomy* 13:1-32.
- Britikov EA, SV Vladimirtseva and NA Musatova (1965)** Transformation of proline in germinating pollen. *F. Rastenii (English translation)* 12: 953-967.
- Brookes A J. (1999)** The essence of SNPs , *Gene*, 234 (1999) 177- 186.
- Burton BM, Baker TA. (2005).** Remodeling protein complexes: insights from the AAA+ unfoldase ClpX and Mu trans posase. *Prot. Si.* 14: 1945-1954
- Bustin, S. A. 2000** Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays, *J. Mol. Endocrinol.*, 25, 169, 2000.
- Calatayud, A., Gorbe, E., Roca, D., & Martinez, P.F. (2008).** Effect of two nutrient solution temperatures on nitrate uptake, nitrate reductase activity, NH₄⁺ concentration and chlorophyll *a* fluorescence in rose plants. *Environmental and Experimental Botany*, 64,
- Camejo, D., Jiméñez, A., Alarcón, J.J., Torres, W., Gómez, J.M., Sevilla, F., (2006).** Changes in photosynthetic parameters and antioxidant activities following heat-shock treatment in tomato plants. *Funct. Plant Biol.* 33, 177–187.
- Camejo, D., Rodríguez, P., Morales, M.A., Dell'amico, J.M., Torrecillas, A., Alarcón, J.J., (2005).** High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *J. Plant Physiol.* 162, 281–289.
- Camejo, D., Rodríguez, P., Morales, M.A., Dell'amico, J.M., Torrecillas, A., Alarcón, J.J., (2005).** High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *J. Plant Physiol.* 162, 281–289.
- Campbell, J.L., Klueva, N.Y., Zheng, H.G., Nieto-Sotelo, J., Ho, T.H.D. and Nguyen, H.T. (2001)** Cloning of new members of heat shock protein HSP101 gene family in wheat (*Triticum aestivum* L. Moench) inducible by heat, dehydration, and ABA. *Biochim.*

- Carvalho-Neibel F. (2005)** MtENOD11 gene activation during rhizobial infection and mycorrhizal arbuscule development requires a common AT-rich-containing regulatory sequence. 2005 Dec;18(12):1269-76
- Cattivelli, L.; Rizza, F.; Badeck, F.-W.; Mazzucotelli, E.; Mastrangelo, A.M.; Francia, E.; Marè, C.; Tondelli, A.; Stanca, A.M. (2008)** Drought tolerance improvement in crop plants: An integrated view from breeding to genomics. *Field Crops Res.* 2008, *105*, 1–14.
- Cauvain, Stanley P. & Cauvain P. Cauvain. (2003)** *Bread Making*. CRC Press. p. 540. ISBN 1-85573-553-9.
- Chaerle, L., & Van Der Straeten, D. (2000).** Imaging techniques and the early detection of plant stress. *Trends in Plant Science*, *5*, 495–501, ISSN 1360-1385
- Chaerle, L., & Van Der Straeten, D. (2001).** Seeing is believing: imaging techniques to monitor plant health. *Biochimica et Biophysica Acta*, *1519*, 153–166, ISSN 0304-4165
- Chaitanya KV, D Sundar and AR Reddy (2001).** Mulberry leaf metabolism under high temperature stress. *Biol. Plant* *44*(3): 379-384.
- Chaki, M., R. Valderrama, A.M. FernÁNdez-OcaÑA, A. Carreras, M.V. GÓmez-RodrÍGuez, J. LÓpez- Jaramillo, J.C. Begara-Morales, B. SÁNchez-Calvo, F. Luque, M. Leterrier, F.J. Corpas & J.B. Barroso, (2011).** High temperature triggers the metabolism of S-nitrosothiols in sunflower mediating a process of nitrosative stress which provokes the inhibition of ferredoxin–NADP reductase by tyrosine nitration. *Plant, Cell & Environment.*, *34*: 1803-1818
- Chao, S., Dubcovsky, J., Dvorak, J., Luo, M.C., Baenziger, S.P., Matnyazov, R., Clark, D.R., Talbert, L.E., Anderson, J.A., Dreisigacker, S., Glover, K., Chen, J., Campbell, K., Bruckner, P.L., Rudd, J.C., Haley, S., Carver, B.F., Perry, S., Sorrells, M.E. and Akhunov, E.D. (2010)** Population- and genome-specific patterns of linkage disequilibrium and SNP variation in spring and winter wheat (*Triticum aestivum* L.). *BMC Genomics*, *11*, 727.
- Chen B, Retzlaff M, Roos T, Frydman J (2011)** Cellular strategies of protein quality control. *Cold Spring Harbor Perspect Biol* 2011, *3*:14.
- Chen, T.H.H., Shen, Z.Y., Lee, P.H., (1982).** Adaptability of crop plants to high temperature stress. *Crop Sci.* *22*, 719–725.
- Christiansen, M.N. (1978)** The physiology of plant tolerance to temperature extremes. p. 173-191. In G.A. Jung (ed.) *Crop tolerance to suboptimal land conditions*. Am. Soc. Agron. Madison. WI.
- CHU TM, D ASPINALL, LG PALEG (1974)** Stress metabolism. VI. Temperature stress and the accumulation of proline in barley leaves. *Aust J Plant Physiol* *1*:87-97
- Chu TM, D Aspinall and LG Paleb (1978)** Accumulation of free proline at low temperatures. *Physiol. Plants.**43*: 256-260.

- CIMMYT, (2005) CIMMYT Business Plan (2006–2010).** Translating the Vision of Seeds of Innovation into a Vibrant Work Plan. Centro Internacional de Mejoramiento de Maíz y Trigo, El Batán, Mexico. 31–37.
- Cochram J., Jones H., Leigh F. J., Sullivan D. O., Powell W., Laurie D. A. and Greenland A. J. (2007)** Control of flowering time in temperate cereals: genes, domestication and sustainable productivity. *J. Exp. Bot.* 58, 1231–1244.
- Cowles, C.R., Hirschhorn, J.N., Altshuler, D. and Lander, E.S. (2002)** Detection of regulatory variation in mouse genes. *Nat. Genet.* 32, 432–437.
- Curtis, Rajaraman and MacPherson (2002).** "Bread Wheat" (<http://www.fao.org/docrep/006/y4011e/y4011e00.htm>). Food and Agriculture Organization of the United Nations.
- D. G. Lee, N. Ahsan, S. H. Lee et al. (2007).** A proteomic approach in analyzing heat-responsive proteins in rice leaves," *Proteomics*, vol. 7, no. 18, pp. 3369–3383, 2007.
- D. Higgins and W. Taylor (2000)** Bioinformatics: Sequence, structure, and databanks. Oxford University Press Inc., New York © Oxford University Press, 2000 ISBN 19 963791 1
- Dias, A. S., and Lidon, F. C. (2009)** Evaluation of Grain Filling Rate and Duration in Bread and Durum Wheat, under Heat Stress after Anthesis DOI: 10.1111/j.1439-037X.2008.00347.
- Díaz-Almeyda E, Thomé PE, El Hafidi M, Iglesias-Prieto R. (2011)** Differential stability of photosynthetic membranes and fatty acid composition at elevated temperature in *Symbiodinium*. *Coral Reefs* 30: 217–225
- Djanaguiraman, M., P.V.V. Prasad, and M. Seppanen. (2010)** Selenium protects sorghum leaves from oxidative damage under high temperature stress by enhancing antioxidant defense system. *Plant Physiol. Biochem.* 48:999–1007.
- Djoukeng, J.D.; Arbona, V.; Argamasilla, R.; Gomez-cadenas, A. (2008)** Flavonoid profiling in leaves of citrus genotypes under different environmental situations flavonoid profiling in leaves of *Citrus* genotypes. *J. Agric. Food Chem.* 2008, 56, 11087–11097.
- Eagles H. A., Cane K., Kuchel H., Hollanby G. J., Vallance V., Eastwood R. F. et al. (2010)** Photoperiod and vernalization gene effects in southern Australian wheat. *Crop Pasture Sci.* 61, 721–730.
- Easterling DR and Apps M (2005)** Assessing the consequences of climate change for food and forest resources: a view from the IPCC. *Clim Change* 70: 165–189
- Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO (2000).** Climate extremes: observations, modeling, and impacts. *Science* 2000;289:2068–2074.
- Edward K J & Mogg R, (2001)** Plant genotyping by analysis of single nucleotide polymorphisms, in *Plant genotyping: The DNA fingerprinting of plants*, edited by R J Henry (CABI Publishing, UK) 2001, 1-13.

- Edwards, K.J., Reid, A.L., Coghill, J.A., Berry, S.T. and Barker, G.L.A. (2009)** Multiplex single nucleotide polymorphism (SNP)-based genotyping in allohexaploid wheat using padlock probes. *Plant Biotechnol. J.* 7, 375–390.
- Efeoglu B, Terzioglu S (2009)** Photosynthetic responses of two wheat varieties to high temperature. *EurAsian Journal BioScience* 3, 97-106
- Efeoglu, B. and S. Terzioglu. (2007)** Varying patterns of protein synthesis in bread wheat during heat shock. *Acta Biologica Hungarica* 58:93-104.
- Ehlers, J.D., Hall, A.E., (1998)** Heat tolerance of contrasting cowpea lines in short and long days. *Field Crops Res.* 55, 11–21.
- Ehlert, B., & Hinch, D.K. (2008).** Chlorophyll fluorescence imaging accurately quantifies freezing damage and cold acclimation responses in *Arabidopsis* leaves. *Plant Methods*, 4, 12, ISSN 1746-4811
- Evan Ibrahim Al-Jebory (2013).** Changes in cellular membrane tolerance due to heat stress during *Triticum sativum* L. seeds germination, magazin o f alkufa university for biolog / VOL.5 / NO.2
- Evenson, R.E. (2003)** Assessing the impact of the green revolution, 1960 to 2000. *Science* 300:758–762.
- Faris, J.D., B. Friebe, and B.S. Gill. (2002)** Wheat genomics: Exploring the polyploid model. *Curr. Genomics* 3:577–591.
- Feder, M.E., Hoffman, G.E., (1999)** Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu. Rev. Physiol.* 61, 243–282.
- Feldman, M. (1976)** Taxonomic classification and names of wild, primitive, cultivated, and modern cultivated wheats. *Wheats*. Pages 120_128 in N. W. Simmonds, ed. *Evolution of crop plants*. Longman, London, UK.
- Feltus, F.A., Wan, J., Schulze, S.R., Estill, J.C., Jiang, N., and Paterson, A.H. (2004).** An SNP resource for rice genetics and breeding based on subspecies *Indica* and *Japonica* genome alignments. *Genome Res.* 14: 1812–1819. doi:10.1101/gr.2479404. PMID:15342564.
- Ferguson, D.L., Guikema, J.A., Paulsen, G.M., (1990).** Ubiquitin pool modulation and protein degradation in wheat roots during high temperature stress. *Plant Physiol.* 92, 740–746.
- Flexas, J. (2002)** Drought-inhibition of photosynthesis in C3 plants: Stomatal and non-stomatal limitations revisited. *Ann. Bot.* 2002, 89, 183–189.
- Flood R. G. and Halloran G. M. (1984)** The nature and duration of gene action for vernalization response in wheat. *Ann. Bot.* 53, 363–368
- Fokar, M., H.T. Nguyen & A. Blum, 1998.** Heat tolerance in spring wheat. I. Estimating cellular thermotolerance and its heritability. *Euphytica* (In Review).

- Frank, G.; Pressman, E.; Ophir, R.; Althan, L.; Shaked, R.; Freedman, M.; Shen, S.; Firon, N. (2009).** Transcriptional profiling of maturing tomato (*Solanum lycopersicum* L.) microspores reveals the involvement of heat shock proteins, ROS scavengers, hormones, and sugars in the heat stress response. *J. Exp. Bot.* 2009, *60*, 3891–3908.
- Georgieva, K. (1999)** Some mechanisms of damage and acclimation of the photosynthetic apparatus due to high temperature. *Bul. J. Plant Physiol.* 25:89–99.
- Gilmore, A.M., & Govindjee (1999).** How higher plants respond to excess light: Energy dissipation in photosystem II, In: *Concepts in Photobiology: Photosynthesis and Photomorphogenesis*, Singhal G.S., Renger G., Irrgang K.D., Govindjee, Sopory S., pp. 513-548, Narosa Publishing House, New Delhi, ISBN 0-7923-5519-9
- Giorno, F.; Wolters-Arts, M.; Grillo, S.; Scharf, K.-D.; Vriezen, W.-H.; Mariani, C. (2010).** Developmental and heat stress-regulated expression of HsfA2 and small heat shock proteins in tomato anthers. *J. Exp. Bot.* 2010, *61*, 453–462.
- Global Agricultural information net work (Gain) (2012)** Egypt Grain and Feed Annual Wheat and Corn Production on the Rise.
- Godzic S and HF Linskens (1974)** Concentration of free amino acids in primary bean leaves after continuous and interrupted SO₂ fumigation and recovery. *Environmental Pollution* 7: 25-38.
- Goncharov N. P. (2004)** Response to vernalization in wheat: its quantitative or qualitative nature. *Cereal Res. Commun.* 32, 323–330.
- Goswami, A.; Banerjee, R.; Raha, S. (2010)** Mechanisms of plant adaptation/memory in rice seedlings under arsenic and heat stress: Expression of heat-shock protein gene HSP70. *AoB Plants* 2010, 2010, doi:10.1093/aobpla/plq023.
- Grassini, P., K.G. Cassman, (2012).** High-yield maize with large net energy yield and small global warming intensity. *Proceedings of the National Academy of Sciences*, 109: 1074-1079.
- Grigorova, B.; Vaseva, I.I.; Demirevska, K.; Feller, U. (2011)** Expression of selected heat shock proteins after individually applied and combined drought and heat stress. *Acta Physiol. Plant.* 2011, *33*, 2041–2049.
- Guidi, L., & Degl’Innocenti, E. (2008).** Ozone effects on high light-induced photoinhibition in *Phaseolus vulgaris*. *Plant Science* 174: 590-596, ISSN 0306-4484.
- Guidi, L., Mori, S., Degl’innocenti, E., & Pecchia, S. (2007).** Effects of ozone exposure or fungal pathogen on white lupin leaves as determined by imaging of chlorophyll a fluorescence. *Plant Physiology and Biochemistry*, 45, 851-857, ISSN 0981-9428
- Guillaumie S, Charmet G, Linossier L, Torney V, Robert N, Ravel C (2004)** Colocation between a gene encoding the bZip factor SPA and an eQTL for a high-molecular-weight glutenin subunit in wheat (*Triticum aestivum*) Genome. 47(4), 705-713 .
- Guo, M., Rupe, M.A., Danilevskaya, O.N., Yang, X. and Hu, Z. (2003)** Genome-wide mRNA profiling reveals heterochronic allelic variation and a new imprinted gene in hybrid maize endosperm. *Plant J.* 36, 30–44.

- Guo, Y.-P., Zhou, H.-F., Zhang, L.-C., (2006).** Photosynthetic characteristics and protective mechanisms against photooxidation during high temperature stress in two citrus species. *Sci. Hort.* 108, 260–267.
- Gupta, S.C., Sharma, A., Mishra, M., Mishra, R., Chowdhuri, D.K., (2010).** Heat shock proteins in toxicology: how close and how far? *Life Sci.* 86, 377–384.
- Gut, I.G. (2001)** Automation in genotyping of single nucleotide polymorphisms. *Hum. Mutat.* 17, 475–492.
- Hall, A.E., (2001)** Crop Responses to Environment. CRC Press LLC, Boca Raton, Florida.
- Halter GM., (2012)** Gene expression analysis of heat shock proteins in *Boechera Sparsiflora*, *Boechera pulchra* and *Boechera depauperata* when exposed to various degrees of heat stress.
- Hamam, K.A. and A.G.A. Khaled, (2009)** Stability of Wheat Genotypes under Different Environments and Their Evaluation under Sowing Dates and Nitrogen Fertilizer Levels *Australian Journal of Basic and Applied Sciences*, 3(1): 206-217 ISSN 1991-8178.
- Hammond-Kosack KE, Jones JDG (2000)** . Response to plant pathogens. In: Buchannan B, Gruissem W, Jones R, editors. *Biochemistry and molecular biology of plants*. Rockville, MD: American Society of Plant Physiologists; 2000. p. 1102-1157.
- Hare PD and WA Cress (1997)** Metabolic implications of stress induced proline accumulation in plants. *Plant Growth Regulation* 21(2): 79-102.
- Hare, P.D., Cress, W.A., Staden, J.V., (1998)** Dissecting the roles of osmolyte accumulation during stress. *Plant Cell Environ.* 21, 535–553.
- Hasan MA, JU Ahmed, MM Bahadur, MM Haque and S Sikder (2007).** Effect of late planting heat stress on membrane thermostability, proline content and heat susceptibility index of different wheat cultivars. *J. Natn. Sci. Foundation Sri Lanka* 35(2): 109-117.
- Helen S. (2013)** Chaperone machines for protein folding, unfolding and disaggregation *Nature Reviews Molecular Cell Biolog* 14, 630–642 (2013).
- Herms DA, Mattson WJ. (1992)** The dilemma of plants—to grow or defend. *Quarterly Review of Biology* 1992;67:283-335.
- Hogewoning, S.W., and Harbinson, J. (2007)** Insights on the development, kinetics, and variation of photoinhibition using chlorophyll fluorescence imaging of a chilled, variegated leaf. *Journal of Experimental Botany*, 58, 453-463, ISSN 0022-0957
- Hong SW, Vierling E (2001)** HSP101 is necessary for heat tolerance but dispensable for development and germination in the absence of stress. *The Plant Journal* 27, 25-35
- Hong, S.W. and Vierling, E. (2000)** Mutants of *Arabidopsis thaliana* defective in the acquisition of tolerance to high temperature stress. *Proc. Natl. Acad. Sci. U. S. A.* 97, 4392–4397
- Hossain MM, H Takeda and T Senboku (1995).** Proline content in *Brassica* under high temperature stress. *JIRCAS J.* 2: 87-93.

- Hossain, Z.; López-Climent, M.F.; Arbona, V.; Pérez-Clemente, R.M.; Gómez-Cadenas, A. (2009)** Modulation of the antioxidant system in *Citrus* under waterlogging and subsequent drainage. *J. Plant Physiol.* 2009, *166*, 1391–1404.
- Howarth, C.J. (2005).** Genetic improvements of tolerance to high temperature. In: Ashraf, M., Harris, P.J.C. (Eds.), *Abiotic Stresses: Plant Resistance Through Breeding and Molecular Approaches*. Howarth Press Inc., New York.
- Hu W, Hu G, Han B (2009)** Genome-wide survey and expression profiling of heat shock proteins and heat shock factors revealed overlapped and stress specific response under abiotic stresses in rice. *Plant Science* 176, 583-590
- Hughes T, Deininger M, Hochhaus A, Branford S, Radich J, Kaeda J, Baccarani M, Cortes J, Cross NC, Druker BJ, Gabert J, Grimwade D, Hehlmann R, Kamel-Reid S, Lipton JH, Longtine J, Martinelli G, Saglio G, Soverini S, Stock W, Goldman JM (2006)** Monitoring CML patients responding to treatment with tyrosine kinase inhibitors: review and recommendations for harmonizing current methodology for detecting BCR-ABL transcripts and kinase domain mutations and for expressing results. *Blood* 108:28–37
- Iba, K., (2002)** Acclimative response to temperature stress in higher plants: approaches of gene engineering for temperature tolerance. *Annu. Rev. Plant Biol.* 53, 225–245.
- Ibrahim, A.M H. and J. S. Quick. (2001)** Heritability of heat tolerance in winter and spring wheat. *Crop Sci.* 41: 1401-1405.
- Imelfort, M., Duran, C., Batley, J. and Edwards, D. (2009)** Discovering genetic polymorphisms in next-generation sequencing data. *Plant Biotechnol. J.* 7,312–317.
- Ismail, A.M., Hall, A.E., (1999)** Reproductive-stage heat tolerance, leaf membrane thermostability and plant morphology in cowpea. *Crop Sci.* 39, 1762–1768.
- Jackson-Constan, D., Akita, M., Keegstra, K., (2001).** Molecular chaperones involved in chloroplast protein import. *Biochim. Biophys. Acta* 1541, 102–113.
- Jehan, T. & Lakhanpaul, S. (2006).** Single nucleotide polymorphism (SNP)-Methods and applications in plant genetics: A review. *Indian Journal of Biotechnology* Vol.5, No.4, (October 2006), pp. 435-439. ISSN 0972-5849
- Johnston JA, Ward CL, Kopito RR: Aggresomes (1998)** A cellular response to misfolded proteins. *J Cell Biol* 1998, 143:1883-1898.
- Jones, P.D., New, M., Parker, D.E., Mortin, S., Rigor, I.G., (1999).** Surface area temperature and its change over the past 150 years. *Rev. Geophys.* 37, 173–199.
- Kanazin, V., Talbert, H., See, D., DeCamp, P., Nevo, E., and Blake, T. (2002).** Discovery and assay of single-nucleotide polymorphisms in barley (*Hordeum vulgare*). *Plant Mol. Biol.* 48: 529–537. Doi :10. 102 3/ A : 1 0 1 4859031781. PMID:11999833.
- Karim, M.A., Fracheboud, Y., Stamp, P., (1997).** Heat tolerance of maize with reference of some physiological characteristics. *Ann. Bangladesh Agri.* 7, 27–33.
- Karim, M.A., Fracheboud, Y., Stamp, P., (1999).** Photosynthetic activity of developing
- Katiyar-Agarwal S, Agarwal M, Grover A (2003)** Heat-tolerant basmati rice engineered by over-expression of *HSP101*. *Plant Molecular Biology* 51, 677- 686

- Katiyar-Agarwal, S., Agarwal, M., Gallie, D.R. and Grover, A. (2001)** Search for cellular functions of plant Hsp100/Clp family proteins. *Crit. Rev. Plant Sci.* 20, 277–295.
- Katiyar-Agarwal, S.; Agarwal, M.; Grover, A. (2003).** Heat-tolerant basmati rice engineered by over-expression of hsp101. *Plant Mol. Biol.* 2003, 51, 677–686.
- Kavi Kishore, P.B., Sangam, S., Amrutha, R.N., Laxmi, P.S., Naidu, K.R., Rao, K.R.S.S., Rao, S., Reddy, K.J., Theriappan, P., Sreenivasulu, N., (2005).** Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. *Curr. Sci.* 88, 424–438.
- Keeler SJ, Boettger CM, Haynes JG, Kuches KA, Johnson MM, Thureen DL, Keeler L Jr., Kitto SL (2000)** Acquired thermotolerance and expression of the *HSP100/ClpB* genes of Lima bean. *Plant Physiology* 123, 1121-1132
- Keresa, S., M. Baric, H. Sarcevic, S. Marchetti and G. Dresner. (2000).** Callus induction and plant regeneration from immature and mature embryos of winter wheat (*Triticum aestivum* L.) genotypes. *Plant Breeding Sustaining the Future.* XVIth EUCARPIA congress, Edinburgh, Scotland.
- Key, J.L., Lin, C.Y., Chen, Y.M., (1981).** Heat shock proteins of higher plants. *P Natl. Acad. Sci. USA* 78, 3526–3530.
- Khan, M.A., M.U. Shirazi, M.A. Khan, S.M. Mujtaba, E. Islam, S. Mumtaz, A. Shereen, R.U. Ansari and M.Y. Ashraf. (2009).** Role of proline, K/Na ratio and chlorophyll content in salt tolerance of wheat (*Triticum aestivum* L.). *Pak. J. Bot.* 41(2):633-638.
- Kherallah, M., N. Miont and P. Gruhn (1999).** Adjustment of wheat production to market reform in Egypt. MSSD Discussion Paper No. 32.
- Kimber, G. and Sears, E. R. (1987)** Evolution in the genus *Triticum* and the origin of cultivated wheat. Pages 154_164. In E. G. Heyne, ed. *Wheat and wheat improvement.* 2nd ed. Agron. Monogr. 13. ASA, CSSA, and SSSA. Madison, WI.
- Kinnersley, A.M., Turano, F.J., (2000).** Gamma aminobutyric acid (GABA) and plant responses to stress. *Crit. Rev. Plant Sci.* 19, 479–509.
- Kirakosyan, A.; Kaufman, P.; Warber, S.; Zick, S. (2004).** Aaronson, K.; Bolling, S.; Chul Chang, S. Applied environmental stresses to enhance the levels of polyphenolics in leaves of hawthorn plants. *Physiol. Plant.* 2004, 121, 182–186.
- Kirby, E.J.M. (2002)** Botany of the wheat plant. In: Curtis, B. C., S. Rajaram and H. G. Macpherson (Eds.) *Bread Wheat Improvement and Production*, FAO Plant Production and Protection Series, No. 30. Food and Agriculture Organization of the United Nations, Mexico, DF, Mexico.
- Kirk B W, Feinsod M, Favis R, Kleiman R M and Barany F. (2002).** Single nucleotide polymorphism seeking long-term association with complex disease, *Nucleic Acids Res*, 30 (2002) 3295-3311.

- Kishor, P.B.K.; Sangam, S.; Amrutha, R.N.; Laxmi, P.S.; Naidu, K.R.; Rao, K.R.S.S.; Rao, S.; Reddy, K.J.; Theriappan, P.; Sreenivasulu, N. (2005).** Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: Its implications in plant growth and abiotic stress tolerance. *Curr. Sci.* 2005, 88, 424–438.
- Kotak, S.; Larkindale, J.; Lee, U.; Koskull-Döring, P.; Vierling, E.; Scharf, K.-D. (2007).** Complexity of the heat stress response in plants. *Curr. Opin. Plant Biol.* 2007, 10, 310–316.
- Kou CG, HM Chen and LH Ma 1986.** Effect of high temperature on proline content in tomato floral buds and leaves. *J. Am. Soc. Hort. Sci.* 11: 734-750.
- Krause, G.H. (1988).** Photoinhibition of photosynthesis. An evaluation of damaging and protective mechanisms. *Physiologia Plantarum*, 74, 566-574, ISSN 0031-9317
- Kregel, K.C., (2002).** Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *J. Appl. Physiol.* 92, 2177–2186.
- Kumar S., Sharma V., Chaudhary S., Kumar A. and Kumari R. (2012)** Agronomic characteristics of autumn and winter seeded photoperiod in sensitive spring wheat in agro-climate of north-west India. *Proc. Ind. Natl. Sci. Acad.* 78, 71–89.
- Kumar Tewari A¹, Charan Tripathy B., (1998).** Temperature-stress-induced impairment of chlorophyll biosynthetic reactions in cucumber and wheat, *Plant Physiol.* 1998 Jul;117(3):851-8.
- Lagudah ES, Dubcovsky J, Powell W (2001)** Wheat genomics. *Plant Physiol Biochem* 39:335–344
- Large, E.C. (1954)** Growth stages in cereals illustration of the feekes scale. *Plant Pathol.* 3:128–129.
- Larkindale, J., Hall, J.D., Knight, M.R., Vierling, E., (2005).** Heat stress phenotypes of *Arabidopsis* mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiol.* 138, 882–897.
- Law, R.D. and C. Brandner. (2001).** High temperature stress increases the expression of wheat leaf ribulose-1, 5- bisphosphate carboxylase/oxygenase activase protein. *Arch. Biochem. Biophys.* 386: 261-267.
- Lee, S., Sowa, M.E., Chai, J.M. and Tsai, F.T.F. (2004)** The ClpB/Hsp104 molecular chaperone – a protein disaggregating machine. *J. Struct. Biol.* 146, 99–105.
- Lee, U., Vie, C., Escobar, M., Williams, B., Hong, S.W. and Vierling, E. (2005)** Genetic analysis reveals domain interactions of Arabidopsis HSP100/Clp and cooperation with the small heat shock protein chaperone system. *Plant Cell* 17, 559–571.
- Lichtenthaler, H.K., Ac, A., Marek, M.V., Kalina, J., & Urban, O. (2007).** Differences in pigment composition, photosynthetic rates and chlorophyll fluorescence images of sun and shade leaves of four tree species. *Plant Physiology and Biochemistry*, 45, 577- 588, ISSN 0981- 9428.
- Livak KJ, Schmittgen TD (2001)** Analysis of relative gene expression data using realtime quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods* 25: 402-408.

- Lobell, D.B., J.I. Ortiz-Monasterio, 2007.** Impacts of Day Versus Night Temperatures on Spring Wheat Yields. *Agron. J.* 99: 469-477.
- Lott N, Ross T, Smith A, Houston T, Shein K (2011).** Billion dollar U.S. Weather disasters, 1980 - 2010. National Climatic Data Center, Asheville, NC, USA. Available online at: <http://www.ncdc.noaa.gov/oa/reports/billionz.html>.
- Lubben, T.H., Donaldson, G.K., Viitanen, P.V., Gatenby, A.A., (1989).** Several proteins imported into chloroplasts form stable complexes with the GroEL-related chloroplast molecular chaperone. *Plant Cell* 1, 1223–1230.
- Luck J, Spackman M, Freeman A, Trebicki P, Griffiths W, Finlay K, Chakraborty S. (2011).** Climate change and diseases of food crops. *Plant Pathology* 2011;60:113-121.
- Lumpkin, T.A. (2011)** Wheat-global alliance for improving food security and the livelihoods of the resource- poor in the developing world. National Agricultural Research Institute.
- M. Krishnan, H. T. Nguyen, and J. J. Burke, (2002).** Heat shock protein synthesis and thermal tolerance in wheat,” *Plant Physiology*, vol. 90, no. 1, pp. 140–145, 1989.
- Mackay, I. M. 2002** Real-time PCR in virology, *Nucleic Acids Res.* , 30, 1292, 2002.
- Madan P, Jagadish SV, Craufurd PQ, Fitzgerald M, Lafarge T, Wheeler TR. (2012).** Effect of elevated CO₂ and high temperature on seed-set and grain quality of rice. *Journal of Experimental Botany* 63, 3843–3852.
- Madgwick JW, West JS, White RP, Semenov MA, Townsend JA, Turner JA, Fitt BD L. (2011).** Impacts of climate change on wheat anthesis and fusarium ear blight in the UK. *European Journal of Plant Pathology* 2011;130:117-131.
- Maestri, E., Klueva, N., Perrotta, C., Gulli, M., Nguyen, H.T., Marmiroli, N., (2002).** Molecular genetics of heat tolerance and heat shock proteins in cereals. *Plant Mol. Biol.* 48, 667–681.
- Mandhania S, Madan S, Sawhney V (2006)** Antioxidant defense mechanism under salt stress in wheat seedlings. *Biol Plant* 50:227-231
- Maraseni, T. N., Mushtaq, S. & Maroulis, J. (2009)** Greenhouse gas emissions from rice farming inputs: a cross-country assessment. *Journal of Agricultural Science, Cambridge* 147, 117–126.
- Marchand, F.L., Mertens, S., Kockelbergh, F., Beyens, L., Nijs, I., (2005).** Performance of high arctic tundra plants improved during but deteriorated after exposure to a simulated extreme temperature event. *Global Change Biol.* 11, 2078–2089.
- Marcum, K.B., (1998)** Cell membrane thermostability and whole plant heat tolerance of Kentucky bluegrass. *Crop Sci.* 38, 1214–1218.
- Mariolina Gulli, Massimiliano Corradia, Patrizia Rampinob, Nelson Marmiroli and Carla Perrotta (2007).** Four members of the HSP101 gene family are differently regulated in *Triticum durum* Desf. 581 (2007) 4841–4849

- Martineau, J.R., Specht, J.E., Williams, J.H., Sullivan, C.Y., (1979).** Temperature tolerance in soybean. I. Evaluation of technique for assessing cellular membrane thermostability. *Crop Sci.* 19, 75–78.
- Massacci, A., Nabiev, S.M., Pietrosanti, L., Nematov, S.K., Chernikova, T.N., Thor, K., & Leipner, J. (2008).** Response of the photosynthetic apparatus of cotton (*Gossypium*)
- McMaster, G.S., (1997).** Phonology, development, and growth of wheat (*Triticum aestivum* L.) shoot apex: A 26. Sofield, I., L.T. Evans and I.F. Wardlaw, 1977. The review. *Adv. Agron.*, 59: 63-118.
- Miernyk, J.A., (1999)** Protein folding in the plant cell. *Plant Physiol.* 121, 695–703.
- Miller, G. and Mittler, R. (2006)** Could heat shock transcription factors function as hydrogen peroxide sensors in plants? *Ann. Bot.* 2006, 98, 279–288.
- Mittler R and Blumwald E. (2010)** Genetic engineering for modern agriculture: challenges and perspectives. *Annual Review of Plant Biology* 2010;61:443-462
- Moaed Almeselmani, P. S. Deshmukh and Viswanathan Chinnusamy (2012).** Effects of Prolonged High Temperature Stress on Respiration, Photosynthesis and Gene Expression in Wheat (*Triticum aestivum* L.) Varieties Differing in their Thermotolerance. Melbourne, pp. 31–61.
- Mochida, K. and Shinozaki, K. (2010)** Genomics and bioinformatics resources for crop improvement. *Plant Cell Physiol.* 51, 497–523.
- Mogk, A. et al. (2003)** Refolding of substrates bound to small Hsps relies on a disaggregation reaction mediated most efficiently by ClpB/ DnaK. *J. Biol. Chem.* 278, 31033–31042
- Mohammadreza Amirjani, (2012)** Estimation of wheat responses to "high" heat stress, *American-Eurasian Journal of Sustainable Agriculture*, 6(4): 222-233, 2012
- Momcilovic, I., Ristic, Z., 2007.** Expression of chloroplast protein synthesis elongation factor, EF-Tu, in two lines of maize with contrasting tolerance to heat stress during early stages of plant development. *J. Plant Physiol.* 164, 90–99.
- Morison JIL, Baker NR, Mullineaux PM, Davies WJ (2008)** . Improving water use in crop production. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2008;363:639-658.
- Mostafa, H.A.M., R.A. Hassanein, S.I. Khalil, S.A. El-Khawas, H.M.S. El-Bassiouny, A.A. Abd El- Monem, (2010).** Effect of arginine or putrescine on growth, yield and yield components of late sowing wheat. *J. of Applied Sci. Res.*, 6(2): 177-183.
- Nakamoto, H. and Vigh, L., (2007)** The small heat shock proteins and their clients. *Cell Mol. Life Sci.* 64, 294–306.
- Neill, Richard. (2002)** *Booze: The Drinks Bible for the 21st Century*. Octopus Publishing Group - Cassell Illustrated. p. 112. ISBN 1-84188-196-1.
- Neilson, K.A. and Gammulla, G. (2010)** Mirzaei, M.; Imin, N.; Haynes, P.A. 2010 Proteomic analysis of temperature stress in plants. *Proteomics* 2010, 10, 828–845.

- Newton AC, Johnson SN, Gregory PJ . (2011).** Implications of climate change for diseases, crop yields and food security. *Euphytica* 2011;179:3-18
- Nicod J and Largiadere C R, (2003)** SNPs by AFLP (SBA): A rapid SNP isolation strategy for non-model organisms, *Nucleic Acids Res*, 31 (2003) e19.
- Nicol JM, Turner SJ, Coyne DL, denNijs L, Hockland S, TahnaMaafi Z.(2011).** Current nematode threats to world agriculture. In: Jones J, Gheysen G, Fenoll C , editors. *Genomics and molecular genetics of plant–nematode interactions*. London: Springer;2011. p. 21-44.
- Nieto-Sotelo J, Martinez LM, Ponce G, Cassab GI, Alagon A, Meeley RB, Nover, L., Baniwal, S.K., (2006).** Multiplicity of heat stress transcription factors controlling the complex heat stress response of plants. In: International Symposium on Environmental Factors, Cellular Stress and Evolution, Varanasi, India, October 13–15, 2006, p. 15.
- Nordborg, M., et al. (2005).** The pattern of polymorphism in *Arabidopsis thaliana*. *PLoS Biol.* 3: e196.
- Osmond, B., Schwartz, O., & Gunning, B. (1999).** Photoinhibitory printing on leaves, visualized by chlorophyll fluorescence imaging and confocal microscopy, is due to diminished fluorescence from grana. *Australian Journal of Plant Physiology*, 26,717– 724., ISSN, 0310-7841
- Ouda, S.A., S.M. El-Marsafawy, M.A. El-Kholy and M.S. Gaballah, (2005).** Simulating the effect of Water Stress and Different Sowing Dates on Wheat Production in South Delta *Journal of Applied Sciences Research*, 1(3): 268-276.
- Overbergh, L., Valckx, D., Waer, D., and Mathieu, C. (1999)** Quantification of murine cytokine mRNAs using real time quantitative reverse transcriptase PCR. *Cytokine* 11 (4), 305–312
- P. Rampino, S. Pataleo, M. De Pascali, C. Gerardi, G.Mita, N. Di Fonzo, C. Perrotta**
P. Rampino, S. Pataleo, M. De Pascali, C. Gerardi, G.Mita, N. Di Fonzo and C. Perrotta (2007) Characterisation of Different Durum Wheat (*Triticum durum* Desf.) Cultivars for Thermotolerant 61, 199-223.
- Paleg LG, TJ Douglas, A Van Daal and DB Keech (1981).** Proline, betaine and other organic solutes protect enzymes against heat inactivation. *Aust. J. Plant Physiol.* 8: 107-114.
- Palmer, John J. (2001)** *How to Brew*. Defenestrative Pub Co. p. 233. ISBN 0-9710579-07.
- Panaretou B, Zhai C (2008)** The heat shock proteins: Their roles as multi-component machines for protein folding. *Fungal Biology Reviews* 22, 110-119
- Panaretou, B., Zhai, C., (2008).** The heat shock proteins: their roles as multi-component machines for protein folding. *Fungal biol. rev.* 22, 110–119.
- Parankiewicz, J., Wang, J. and Clarke, A.K. (1999)** New insights into the ATP-dependent Clp protease: *Escherichia coli* and beyond. *Mol. Microbiol.* 32, 449–458.
- Pareek, A., Singla, S.L., Grover, A., (1998).** Plant Hsp90 family with special reference to rice. *J. Biosci.* 23, 361–367.

- Park J, H Ro, K Hwang and MS Yiem (2001).** Effect of water stress induced by polyethylene glycol and root zone temperature on growth and mineral contents of Fuji/M. 26 apple. *J. Korean Soc. Hort. Sci.* 42(4): 435-438.
- Park, H.-S.; Jeong, W.-J.; Kim, E.-C.; Jung, Y.; Lim, J.M.; Hwang, M.S.; Park, E.-J.; Ha, D.-S.; Choi, D.-W. (2012).** Heat shock protein gene family of the *Porphyra seriata* and enhancement of heat stress tolerance by PsHSP70 in *Chlamydomonas*. *Mar. Biotechnol.*
- Parsell DA, Kowal AS, Singer MA, Lindquist S (1994).** Protein disaggregation mediated by heat-shock protein HSP104. *Nature* 372, 475-478
- Parsell, P.A., Lindquist, S., (1993).** The function of heat-shock proteins in stress tolerance. degradation and reactivation of damaged proteins. *Annu. Rev. Genet.* 27, 437-496.
- Peet, M.M., Willits, D.H., (1998).** The effect of night temperature on greenhouse grown tomato yields in warm climate. *Agric. Forest Meteorol.* 92, 191-202.
- Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X. (2004).** Rice yields decline with higher night temperature from global warming. *Proc Natl Acad Sci* 2004;101:9971-5.
- Peres Ben-Zvi, A. and Goloubinoff, P. (2001)** Mechanisms of disaggregation and refolding of stable protein aggregates by molecular chaperones. *J. Biol. Chem.* 135, 84-93
- Perkel, J. (2008)** SNP genotyping: Six technologies that keyed a revolution. *Nat. Methods*, 5, 447-453.
- Porter, D.R., H.T. Nguyen and J.J. Burke. (1995).** Genetic control of acquired high temperature tolerance in winter wheat. *Euphytica.* 83: 153-157.
- Porter, J.R., (2005)** Rising temperatures are likely to reduce crop yields. *Nature* 436, 174.
- Pradhan, G.P., P.V.V. Prasad, A.K. Fritz, M.B. Kirkham & B.S. Gill .(2012).** Effects of drought and high temperature stress on synthetic hexaploid wheat. *Functional Plant Biology*, 39: 190-198.
- Prasad, P.V.V., S.R. Pisipati, Z. Ristic, U. Bukovnik & A.K. Fritz, (2008).** Impact of Nighttime Temperature on Physiology and Growth of Spring Wheat. *Crop Sci.*, 48: 2372-2380.
- Pratt, W.B., Galigniana, M.D., Harrell, J.M., Deranco, D.B., (2004).** Role of hsp90 and the hsp90-binding immunophilins in signaling protein movement. *Cell Signal.* 16, 857-872.
- Pratt, W.B., Toft, D.O., (2003).** Regulation of signaling protein function and trafficking by the hsp90/hsp70-based chaperone machinery. *Exp. Biol. Med.* 228, 111-133.
- Quan, R., Shang, M., Zhang, H., Zhao, Y., Zhang, J., (2004).** Engineering of enhanced glycine betaine synthesis improves drought tolerance in maize. *Plant Biotech. J.* 2, 477-486.
- Queitsch C, Hong SW, Vierling E, Lindquist S (2000)** Heat shock protein 101 plays a crucial role in thermotolerance in *Arabidopsis*. *Plant Cell* 12, 479-492

- Rafalski A, (2002a)** Applications of single nucleotide polymorphisms in crop genetics. *Curr Opin Plant Biol* 5:94–100.
- Rafalski, J.A. (2002b).** Novel genetic mapping tools in plants: SNPs and LD-based approaches. *Plant Sci.* 162, 329–333.
- Rasheed, I. M., Aslam, D. H., & Sarwar, S. (2010).** Motivational Issues for Teachers in Higher Education: A Critical Case of IUB. *Journal of Management Research* ISSN 1941-899X 2010, Vol. 2, No. 2: E3.
- Rayan, A.A., S.M. El-Marsafawy and K.A. Mohamed, (1999).** Response of some wheat varieties to different sowing dates and irrigation regimes in Upper Egypt. 3rd Conference of On-Farm Irrigation and Agro climatology, SWERI, ARC Egypt
- Remington, D.L., Thornsberry, J.M., Matsuoka, Y., Wilson, L.M., Whitt, S.R., Doebley, J., Kresovich, S., Goodman, M.M., and Buckler IV, E.S. (2001).** Structure of linkage disequilibrium and phenotypic associations in the maize genome. *Proc. Natl. Acad. Sci. USA* 98: 11479–11484.
- Renu, M., S.S. Dhanda, R.K. Rana and I. Singh. (2004).** Membrane thermostability as an indicator of heat tolerance at seedling stage in bread wheat. *Nat. J.Pl.Improv.*6 (2): 133-135.
- Reynolds, M.P. and R.M. Trethowan, (2007).** Physiological interventions in breeding for adaptation to abiotic stress. In: *Scale and complexity in plant system research: Geneplant crop relations* (Eds: Spiertz JHJ, Struk PC & Van Laar HH) Wageningen UR Frontiers series Springer. Pp. 129- 146
- Reynolds, M.P., M.I. Delgado B, M. Gutiérrez-Rodríguez, and A. Larqué-Saavedra. (2000).** Photosynthesis of wheat in a warm, irrigated environment: I: Genetic diversity and crop productivity. *Field Crops Res.* 66:37–50.
- Ribaut JM, Yang R (2002)** Maize HSP101 plays important roles in both induced and basal thermotolerance and primary root growth. *Plant Cell* 14, 1621-1633
- Ristic, Z., U. Bukovnik, and P.V.V. Prasad. (2007).** Correlation between heat stability of thylakoid membranes and loss of chlorophyll in winter wheat under heat stress. *Crop Sci.* 47:2067–2073.
- Ritossa F. (1962).** A new puffing pattern induced by temperature shock and DNP in *Drosophila*. *Experientia.* ;18:571–573. doi: 10.1007/BF02172188.
- Rockstrom J, Falkenmark M (2000).** Semiarid crop production from a hydrological perspective: gap between potential and actual yields. *Critical Reviews in Plant Sciences*2000;19:319-346
- Ronde J, A Mescht and HSF Steyn (2001).** Proline accumulation in response to drought and heat stress in cotton. *African Crop Sci. J.* 8(1): 85-91.
- Rosegrant, M.W. and Agcaoili, M. (2010),** Global Food Demand, Supply, and Price Prospects to 2010. International Food Policy Research Institute: Washington, DC.
- Roussel, V., Koenig, J., Beckert, M., and Balfourier, F. (2004).** Molecular diversity in French bread wheat accessions related to temporal trend and breeding programmes. *Theor. Appl. Genet.* 108: 920–930. doi:10.1007/s00122-003-1502-y. PMID:14614567.

- Rouch, J.M., Bingham, S.E., Sommerfeld, M.R., (2004).** Protein expression during heat stress in thermo-intolerance and thermotolerance diatoms. *J. Exp. Mar. Biol. Ecol.* 306, 231–243.
- Routaboul, J. M., S. F. Fischer, and J. Browse. (2000).** Trienoic fatty acids are required to maintain chloroplast function at low temperatures. *Plant Physiol.* 124:1697–1705.
- Ruelland E, Zachowski A (2010).** How plants sense temperature, *Environmental and Experimental Botany* 69(3): 225-232.
- Saadalla, M. M., J. F. Shanahan and J. S. Quick. (1990a).** Heat tolerance in winter wheat. I. Hardening and genetic effects on membrane thermostability. *Crop Sci.* 30: 1243-1247.
- Saadalla, M. M., J. F. Shanahan and J. S. Quick. (1990b).** Heat tolerance in winter wheat.II. Membrane thermostability and field performance. *Crop Sci.* 30: 1248-1251.
- Sabehat, A. and Weiss, D. (1998)** Lurie, S. Heat-shock proteins and cross-tolerance in plants. *Physiol. Plant.* 1998, 103, 437–441.
- Saidi, Y. et al. (2009)** The heat shock response in moss plants is regulated by specific calcium-permeable channels in the plasma membrane. *Plant Cell* 21, 2829–2843
- Saidi, Y. et al. (2010)** Membrane lipid composition affects plant heat sensing and modulates Ca²⁺-dependent heat shock response. *Plant Signal. Behav.* 5, 1530–1533
- Sairam, R.K., Tyagi, A., (2004).** Physiology and molecular biology of salinity stress tolerance in plants. *Curr. Sci.* 86, 407–421.
- Sakamoto, A., Murata, N., (2002).** The role of glycine betaine in the protection of plants from stress: clues from transgenic plants. *Plant Cell Environ.* 25, 163–171.
- Salvucci, M.E., Crafts-Brandner, S.J., (2004).** Inhibition of photosynthesis by heat stress: the activation state of Rubisco as a limiting factor in photosynthesis. *Physiol Plant* 120: 179–186
- Sato, S., Kamiyama, M., Iwata, T., Makita, N., Furukawa, H., Ikeda, H., (2006).** Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. *Ann. Bot.* 97, 731–738.
- Savchenko, G.E., Klyuchareva, E.A., Abrabchik, L.M., Serdyuchenko, E.V., (2002).** Effect of periodic heat shock on the membrane system of etioplasts. *Russ. J. Plant Physiol.* 49, 349–359.
- Schöffl, F., Prandl, R., Reindl, A., (1999).** Molecular responses to heat stress. In: Shinozaki, K., Yamaguchi-Shinozaki, K. (Eds.), *Molecular Responses to Cold, Drought, Heat and Salt Stress in Higher Plants*. R.G. Landes Co., Austin, Texas, pp. 81–98.
- Schirmer, E.C. et al. (1996)** Hsp100/Clp proteins: a common mechanism explains diverse functions. *Trends Biochem. Sci.* 21, 289–296
- Schlesinger, M.J., (1990).** Heat shock proteins. *J. Biol. Chem.* 265,12111–12114.

- Schoffl, F., Prandl, R., Reindl, A., (1999).** Molecular responses to heat stress. In: Shinozaki, K., Yamaguchi-Shinozaki, K. (Eds.), *Molecular Responses to Cold, Drought, Heat and Salt Stress in Higher Plants*. R.G. Landes Co., Austin, Texas, pp. 81–98.
- Scholes, J.D., & Rolfe, S.A. (1996).** Photosynthesis in localized regions of oat leaves infected with crown rust (*Puccinia coronata*). Quantitative imaging of chlorophyll fluorescence. *Planta*, 199, 573-582, ISSN, 0032-0943
- Scholz, M.; Gatzek, S.; Sterling, A.; Fiehn, O.; Selbig, J. (2004).** Metabolite fingerprinting: Detecting biological features by independent component analysis. *Bioinformatics* 2004, 20, 2447–2454.
- Schroda, M., Vallon, V., Wollman, F., Beck, C.F., (1999).** A chloroplast-targeted heat shock protein 70 (HSP70) contributes to the photoprotection and repair of photosystem II during and after photoinhibition. *Plant Cell* 11, 11165–11178.
- Seo, J.S., Lee, Y.M., Park, H.G., Lee, J.S., (2006).** The inter tidal copepod *Tigriopus japonicus* small heat shock protein 20 gene (Hsp20) enhances thermotolerance of transformed *Escherichia coli*. *Biochem. Biophys. Res. Commun.* 340, 901–908.
- Shakun, J.D., P.U. Clark, F. He, S.A. Marcott, A.C. Mix, Z. Liu, B. Otto-Bliesner, A. Schmittner & E. Bard, (2012).** Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature*. 2012 Apr 4;484(7392):49-54. doi: 10.1038/nature10915.
- Shamsi Keyvan, (2010).** The effects of drought stress on yield, relative water content, proline, soluble carbohydrates and chlorophyll of bread wheat cultivars *Journal of Animal & Plant Sciences*, 2010. Vol. 8, Issue 3: 1051- 1060.
- Shanahan, J. F., I. B. Edwards, J. S. Quick and J. R. Fenwick. (1990).** Membrane thermostability and heat tolerance of spring wheat. *Crop Sci.* 30: 247-251.
- Shao HB, Chu LY, Jaleel CA, Zhao CX (2008)** Water-deficit stress-induced anatomical changes in higher plants. *Comptes Rendus Biologies* 2008;331:215-225.
- Shao, H.B., Guo, Q.J., Chu, L.Y., Zhao, X.N., Su, Z.L., Hu, Y.C., Cheng, J.F., (2007).** Understanding molecular mechanism of higher plant plasticity under abiotic stress. *Colloids Surf. B: Biointerfaces* 54, 37–45.
- Sharon J. Keeler and co-workers (2000)** Acquired thermotolerance and expression of the HSP100/ClpB genes of Lima bean. *Plant Physiol.* 123, 1121–1132
- Sheikh S, Behl R K, Dhanda SS, Munjal R (2010)** Membrane thermostability and chlorophyll fluorescence as indices of high temperature stress tolerance and performance in wheat (*Triticum aestivum* L.). *Cereal Res Commun* 38(3):335-344
- Shewry PR. (2007)** Improving the protein content and composition of cereal grain. *Journal of Cereal Science*, 46: 239–250.
- Shewry PR, (2009)** The HEALTHGRAIN programme opens new opportunities for improving wheat for nutrition and health. *Nutrition Bulletin*, 34(2): 225–231.
- Shitsukawa, N., A. Takagishi, C. Ikari, S. Takumi, and K. Murai (2006).** WFL, a wheat FLORICAULA/LEAFY ortholog, is associated with spikelet formation as lateral branch of the inflorescence meristem. *Genes Genet. Syst.* 81:3–20.

- Shitsukawa, N., H. Kinjo, S. Takumi, and K. Murai. (2009).** Heterochronic development of the floret meristem determines grain number per spikelet in diploid, tetraploid and hexaploid wheats. *Ann. Bot.* 104:243–251.
- Siam Gamal M. (2006).** An assessment of the impact of increasing wheat self-sufficiency and promoting cash-transfer subsidies for consumers in Egypt: A multi-market; model ESA Working Paper No. 06-03 February 2006.
- Simmonds DH, (1989).** Inherent Quality Factors in Wheat. Wheat and Wheat Quality in Australia. Australia Wheat Board,
- Slafer, G.A., and H.M. Rawson. (1994).** Sensitivity of wheat phasic development to major environmental factors: A re-examination of some assumptions made by physiologists and modellers. *Aust. J. Plant Physiol.* 21:393–426.
- Smith, Albert E. (1995)** *Handbook of Weed Management Systems*. Marcel Dekker. p. 411. ISBN 0-8247-9547-4
- Smith, D. B. and Flavell, R. B. (1975).** Characterization of the wheat genome by renaturation kinetics. *Chromosoma* 50: 223_242.
- Smith, P. & OLESEN, J. E. (2010)** Synergies between the mitigation of, and adaptation to, climate change in agriculture. *Journal of Agricultural Science, Cambridge* 148, 543–552.
- Song, H.M.; Wang, H.Z.; Xu, X.B. (2012)** Overexpression of AtHsp90.3 in *Arabidopsis thaliana* impairs plant tolerance to heavy metal stress. *Biol. Plant.* 2012, 56, 197–199.
- Song, J., Y. Xu, S. White, K. W. P. Miller, and M. Wolinsky. (2005).** SNPsFinder--a web-based application for genome-wide discovery of single nucleotide polymorphisms in microbial genomes. *Bioinformatics* 21:2083-2084.
- Stevenson D, Doherty R, Sanderson M, Johnson C, Collins B, Derwent D. (2005).** Impacts of climate change and variability on tropospheric ozone and its precursors. *Faraday Discussions* 130, 41–57.
- Stewart GR and JA Lee (1974).** The role of proline accumulation in halophytes. *Planta* 120: 279-289.
- Stewart GR and SF Boggess (1978).** Metabolism of 5-3H proline by barley leaves and its use in measuring the effect of water stress on proline oxidation. *Plant Physiol.* 61: 654-657.
- Stoskopf, N. C. (1985).** Cereal grain crops. Pp. 187-200 and 388-400. Reston Publishing Co., Inc. Reston, Virginia. Sustainability. *Cur. Sci.* 87: 1334-1335.
- Strand, M., & Öquist, G. (1985).** Inhibition of photosynthesis by freezing temperatures and high light levels in cold-acclimated seedlings of Scots pine (*Pinus sylvestris*). - II. Effects on chlorophyll fluorescence at room temperature and 77 K. *Physiologia Plantarum*, 65, 117-123, ISSN 0031-9317
- Stupar, R.M. and Springer, N.M. (2006)** Cis-transcriptional variation in maize inbred lines B73 and Mo17 leads to additive expression patterns in the F1 hybrid. *Genetics*, 173, 2199–2210.

- Su, P.-H., Li, H.-m., (2008).** *Arabidopsis* stromal 70-kD heat shock proteins are essential for plant development and important for thermotolerance of germinating seeds. *Plant Physiol.* 146, 1231–1241.
- Sullivan, C.Y. and W.M. Ross. (1979).** Selecting for drought and heat resistance in grain sorghum. In Mussell, H. and Staples, R (eds) *Stress physiology in Crop plants*. New York, USA: John Wiley.
- Sun WN, Van Montagu M, Verbruggen N. (2002)** Small heat shock proteins and stress tolerance in plants. *BBA-Gene Struct Expr* 2002, 1577:1-9.
- Sung, D.-Y., Kaplan, F., Lee, K.-J., Guy, C.L., (2003).** Acquired tolerance to temperature extremes. *Trends Plant Sci.* 8, 179–187.
- Sung, D.Y.; Guy, C.L. (2003)** Physiological and molecular assessment of altered expression of Hsc70-1 in *Arabidopsis*. Evidence for pleiotropic consequences. *Plant Physiol.* 2003, 132, 979–987.
- Swindell, W.R.; Huebner, M.; Weber, A.P. (2007)** Transcriptional profiling of *Arabidopsis* heat shock proteins and transcription factors reveals extensive overlap between heat and non-heat stress response pathways. *BMC Genomics* 2007, 8, 1–15.
- Tabassum Jehan and Suman Lakhanpaul. (2006)** Single nucleotide polymorphism (SNP)–Methods and applications in plant genetics: A review
- Tahir, M. and M. Singh. (1993)** Assessment of screening techniques for heat tolerance in wheat. *Crop Sci.* 33: 740-744.
- Taiz, L., and E. Zeiger. (2006)** *Stress physiology*. 4th ed. Sinauer Associates, Inc., MA, USA. pp. 671–702.
- Taiz, L., Zeiger, E., (2006)** *Plant Physiology*. Sinauer Associates Inc. Publishers, Massachusetts.
- Takeda H, U Cempukelee, YS Chauhan, A Srinivasan, MM Hossain, MH Rashad, B Lin, HS Tolwar, T Senboku, S Varhima, S Yanagihara, M Shono, S Ancho, BQ Lin, M Yajima and T Hayashi (1999).** Studies in heat tolerance of *Brassica* vegetables and legumes at the International Collaboration Research Station from 1992 to 1996. Proc. Workshop on heat tolerance of crops. Okinawa. Japan, 7-9 October 1997. JIRCAS working Report 14: 17-29.
- Takeda S, Matsuoka M. (2008)** Genetic approaches to crop improvement: responding to environmental and population changes. *Nature Reviews Genetics* 2008;9:444-457.
- Tardy, A. Creach, and M. Havaux, (1998).** “Photosynthetic pigment concentration. Organization and enter conversions in a pale green Syrian landrace of barley adapted to harsh climatic conditions”. *Plant Cell Environment*, 21: 479-489, 1998.
- Thao, N.P., Chen, L., Nakashima, A., Hara, S., Umemura, K., Takahashi, A., Shirasu, K., Kawasaki, T., Shimamoto, K., (2007).** RAR1 and HSP90 form a complex with Rac/Rop GTPase and function2 in innate-immune responses in rice. *Plant Cell* 19, 4035–4045.

- Tenaillon, M.I., Sawkins, M.C., Long, A.D., Gaut, R.L., Doebley, J.F., and Gaut, B.S. (2001).** Patterns of DNA sequence polymorphism along chromosome 1 of maize (*Zea mays* ssp. *mays* L.). *Proc. Natl. Acad. Sci. USA* 98: 9161–9166.
- Theocharis, A.; Clément, C.; Barka, E.A. (2012)** Physiological and molecular changes in plants grown at low temperatures. *Planta* 2012, 235, 1091–1105.
- Tewari, A.K., Tripathy, B.C. (1998)** Temperature-stress-induced impairment of chlorophyll biosynthetic reactions in cucumber and wheat. – *Plant Physiol.* 117: 851-858, 1998.
- Timperio AM., Anna M., Maria G., (2008).** Proteomics applied on plant abiotic stresses: Role of heat shock proteins (HSP). *Journal of proteomics* 71 (2008) 391– 411
- Tissieres, A., Mitchell, H.K., Tracy, U.M., (1974).** Protein synthesis in salivary glands of *D. Melanogaster*. Relation to chromosome puffs. *J. Mol. Biol.* 84, 389–398.
- Tonsor SJ, Scott C, Boumaza I, Liss TR, Brodsky JL, Vierling E. (2008)** Heat shock
- Treichel S (1975)** The effect of NaCl on the concentration of proline in different halophytes. *Zestschrifti Pflanzenphysiologie* 76: 56-68.
- Trevaskis B. (2010)** The central role of the *VERNALIZATION1* gene in the vernalization of cereals. *Funct. Plant Biol.* 37, 479– 487.
- Trevaskis B., Hemming M. N., Peacock W. J. and Dennis E. S. (2007)** The molecular basis of vernalization-induced flowering in cereals. *Trends Plant Sci.* 12, 352–357.
- Tripp J, Mishra SK, Scharf KD (2009)** Functional dissection of the cytosolic chaperone network in tomato mesophyll protoplasts. *Plant Cell Environment* 32, 123-133
- Tripp, J., Mishra, S.K., Scharf, K.-D., (2009)** Functional dissection of the cytosolic chaperone network in tomato mesophyll protoplasts. *Plant Cell Environ.* 32, 123–133.
- Troll W. and Lindsey J. (1955),** A photometric method for determination of proline. *J. Biol. Chem.* 215, 655-660.
- Tubiello, F.N., Rosenzweig, C., Goldberg, R.A., Jagtap, S., Jones, J.W., (2000)** U.S. National Assessment Technical Report Effects of Climate Change on U.S. Crop Production Part I: Wheat, Potato, Corn, and Citrus.
- Tyedmers J., Mogk A., Bukau B. (2010).** Cellular strategies for controlling protein aggregation. *Nat. Rev. Mol. Cell Biol.* 11 777–788.
- United Nations Population Division Department of Economic and Social Affairs (DoEaSA) (2009)** World Population Prospects: the 2008 Revision. Available at: <http://esa.un.org/unpp> (accessed 3 August 2009).
- Vavilov, N. I. (1951).** The origin, variation, immunity, and breeding of cultivated plants. *Chronica Bot.* 13: 1_366.
- Verbruggen, N.; Hermans, C. Proline accumulation in plants (2008)** A review. *Amino Acids* 2008, 35, 753–759.
- Verslues, P.E.; Agarwal, M.; Katiyar-Agarwal, S.; Zhu, J.; Zhu, J.-K. (2006).** Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J.* 2006, 45, 523–539.

- Vierling, E. (1991)** The roles of heat shock proteins in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 1991, 42, 579–620.
- Vinocur, B.; Altman, A. (2005)** Recent advances in engineering plant tolerance to abiotic stress: Achievements and limitations. *Curr. Opin. Biotechnol.* 2005, 16, 123–132.
- Visser, E.J.W. (2005)** Voeselek, L.A.C.J. Acclimation to soil flooding-sensing and signal-transduction. *Plant Soil* 2005, 274, 197–214.
- Von Koskull-Doering, P., Scharf, K.D., Nover, L., (2007)** The diversity of plant heat stress transcription factors. *Trends Plant Sci.* 12, 452–457.
- Wahid, A., (2007)** Physiological implications of metabolites biosynthesis in net assimilation and heat stress tolerance of sugarcane sprouts. *J. Plant Res.* 120, 219–228.
- Wahid, A., Close, T.J., (2007).** Expression of dehydrins under heat stress and their relationship with water relations of sugarcane leaves. *Biol. Plant.* 51, 104–109.
- Wahid, A., Ghazanfar, A., (2006).** Possible involvement of some secondary metabolites in salt tolerance of sugarcane. *J. Plant Physiol.* 163, 723–730.
- Wahid, A., S. Gelani, M. Ashraf, and M. Foolad. (2007).** Heat tolerance in plants: An overview. *Environ. Exp. Bot.* 61:199–223.
- Wahid, A., Shabbir, A., (2005).** Induction of heat stress tolerance in barley seedlings by pre-sowing seed treatment with glycinebetaine. *Plant Growth Reg.* 46, 133–141.
- Wakeley J. (1996).** The excess of transitions among nucleotide substitutions: New methods of estimating transition bias underscore its significance. *TREE* 11:158–163
- Wang Y., M. Wisniewski, R. Meilan, M. Cui, L. Fuchigami. (2006).** Transgenic tomato (*Lycopersicon esculentum*) overexpressing cAPX exhibits enhanced tolerance to UV-B and heat stress, *J. Appl. Horticult.* 8 (2006) 87e90.
- Wang, W., Vinocur, B., Altman, A., (2003).** Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218, 1–14.
- Wang, W.; Vinocur, B.; Shoseyov, O.; Altman, A. (2004)** Role of plant heat shock proteins and molecular chaperones in the abiotic stress response. *Trends Plant Sci.* 2004, 9, 244–252.
- Wettstein, D., (1957)** Chlorophyll-letale und der Submikroskopische Formwechsel der Plastiden. *Experimental Cell Research* 12, 427–506.
- Wheeler, T.R., P.Q. Craufurd, R.H. Ellis, J.R. Porter & P.V. Vara Prasad, (2000).** Temperature variability and the yield of annual crops. *Agriculture, Ecosystems & Environment.*, 82: 159-167.
- Wiese, M. V. (1987)** Compendium of wheat diseases, 2nd ed. Pp. 1-9. APS Press, St. Paul, MN.
- Win J., Kamoun S., Jones A.M. (2011).** Purification of effector-target protein complexes via transient expression in *Nicotiana benthamiana*. *Methods Mol. Biol.* 712: 181–194

- Wise, R.R., A.J. Olson, S.M. Schrader, and T.D. Sharkey. (2004).** Electron transport is the functional limitation of photosynthesis in field-grown pima cotton plants at high temperature. *Plant Cell Environ.* 27:717–724.
- Wolters P, Powell W, Lagudah E, Snape J, et al (2000).** Nucleotide Diversity at Homologous Loci in Wheat. In: *Plant and Animal Genome VIII Conference*, San Diego, 9-12.
- Woo, N.S., Badger, M.R., & Pogson, B.J. (2008).** A rapid, non-invasive procedure for quantitative assessment of drought survival using chlorophyll fluorescence. 11;4:27. doi: 10.1186/1746-4811-4-27.
- Xu-QA., (1991)** Biochemical and ultra structural changes in wheat (*Triticum aestivum*) plant exposed to high temperature and phytohormones. *Dissertation-Abstracts International. B. Science-and Engineering. U.S.A.* Pp.171.
- Yamada, A.; Sekiguchi, M.; Mimura, T.; Ozeki, Y. (2002)** The role of plant CCT α in salt- and osmotic-stress tolerance. *Plant Cell Physiol.* 2002, 43, 1043–1048.
- Yamada, M., Hidaka, T., Fukamachi, H., (1996)** Heat tolerance in leaves of tropical fruit crops as measured by chlorophyll fluorescence. *Sci. Hort.* 67, 39–48.
- Yang, J., R.G. Sears, B.S. Gill, and G.M. Paulsen (2002)** Growth and senescence characteristics associated with tolerance of wheat-alien amphiploids to high temperature under controlled conditions. *Euphytica* 126:185–193.
- Yang, X.; Chen, X.; Ge, Q.; Li, B.; Tong, Y.; Zhang, A.; Li, Z.; Kuang, T.; Lu, C. (2006)** Tolerance of photosynthesis to photoinhibition, high temperature and drought stress in flag leaves of wheat: A comparison between a hybridization line and its parents grown under field conditions. *Plant Sci.* 2006, 171, 389–397.
- Ye, S.F.; Yu, S.W.; Shu, L.B.; Wu, J.H.; Wu, A.Z.; Luo, L.J. (2012).** Expression profile analysis of 9 heat shock protein genes throughout the life cycle and under abiotic stress in rice. *Chin. Sci. Bull.* 2012, 57, 336–343.
- Young TE, Ling J, Geisler-Lee CJ, Tanguay RL, Caldwell C, Gallie DR (2001)** Developmental and thermal regulation of the maize heat shock protein, HSP101. *Plant Physiology* 127, 777-791
- Zhang, C. and Guy, C.L. (2005)** Co-immunoprecipitation of Hsp101 with cytosolic Hsc70. *Plant Physiol. Biochem.* 43, 13–18.
- Zhang, J.-H., Wang, L.-J., Pan, Q.-H., Wang, Y.-Z., Zhan, J.-C., Huang, W.-D., (2008).** Accumulation and subcellular localization of heat shock proteins in young grape leaves during cross-adaptation to temperature stresses. *Scientia Horticulturae* 117, 231–240.
- Zhang, X., B. Wollenweber, D. Jiang, F. Liu & J. Zhao, (2008).** Water deficits and heat shock effects on photosynthesis of a transgenic *Arabidopsis thaliana* constitutively expressing ABP9, a bZIP transcription factor. *J Exp Bot.*, 59: 839-548.
- Zhu, C., Gore, M., Buckler, E.S., and Yu, J. (2008).** Status and prospects of association mapping in plants. *Plant Genome* 1: 5–20.
- Zou J, Liu C, Chen X. (2011)** Proteomics of rice in response to heat stress and advances in genetic engineering for heat tolerance in rice. *Plant Cell Reports* 30, 2155–2165.