

CHAPTER SEVEN

DISCUSSION AND CONCLUSIONS

CHAPTER SEVEN

DISCUSSION AND CONCLUSIONS

The mineral and petrographic characteristics, the diagenetic history, as well as the hydrocarbon reservoir quality of the Naqus Formation (Upper Cambrian) and the uppermost part of Araba Formation (Lower Cambrian) at Gebel Naqus, southwestern Sinai, together with some subsurface core samples are the main subjects treated in this work. A sedimentary section at Gebel Naqus to the north of El Tor was selected for this study. The section is about 370 m thick and is made up of the entire Naqus Formation underlain by the uppermost part of the underlying Araba Formation. One hundred and fourteen rock samples representing all the possible varieties were collected. Lithological variations and color changes of strata were taken into consideration during sampling. Observations were also focused on sedimentary textures, structures and facies changes.

Subsurface data from Ras Budran oil field were provided by Suez Oil Company. The field is located in the northern offshore area, 4 km west of the eastern coast of the Gulf of Suez. It has three main blocks A, B and C separated by two main faults. Six key wells were selected for this study: two wells from block A (RB-A1 and RB-A5); three wells from block B (RB-B3, RB-B4 and RB-B7) and one well from block C (RB-C2). Data provided included representative Paleozoic core samples (60 samples), core photographs, core analysis data and electric logs.

One hundred and fifty rock samples were selected to represent the studied sandstones. Petrographic and mineral examinations were considered using the conventional polarizing microscope, X-ray

diffraction (XRD) analysis, heavy mineral analysis, scanning electron microscopy (SEM) supplemented by X-ray energy dispersive analysis (EDAX). Some important petrophysical parameters were also measured.

Different types of sedimentary structures are recognized in the studied sandstones. They include: erosional surfaces, channel-fill structures, massive bedding, graded bedding, flat bedding and planar lamination, cross-bedding (tabular and trough) and cross-lamination, deformed structures (convolute bedding and recumbent foresets), chemical structures (differential coloration, lieegang bands, mottling, calcite and gypsum veinlets, and silica-filled fractures), mechanical structures (differential weathering, joints, open fractures) and biogenic structures (burrowing organisms, mainly *Skolithos* tubes). Deformed structures are widely recorded in Naqus Formation and its subsurface equivalents, whereas biogenic structures characterize Araba Formation and its subsurface equivalents.

Textural analysis indicates that the studied sandstones are very fine to medium-grained, occasionally coarse-grained, and poorly sorted to moderately well sorted, but the majority are moderately sorted. Grain shape varies from angular to well rounded, but the majority of the sand-size grains are subangular to subrounded.

Microscopic investigation of the studied sandstone samples showed that they are composed mainly of three architectural components; framework grains (F), cement (C) and porosity (P). Detrital matrix is very rare and constitutes a negligible proportion of the sandstone composition. Although infiltrated clay minerals are present in some samples, they are mostly of diagenetic origin. In this concern, Naqus Sandstone ranges in composition of the three components from $F_{57.50}C_{2.50}P_{0.0}$ to $F_{79.25}C_{42.50}P_{27.75}$ with an average of $F_{69.49}C_{12.94}P_{17.57}$. Sandstone of Araba

Formation ranges in composition of the three components from $F_{51.25}C_{19.25}P_{0.0}$ to $F_{75.75}C_{48.75}P_{11.50}$ with an average of $F_{62.29}C_{34.27}P_{3.45}$. On the other hand, the three components in the subsurface sandstone range from $F_{75.25}C_{2.00}P_{0.0}$ to $F_{88.25}C_{24.75}P_{14.75}$ with an average of $F_{82.03}C_{13.12}P_{4.86}$. Differences observed among the studied units, and in samples from the same unit, are primarily due to variations in diagenetic processes.

Because clay matrix is nearly absent, and quartz comprises the only framework mineral, the studied sandstones are considered as quartz arenites. Concerning mineral composition, they are mature to supermature. Texturally speaking, they are submature to mature as evidenced by the abundance of subrounded and rounded grains, and the moderate-to-high degree sorting. The scarcity of detrital clay minerals, rock fragments and feldspars may, however, be a compositional maturity obtained by dissolution during diagenesis (diagenetic maturation), thus, the studied sandstones may be described as diagenetic quartz arenites.

Quartz is the most abundant detrital component in the studied sandstones. The average values for quartz in the Naqus, Araba and subsurface rocks are 99.58, 99.42 and 99.60 %, respectively. Some of the studied samples have polymodal grain size distributions. Coarse sand grains are mixed together with very fine sand grains with some fragments in the intermediate size ranges. Such a texture may result from mixing of sediment from two different environments, storm mixing of material in a high-energy environment, multiple sources of sand supply or, in some cases, may be produced by burrowing or other *in situ* mixing processes. Banding on the scale of a thin section is locally observed.

Quartz grains are either monocrystalline or polycrystalline; the former are the most abundant; averaging 88.13, 97.02 and 96.34 % of the total quartz in the Naqus, Araba and subsurface sandstones, respectively.

Monocrystalline quartz with straight extinction (nonundulose) is the most common (averages 91.23, 96.58 and 95.31 %; respectively). The average values of undulose quartz are 8.77, 3.42 and 4.69 % in the Naqus, Araba and subsurface sandstones, respectively. The average ratios of the undulose to nonundulose types are 0.10, 0.04 and 0.05.

The average values of polycrystalline quartz in total quartz are 11.87, 2.98 and 3.66 % in the Naqus, Araba and subsurface sandstones; respectively. The average ratios of polycrystalline to monocrystalline types are 0.14, 0.03 and 0.04.

The complete spectrum of quartz types present in the studied sandstones is: common, volcanic, recrystallized metamorphic, stretched metamorphic and quartz vein. Petrographic observations indicate that common quartz, typical of plutonic and some metamorphic rocks, is by far the most abundant.

The predominance of quartz in the studied sandstones may indicate that parts of the source region had low relief and underwent deep weathering and the sediment is at least partly multi-cyclic or that transport distances were long. Furthermore, the assemblage of ultrastable heavy minerals indicates high degree of reworking during a multi-cyclic origin of these sands. Moreover, the presence of rounded detrital quartz grains, rounded sedimentary lithic fragments, and rounded grains of zircon and tourmaline suggest that a major component of the provenance is older (pre-existing) sedimentary rocks.

Feldspar minerals are essentially absent in the studied sandstones, except a few extensively altered to completely dissolved grains on the scale of a thin section. The scarcity of feldspar may have resulted from one or more of the following: 1) deposition in a high-energy regime which selectively destroyed feldspar grains; 2) poverty of source rock-

bearing feldspars; 3) low topography of the source area associated with the predominance of a warm, humid climate; and 4) extensive diagenetic modification due to intrastratal solution. In the present study, it is believed that the original feldspar minerals were lost due to grain dissolution by ground water in an active hydrologic regime and/or due to excessive alteration during burial diagenesis. Many feldspar grains were completely dissolved leaving behind skeletal remnants or showing no evidence of their former presence. In the later case, the dissolved grains left only oversized pores that indicate their original presence. The proportions of feldspars lost by the effect of these processes were determined assuming that 80 % of the oversized pores and of grains pseudomorphed by kaolinite were originally feldspars. The volume of feldspars lost averages 6.69, 4.64 and 0.77 % in the Naqus, Araba and subsurface sandstones, respectively.

Rock fragments are a minor constituent in the studied sandstones. Their average values in the Naqus, Araba and subsurface rocks are 0.22, 0.18 and 0.14 %; respectively. The recorded rock fragments are sedimentary and, much less commonly, metamorphic and volcanic.

Sedimentary rock fragments consist mainly of chert, fine-grained siliceous and ferruginous sandstones, siltstones and mudstones. Chert fragments are mostly made up of subrounded to well-rounded microcrystalline quartz suggestive of a polycyclic origin. They were generally mechanically and chemically stable during diagenesis. However, some chert fragments have undergone partial dissolution that resulted in microporosity development. Some chert fragments are embayed by quartz grains indicating a higher stability of quartz relative to chert. Silica-cemented siltstone and sandstone clasts are present locally. Hematite-cemented sandstone fragments are occasionally observed.

Metamorphic rock fragments are mainly represented by quartz mica schist and quartz gneiss. Different ductile grains, represented mainly by mica flakes are also recorded. These grains are relatively stable chemically, but they are easily deformed during compaction. Muscovite flakes are by far more common than biotite flakes, due to their abundance in source rocks and greater resistance to chemical weathering. Gypsum flakes are also recorded.

The petrographic data indicate that the source terrane or terranes for the studied sandstones include volcanic, plutonic, metamorphic, and sedimentary rocks. The quartz-rich character of the sandstones, together with the general scarcity of volcanic rock fragments, appears to rule out quartz-poor volcanic rocks as major source rocks. Neither plutonic nor metamorphic source rocks were major sediment contributors to most of the studied sandstones. Further, because much evidence of recycling was observed, quartz-rich sedimentary rocks are supposed to be the major source rocks. Analysis of quartz types further indicates that plutonic rocks were more important than metamorphic rocks as sources for the studied sandstones. However, scant evidences, based on relative rock-fragment abundance and polycrystalline quartz grains characteristics, suggest that metamorphic rocks may have been more important sediment sources than plutonic rocks. Volcanic rocks are the least.

Detrital clays are essentially absent in the studied sandstones except as a few claystone clasts. The average values of detrital clays in the Naqus, Araba and subsurface sandstones are 0.05, 0.4 and 0.23 %, respectively.

Heavy minerals may reach up to 1.45 % (average of 0.15 %) of the framework composition of Naqus Formation, whereas, they constitute a minor proportion of those of Araba Sandstone. On the other hand, heavy minerals in the subsurface sandstone may comprise up to 0.33 % with an average of 0.03 % of the framework composition.

Opaque minerals averages 54.18 and 52.49 % of the total heavies of Naqus and Araba sandstones, respectively. Opaques seem to be size-dependent, constituting the great proportion of the total heavy minerals (averaging 63.24 %) of the very fine sand fraction. This may be attributed to selective sorting during transport and deposition. In the studied sandstones, opaques are mainly represented by hematite, goethite, hydrogoethite and Fe-Ti oxides.

Non-opaque minerals averages 45.82 and 47.51 % of the total heavies of Naqus and Araba sandstones, respectively. Non-opaques also seem to be size-dependent showing percentages which are minimum (average 36.76 %) in the very fine sand fraction and maximum (average 49.01 %) in both the fine and medium sand fraction. Ten non-opaque, ultrastable (zircon, tourmaline and rutile) and metastable (garnet, sphene, epidote, staurolite, sillimanite, monazite and hornblende) heavy minerals were identified. Micas were excluded from the heavy mineral assemblage because of their extremely platy shape, which results in anomalous behavior during transport. The ultrastable minerals are the most abundant nonopaques in the studied sandstones. They were recorded in all the examined samples with a relatively high frequencies. These minerals can survive prolonged transport and were mostly recycled from older sedimentary rocks. The lack of other heavies may be attributed to their rarity in the source rocks and/or their relatively low stability during transport and burial.

Ternary diagrams of quartz, feldspars and rock fragments (QFR) illustrate average present-day composition of $Q_{99.78}F_{0.0}R_{0.22}$ for Naqus Sandstone, $Q_{99.82}F_{0.0}R_{0.18}$ for Araba Sandstone and $Q_{99.86}F_{0.0}R_{0.14}$ for the subsurface sandstone. The present composition of the studied sandstones is different- of course- from their composition at the time of deposition,

because these sandstones were progressively subjected to diagenetic processes and weathering. The restored average original composition is $Q_{89.54}F_{8.22}R_{2.24}$ for Naqus Sandstone, $Q_{91.21}F_{6.9}R_{1.89}$ for Araba Sandstone and $Q_{98.70}F_{0.93}R_{0.37}$ for the subsurface sandstone.

Cements and replacement minerals in the studied sandstones vary in abundance. They average 12.94, 34.27 and 13.12 % of the whole rock volume in the Naqus, Araba and subsurface sandstones, respectively. Authigenic cements include quartz, K-feldspars, clay minerals (mainly kaolinite), iron minerals (mostly hematite), calcite, gypsum and halite. Allogenic cement was also recorded and is represented mainly by infiltrated clays.

Quartz cement averages 1.33, 0.53 and 0.78 % of the whole rock volume in the Naqus, Araba and subsurface sandstones, respectively. It is represented mainly by syntaxial overgrowths. Microcrystalline quartz is another type that was recognized in few samples. It occurs as either mosaic cement or isopachous drusy coatings.

Authigenic K-feldspars constitute a negligible amount in the studied sandstones. Commonly they are found as coarse crystals cementing quartz grains. These crystals are not connected to detrital feldspar cores, but occur as isolated crystals partially filling pore spaces. Such feature is known to develop when the pore water becomes supersaturated with respect to K, Al and Si. Dissolution and alteration of authigenic K-feldspars in the studied sandstones have been locally recorded.

Allogenic clay minerals are present in minor proportions. They average 1.21, 1.80 and 4.42 % of the whole rock volume in the Naqus Araba and subsurface sandstones, respectively.

Authigenic clays in the studied sandstones are represented by kaolinite together with traces to minor proportions of illite, smectite, illite-smectite

mixed layer and chlorite as indicated by XRD and SEM examinations. They occur as pore-filling and pore-lining cements, and as pseudomorphous replacements. Kaolinite averages 7.73, 14.20 and 1.29 % of the whole rock volume in the Naqus, Araba and subsurface sandstones, respectively.

Iron minerals (mainly hematite) are abundant and ubiquitous cement in the studied sandstones. They vary largely in abundance among the studied sandstone units, and in samples from the same unit. They may comprise up to 6.50 % of the whole rock volume of Naqus Sandstone and up to 32.25 % of those of Araba Sandstone. The average values of iron minerals cement in the two formations are 1.55 and 17.22 %, respectively. On the other hand, iron minerals cement in the subsurface sandstone may reach 13.50 % with an average of 4.04 % of the whole rock volume. Some subsurface sandstone samples may have up to 70 % by volume of iron minerals cement (mainly hematite); these samples were not included in the point counting.

Calcite is the main carbonate mineral present as a cement in the studied sandstones. It averages 0.12, 0.10 and 0.17 % of the whole rock volume in the Naqus, Araba and subsurface rocks respectively.

The average values of halite in the Naqus, Araba and subsurface sandstones are 1.01, 0.52 and 2.3 %, respectively. However, halite is presently undergoing dissolution, and hence, its original extent is unknown.

Gypsum forms scattered patches of cement in few samples. Most patches are at the millimeter scale, but locally, they may be at the centimeter scale.

The detailed and comprehensive petrographic studies conducted on more than 150 thin sections and 100 samples examined by scanning

electron microscope showed that the studied sandstones exhibit a variety of diagenetic features that include: 1) mechanical infiltration of clay minerals, 2) dissolution of unstable detrital silicate grains (such as feldspars and ferromagnesian minerals) and the consequent development of secondary porosity, 3) development of an authigenic mineral suite that includes quartz, calcite, iron minerals (mostly hematite) and clays (mainly kaolinite), 4) compaction features such as closer packing, mechanical fracturing, deformation and bending of ductile and cleavable grains, 5) dissolution of early-formed carbonate cement followed by fracturing and secondary porosity initialization and 6) development of late stage cements made up of quartz, K-feldspars, kaolinite, iron minerals, calcite, gypsum and halite.

The paragenetic sequence of the diagenetic processes began shortly after deposition of the sediment by a limited infiltration of clays. A phase of precipitation of quartz cement occurred soon after deposition at shallow depths before the commence of real burial compaction. Cementation with iron minerals occurred early in the burial history and continued for a long time. However, recent studies revealed that iron minerals in sandstones may be introduced at any stage, viz, burial compaction, cementation or at any stage in diagenetic events, therefore the existence of iron minerals is valueless for any diagenetic sequence interpretation. The major part of cementation by calcite preceded significant mechanical compaction and followed quartz cementation. Dissolutions of feldspars and rock fragments probably occurred after the main episode of calcite cementation. Kaolinite, K-feldspars and late stage iron minerals cement were deposited following the dissolution process at late mesodiagenesis. Kaolinite commonly breaches secondary pore spaces and is mainly interpreted as a late diagenetic cement. Based on their

textural relationships with other cements and their fresh surfaces, halite and gypsum are interpreted to be formed very late in the paragenetic sequence.

In spite of their age and the large volumes of ground water that probably passed through them, the studied sandstones retain, with a few exceptions, sufficient porosity and permeability to possess a good reservoir quality. In the present study, porosity was measured petrophysically by using a helium porosimeter and in thin sections by point counting. Naqus Sandstone has helium porosity that averages 23.1 %. About 84 % of Naqus samples have porosity greater than 20 %. Sandstone of this porosity magnitude makes it as an excellent reservoir rock. In Araba Sandstone, helium porosity averages 17.8 %. This sandstone has a high kaolinite content and its porosity is usually intercrystalline. On the other hand, helium porosity in the subsurface sandstone averages 10.4 %.

Thin section porosity in Naqus Sandstone has a mean value of 17.7 %, while in Araba Sandstone it averages 3.6 %. On the other hand, thin section porosity in the subsurface sandstone averages 4.9 %. Of the total present thin section porosity, primary porosity averages 75.06, 54.75 and 95.70 %, and secondary porosity averages 24.94, 45.25 and 4.3 % in the Naqus, Araba and subsurface sandstones, respectively.

In Naqus Sandstone, permeability ranges from 1 md to 19526 md with an average of 2818 md. About 46 % of Naqus samples have permeability greater than 1000 md. Sandstone having this permeability magnitude makes an excellent reservoir characters. On the other hand, in Araba Sandstone, permeability ranges from 3 md to 162 md with an average of 50 md. 50 % of Araba samples have permeability values between 10 and 50 md, indicating a moderate reservoir quality. The subsurface sandstone

has a horizontal permeability ranging from 0.0 md to 1270 md and averaging 92 md. On the other hand, its vertical permeability varies from 0.0 md to 1340 md with an average of 100 md. This slight difference in permeability magnitude is attributed to the presence of small-scale open vertical fractures, which enhance vertical permeability. These fractures are sometimes only detectable in thin sections (hairline fractures). Sandstone having this permeability magnitude makes a good reservoir characters.

The reservoir quality of the studied sandstones was controlled mainly by compaction and secondarily by the degree of cementation (except in Araba Sandstone where cementation was more effective than compaction in reducing porosity). From an estimated initial porosity of 45 %, thin section porosity in Naqus, Araba and subsurface sandstones is reduced to the present value of 17.7, 3.6 and 4.9 %, respectively. The average Naqus, Araba and subsurface sandstones lost 31.69, 19.03 and 33.66 % porosity, respectively, by compaction and 5.45, 24.16 and 8.33 % porosity, respectively, by cementation.

Development of secondary porosity (about 4.39, 1.54 and 0.21 % total rock volume in Naqus, Araba and subsurface sandstones, respectively) locally enhanced the reservoir quality of the studied sandstones. This porosity resulted mainly from the dissolution of feldspars, rock fragments and calcite cement. Minor proportion of secondary porosity was also created by fracturing due to tectonic stresses or overpressure (hydrofracturing).

The sandstones of Naqus Formation are believed to have a non-marine mode of deposition. The numerous quartz pebbles that are either randomly distributed in Naqus Sandstone or preferentially oriented along cross-bedding planes may suggest a high-energy shallow agent of

transportation, probably a meandering fluvial system. The streams were probably very erratic and intermittent with coarse load at the bottom followed by the finer sediments with the diminishing of the flood. The proposed fluvial origin for Naqus Sandstone is strengthened by the presence of convolute bedding and recumbent foresets.

The presence of cross-bedding, fossil tracks of *Cruziana* and *Skolithos*, as well as traces of burrowing organisms in Araba Formation may support a shallow marine depositional environment. The *Skolithos* ichnofacies clearly indicates clean, well-sorted near-shore sands with high levels of wave and current energy (intertidal zone). The varicoloration of the sandstone beds could be the result of oxidation processes due to the wind activation of oxygen dissolved in the seawater or the alteration of iron bearing minerals such as mica and feldspars. Biotite and hornblende may have been altered by weathering to hematite and goethite, which give the red coloration. The dominant sedimentary structure in the varicolored sandstones is conventional horizontal bedding, which also suggests a quiet water agent. The observed small-scale cross-beds with horizontal planes in between could be due to an abrupt decrease in the current velocity and increase of the water depth.

REFERENCES

REFERENCES

- Abdallah, A. M., and Adindani, A., 1963**, Stratigraphy of Upper Paleozoic rocks, western side of the Gulf of Suez: Geological Survey of Egypt, 25, 18 p.
- Abdallah, A. M., Darwish, M., El-Araby, A., Gaupp, R., 1993**, Diagenesis related to unconformities in Late Carboniferous/Permian-Triassic sediments of Northern Galala, Gulf of Suez, Egypt: In: Thorweihe, U., and Schandelmeier, H., eds., Geoscientific Research in Northeast Africa: Balkema, Rotterdam, p. 345-349.
- Abdel-Hameed, A. T., Allam, A., and Khalil, H., 1986**, Sedimentological studies on the Pre-Cenomanian sandstones, north Gebel Qabeliat-east Gebel Abu Durba, southwest Sinai, Egypt: Egyptian Journal of Geology, v. 30/1-2, p. 25-45.
- Abdel-Wahab, A. A., 1985**, Sedimentological and diagenetic studies of the Nubia Formation in the Central Eastern Desert, Egypt: Unpublished Ph. D. thesis, Aston. University, B'ham, U. K., 507 p.
- Abdel-Wahab, A. A., 1990**, Fabric and compaction analysis of Araba and Naqus quartzarenites (Cambrian), Gebel Araba-Qabeliat, southwest Sinai, Egypt: Delta Journal of Science, v.14/2, p. 666-707.
- Abdel-Wahab, A. A., 1999**, Petrography, fabric analysis and diagenetic history of Upper Cretaceous sandstone, Kharga Oasis, Western Desert, Egypt: Sediment. Egypt, v. 7, p. 99-117.
- Abdel-Wahab, A. A., and Abu El-Maati, M. A., 1989**, Petrography and diagenesis of the Nubia Formation at Safaga Qena District, Central Eastern Desert, Egypt: 1st Conf. Geochem., Alexandria University (abs.), p. 456.

- Abdel-Wahab, A. A., and McBride, E. F., 1991**, Diagenetic control of reservoir quality of Araba and Naqus diagenetic quartzarenites (Cambrian), Gebel Araba-Qabeliat, southwest Sinai, Egypt: *Delta Journal of Science*, v.15, p. 160-203.
- Abdel-Wahab, A. A., and Turner, P., 1991**, Diagenesis of the Nubia Formation, central eastern desert, Egypt: *Journal of African Earth Sciences*, v. 13/3/4, p. 343-358.
- Abdel-Wahab, A. A., Allam, A., Kholief, M., and Salem, A., 1988**, Provenance of Gebel El-Zeit sandstones, Gulf of Suez, Egypt: 26th Annual Meeting, Geological Society of Egypt, Cairo (abs.), 12-16 Nov., p. 30.
- Abdel-Wahab, A. A., Allam, A., Kholief, M., and Salem, A., 1992**, Sedimentological and palaeoenvironmental studies on the clastic sequence of Gebel El-Zeit area, Gulf of Suez, Egypt: *Journal of African Earth Sciences*, v. 14/1, p. 121-129.
- Abdel-Wahab, A. A., Salem, A. M. K., and McBride, E. F., 1998**, Quartz cement of meteoric origin in silcrete and nonsilcrete sandstone, Lower Carboniferous, western Sinai, Egypt: *Journal of African Earth Sciences*, v. 27/2, p. 277-290.
- Abdelwahab, H. Sh., 1972**, The geology and mineralogy of Nubian Sandstones from Wadi Natash and Kom Omb districts: Ph. D. thesis, Faculty of Science, Ain Shams University, Cairo, Egypt.
- Abdelwahab, H. Sh., 1998**, Petrography, mineralogy and geochemistry of sandstones of Gebel Gunna, central Sinai, Egypt: *Sedimentology of Egypt*, v. 6, p. 33-50.

- Ahmed, M. S., and Osman, R. A., 1998,** A field appraisal of sedimentary facies and environment of the Pre-Carboniferous, SW Sinai, Egypt: *Sedimentology of Egypt*, v. 6, p. 21-32.
- Ali, R. and Khairy, A., 1996,** Mechanism of Mid-Clysmic event and its effect on the Pre and Post Miocene structure of Ras Budran field, Gulf of Suez, Egypt: *Proceedings of Thirteenth Exploration and Production Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, 1996, Published by Egyptian General Petroleum Corporation, Exploration Part*, v. 2/2, p. 396-409.
- Allam, A., 1989,** The Paleozoic sandstones in Wadi Feiran–El Tor area, Sinai, Egypt: *Journal of African Earth Sciences*, v. 9/1, p. 49-57.
- Allam, A., and Khalil, H., 1987,** Geological and stratigraphical study on Gebel Qabeliat area, southwest Sinai, Egypt: *Fourth Sym. Phan. & Develop., Nat. Com. Geol. Sc. (IGCP), Cairo 22 April*.
- Allen, J. L. R., and Banks, N. L., 1972,** An interpretation and analysis of recumbent-folded deformed cross-bedding: *Sedimentology*, v. 19, p. 257-283.
- Almon, W. R., and Davies, D. K., 1978,** Clay technology and well stimulation: In : *Abstracts Transactions Gulf Coast Association Geological Societies, V. 28, 1 p.:* Gulf Coast Association of Geological Studies, New Orleans, USA.
- Andel, T. H. van, 1959,** Reflections on the interpretation of heavy mineral analyses: *Journal of Sedimentary Petrology*, v. 29, p. 153-163.
- API-RP-27,** Recommended practice for determining permeability of porous media: American Petroleum Institute, Dalls.
- API-RP-40,** Recommended practice for core analysis procedures: American Petroleum Institute, Dallas.

- Atkins, J. E., 1989**, Porosity and packing of Holocene river, dune and beach sands: Unpublished M.S. thesis, University of Texas, Austin, Texas, 224 p.
- Atkins, J. E., and McBride, E.F., 1992**, Porosity and packing of Holocene river, dune and beach sands: The American Association of Petroleum Geologists Bulletin, v. 76/3, p. 339-355.
- Awad, G. H., and Said, R., 1965**, Lexique stratigraphique international, Afrique (Egypte): Congr. Geol. Intern. 4, Fasc. 4b, 1-73.
- Bailey, E. H., and Stevens, R. E., 1960**, Selective staining of k-feldspar and plagioclase on rock slabs and thin section: American Mineralogist, v. 45, p. 1020-1025.
- Ball, J., 1916**, The geography and geology of west central Sinai: Egyptian Survey Department, Cairo, 219 p.
- Barakat, M. G., Darwish, M., and El-Barkooky, A. N., 1986**, Litostratigraphy of the Post Carboniferous-Pre Cenomanian clastics in west central Sinai and Gulf of Suez, Egypt: Proceedings of Eighth Exploration Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, November 17-23.
- Barron, T., 1907**, The topography and geology of the Peninsula of Sinai (western portion): Egyptian Survey Department, Cairo, 241 p.
- Basu, A., 1985**, Influence of climate and relief on composition of sands released at source areas: In: Zuffa, G. G., ed., Provenance of Arenites, Reidel Publication, Dordrecht, Holland, p. 1-18.
- Basu, A., Young, S. W., Suttner, L. J., James, W. C., and Mack, G. H., 1975**, Re-evaluation of the use of undulatory extinction and polycrystallinity in detrital quartz for provenance interpretation: Journal of Sedimentary Petrology, v. 45/4, p. 873-882.

- Beadnell, H. J. L., 1927**, The wilderness of Sinai: Arnold, London, 180 p.
- Beard, D. C., and Weyl, P. K., 1973**, The influence of texture on porosity and permeability of unconsolidated sand: The American Association of Petroleum Geologists Bulletin, v. 57, p. 349-369.
- Beets, C. B., 1948**, Correlation of the Paleozoic and Mesozoic rocks in Egypt: Anglo-Egyptian Oilfields Company, Internal Report No. 679 (Files of The Egyptian General Petroleum Corporation, Cairo), 33 p.
- Beleity, A. M., Ghoneim, M., Hinawi, M., Fathi, M., Gebali, H., and Kamel, M., 1986**, Paleozoic stratigraphy, paleogeography and paleotectonics in the Gulf of Suez, Egypt: Proceedings of Eighth Exploration Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, November 17-23, 1986, Dar Al-Madina Al-Monawara Press, 1988, v. 1, p. 96-116.
- Bell, P. H., and Etheridge, M. A., 1973**, Microstructures of mylonites and their descriptive terminology: Lithos, v. 6, p. 337-348.
- Berner, R. A., 1969**, Goethite stability and the origin of red beds: Geochim. Cosmochim. Acta, v. 33, p. 267-273.
- Bhattacharyya, D. P., and Dunn, L. G., 1986**, Sedimentologic evidence for repeated pre-Cenozoic vertical movements along the northeast margin of the Nubian Craton: Journal of African Earth Sciences, v. 5/2, p. 147-153.
- Birkeland, P. W., 1984**, Soils and geomorphology: Oxford University Press, New York.
- Bjørlykke, K., 1983**, Diagenetic reactions in sandstones: In: Parker, A., and Sellwood, B. W., eds., Sediment Diagenesis: Reidel Publication, Dordrecht, Holland, p. 169-213.

- Bjørlykke, K., 1984**, Formation of secondary porosity: How important is it?: In: McDonald, D. A., and Surdam, R. C., eds., *Clastic Diagenesis: The American Association of Petroleum Geologists, Memoir No. 37*, p.277-286.
- Blatt, H., 1967a**, Original characteristics of clastic quartz grains: *Journal of Sedimentary Petrology*, v. 37/2, p. 401-424.
- Blatt, H., 1979**, Diagenetic processes in sandstones: In: Scholle, P. A., and Schluger, P. R., eds., *Aspects of Diagenesis: Society of Economic Palaeontologist and Mineralogist, Special Publication No. 26*, p. 141-157.
- Blatt, H., 1992**, *Sedimentary petrology: 2nd edition*, New York, 514 p.
- Blatt, H., and Christie, J. M., 1963**, Undulatory extinction in quartz of igneous and metamorphic rocks and its significance in provenance studies of sedimentary rocks: *Journal of Sedimentary Petrology*, v. 33/3, p. 559-579.
- Blatt, H., Middleton, G. V., and Murray, R. C., 1980**, *Origin of sedimentary rocks: 2nd edition*, Prentice-Hall, Englewood Cliffs, New Jersey, 782 p.
- Bloch, S., and Helmold, L., 1995**, Approaches to predicting reservoir quality in sandstones: *American Association of Petroleum Geologists Bulletin*, v. 79/1, p. 97-115.
- Blodgett, R. H., Crabaugh, J. P., and McBride, E. F., 1993**, The color of red beds: A geologic perspective: *Soil Sci. Soc. Amer.*, Special Publication No. 31, p. 127-159.
- Boles, J. R., and Franks, S. G, (1979)** Clay diagenesis in Wilcox sandstones of southwest Texas: Implications of smectite diagenesis on

sandstone cementation: *Journal of Sedimentary Petrology*, v. 49, 55-70.

Brenchley, P. J., Newall, G., and Stanistreet, I. G., 1979, A storm surge origin for sandstone beds in an epicontinental platform sequence, Ordovician, Norway: *Sedimentary Geology*, v. 22, p. 185-217.

Brenckle, P., L., and Marchant, T. R., 1982, Calcareous microfossils from the Um Bogma Formation (Lower Carboniferous) of the Lower Nubia Group at Gebel Nukhul, Sinai, Egypt: *Proceedings of Sixth Exploration Seminar sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt*, p. 302-309.

Brown, R. N., 1980, History of exploration and discovery of Morgan, Ramadan and July oil fields, Gulf of Suez, Egypt: In : Miall, A. M., ed., *Facts and principles of world petroleum occurrence*, Canadian Society of Petroleum Geologists, Memoir No. 6, p. 733-764.

Burley, S. D., and Kantorowicz, J. D., 1986, Thin section and SEM. textural criteria for the recognition of cement-dissolution porosity in sandstones: *Sedimentology*, v.33, p. 587-604.

Busch, J. M., Fortney, W. G., and Berry, L. N., 1987, Determination of lithology from well-logs by statistical analysis: *SPE Formation Evaluation*, p. 412-418.

Cade, C. A., Evans, I. J., and Bryant, S. L., 1994, Analysis of permeability controls: A new approach: *Clay Minerals*, v. 29, p. 491-501.

Callender, D. L., and Folk, R. L., 1958, Idiomorphic zircon: Key to volcanism, the Lower Tertiary sands of central Texas: *American Journal of Science*, v. 256, p. 257-269.

- Carver, R. E., ed., 1971, Procedures in sedimentary petrology: New York, Wiley-Interscience, 672 p.**
- Chowdhary, L. R., and Taha, S., 1986, History of exploration and geology of Ras Budran, Ras Fanar, and Zeit Bay oil fields, Gulf of Suez, Egypt: Proceedings of Eighth Exploration Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, November 17-23, 1986, Dar Al-Madina Al-Monawara Press, 1988, v. 1, p. 149-182.**
- Chowdhary, L. R., Shaheen, S., and Naggar, A. A., 1986, Structure and structural evolution of Ras Budran, Gulf of Suez, Egypt: Proceedings of Eighth Exploration Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, November 17-23, 1986, Dar Al-Madina Al-Monawara Press, 1988, v. 1, p. 308-321.**
- Clarke, R. H., 1979, Reservoir properties of conglomerates and conglomeratic sandstones: The American Association of Petroleum Geologists Bulletin, v. 63, p. 799-809.**
- Colletta, B., Le Quellec, P., Letouzy, J., and Moretti, I., 1988, Longitudinal evolution of the Suez rift structure, Egypt: Tectonophysics, v. 153, p. 221-233.**
- Collinson, J. D., and Thompson, D. B., 1982, Sedimentary structures: George Allen and Unwin (Publishers) Ltd, London, UK, 193 p.**
- Crone, A. J., 1975, Laboratory and field studies of mechanically infiltrated matrix clay in arid fluvial sediments: Unpublished Ph .D thesis, University of Colorado, USA, 162 p.**
- Curtis, C. D., Hughes, C. R., Whiteman, J. A., and Whittle, C. K., 1985, Compositional variation within some sedimentary chlorites and some comments on their origin: Mineral, Mag., v. 49, p. 375-386.**

- De Boer, R. D., 1977a**, On the thermodynamics of pressure solution interaction between chemical and mechanical forces: *Geochim. Cosmochim. Acta*, v. 41, p. 249-256.
- De Boer, R. D., 1977b**, Pressure solution: Theory and experiments: *Tectonophysics*, v. 39, p. 287-301.
- Deer, W. A., Howie, R. A. and Zussman, J., 1992**, *Rock-Forming Minerals*: Longman Group Ltd., London, 2nd ed., 696 p.
- Dickinson, W. R., 1988**, Provenance and sediment dispersal in relation to paleotectonics and paleogeography of sedimentary basins: In: Kleinspehn, K. L., and Paola, C., eds., *New Perspectives in Basin Analysis*: Springer-Verlag, New York, p. 3-25.
- Dietz, V., 1973**, Experiments on the influence of transport on shape and roundness of heavy minerals: *Contr. Sed.*, v.1, p. 69-102.
- Dott, R. H., 1964**, Wacke, graywacke and matrix - what approach to immature sandstone classification? *Journal of Sedimentary Petrology*, v. 34, p. 625-632.
- Dutton, S. P., 1986**: Diagenesis and burial history of the Lower Cretaceous Travis Peak Formation, East Texas: Unpublished Ph. D. thesis, University of Texas, Austin, Texas, 165 p.
- Dutton, S. P., and Land, L. S., 1988**, Diagenetic history of a well-cemented quartzarenite, Lower Cretaceous Travis Peak Formation, East Texas: *Geological Society of America Bulletin*, v. 100, p. 1271-1282.
- Eames, L. E., 1984**, Palynologic definition of Paleozoic unconformity bounded sequences, Gulf of Suez region, Egypt: *Proceedings of Sevenths Exploration and Production Conference sponsored by The Egyptian General Petroleum Corporation*, Cairo, v. 1, p. 117-125.

- Ehrenberg, S. N., 1989**, Assessing the relative importance of compaction processes and cementation to reduction of porosity in sandstones: Discussion; compaction and porosity evolution of Pliocene sandstones, Ventura Basin, California: Discussion: The American Association of Petroleum Geologists Bulletin, v. 73/10, p. 1274-1276.
- Ehrenberg, S. N., 1990**, Relationship between diagenesis and reservoir quality in sandstones of the Garn Formation, Haltenbanken, mid-Norwegian continental shelf: The American Association of Petroleum Geologists Bulletin, v. 74, p. 1538-1558.
- El-Hinnawi, E. E., Kabesh, M. L., and Zahran, I., 1973**, Mineralogy and chemistry of Nubian sandstone from the central Eastern Desert of Egypt: N. Jb. Min. Abh., v. 118, p. 211-234.
- El-Mansey, I. M., Hassanien, I. M., and Kholeif, M. M., 1988**, Characteristic of primary and secondary structures of some Paleozoic sandstones in southwest Sinai, Egypt: Proceedings of the Egyptian Academy of Science, v. 37, p. 111-125.
- El-Shahat, A., and Kora, M., 1986**, Petrology of the Early Paleozoic rocks of Um Bogma area, Sinai: Mansoura Science Bulletin, v. 13, p. 151-184.
- Ernst, W. G., and Blatt, H., 1964**, Experimental study of quartz overgrowths and synthetic quartzites: Journal of Geology, v. 72, p. 461-470.
- Eyal, M., Bartov, Y., Shimron, A. E., and Bentor, Y. K., 1980**, Geological Map of Sinai, scale 1: 500 000.
- Fisher, R. S., 1982**, Diagenetic history of Eocene Wilcox sandstones and associated formation waters, south-central Texas: Unpublished Ph. D. thesis, University of Texas, Austin, Texas, 185 p.

- Folk, R. L., 1980**, Petrology of sedimentary rocks: Hemphill Publishing Company, Austin, Texas, 3rd ed., 182 p.
- Folk, R. L., and Ward, W. C., 1957**, Brazos river bar: A study in the significance of grain size parameters: *Journal of Sedimentary Petrology*, v. 27/1, p. 3-26.
- Force, E. R., 1980**, The provenance of rutile: *Journal of Sedimentary Petrology*, v. 50/2, p. 485-488.
- Fothergill, C. A., 1955**, The cementation of oil reservoir sands and its origin: *Proceedings of Fourth World Petroleum Congress, Section 1B, Paper 1*, p. 301-314.
- Fraser, H. J., 1935**, Experimental study of the porosity and permeability of clastic sediments: *Journal of Geology*, v. 43, p. 910-1010.
- Friedman, G., M., 1971**, Staining, in: Carver, R. E., ed., *Procedures in sedimentary petrology*, New York, John Wiley and Sons, p. 511-530.
- Füchtbauer, H., 1967**, Influence of different types of diagenesis on sandstone porosity: *Proceedings of Seventh World Petroleum Congress, Mexico*, v. 2, p. 353-369.
- Füchtbauer, H., 1974a**, *Sediments and Sedimentary Rocks: Schweitzerbart*, Stuttgart, 2nd edition, 464 p.
- Gass, I. G., 1981**, Pan-African (Upper Proterozoic) plate tectonics of the Arabian Nubian shield: In: Kroener, A., ed., *Precambrian plate tectonics*, Elsevier, p. 387-405.
- Ghanem, M. F., and El-Mansey, I. M., 1991**, Lithostratigraphy and petrography of the Paleozoic sequences in west Sinai and Gulf of Suez, Egypt: *Egyptian Journal of Geology*, v. 35/1-2, p. 75-92.
- Girard, J. P., Savin, S. M., and Aronson, J. L., 1989**, Diagenesis of the Lower Cretaceous Arkoses of the Angola Margin: *Petrologic, K/Ar*

- dating and $^{18}\text{O}/^{16}\text{O}$ evidence: *Journal of Sedimentary Petrology*, v. 59, p. 519-538.
- Goudie, A. S., ed., 1985**, The encyclopedia dictionary of physical geography: Blackwell Reference, Oxford.
- Grim, R. E., 1953**, Clay Mineralogy: McGraw-Hill, New York, 384 p.
- Grim, W. D., 1973**, Stepwise heavy mineral weathering in the residual quartz gravels, Bavarian Molasse (Germany): *Contrib. Sedimentol.*, v. 1, p. 103-125.
- Harms, J. C., 1975**, Stratification and sequence in prograding shoreline deposits: Society of Economic Palaeontologist and Mineralogist, Short Course No. 2, p. 81-102.
- Harper, T. R., 1984**, Recommendations for improvement of well productivity Unit I, Ras Budran reservoir, Egypt: For Suco by British Petroleum.
- Hassan, A. A., 1967**, A new Carboniferous occurrence in Abu-Durba, Sinai, Egypt: Proceedings of Sixth Arab Petroleum Congress, Baghdad, 2, 39 (B3).
- Hawkins, P. J., 1978**, Relationship between diagenesis porosity reduction and oil emplacement in late Carboniferous sandstone reservoirs, Bothamsall Oil field, E. Midlands: *Journal of Geological Society of London*, v. 135, p. 7-24.
- Hayes, M. J., and Boles, J. R., 1992**, Volumetric relations between dissolved plagioclase and kaolinite in sandstones: Implications for aluminum mass transfer in the San Joachim Basin, California: In: Houseknecht, D. W., and Pittman, E. D., eds., *Origin, Diagenesis and Petrophysics of Clay Minerals in Sandstones*: Society of Economic

- Palaeontologist and Mineralogist, Special Publication No. 47, p. 111-124, Tulsa.
- Heald, M. T., 1950**, Authigenesis in West Virginia sandstones: *Journal of Geology*, v. 58, p. 624-633.
- Heald, M. T., 1955**, Stylolites in sandstones: *Journal of Geology*, v. 63, p. 101-114.
- Heald, M. T., 1956**, Cementation of Simpson and St. Peter sandstones in parts of Oklahoma, Arkansas and Missouri: *Journal of Geology*, v. 64, p. 16-30.
- Heald, M. T., and Larese, R. E., 1974**, Influence of coatings on quartz cementation: *Journal of Sedimentary Petrology*, v. 44, p. 1269-1274.
- Heckel, P. H., 1972**, Recognition of ancient shallow marine environments: In: Rigby, J. K., and Hamlin, W. K., eds., Recognition of ancient sedimentary environments: Society of Economic Palaeontologist and Mineralogist, Special Publication No. 16, p. 226-286, Tulsa.
- Houseknecht, D. W., 1987**, Assessing the relative importance of compaction processes and cementation to reduction of porosity in sandstones: *The American Association of Petroleum Geologists Bulletin*, v. 71/6, p. 633-642.
- Houseknecht, D. W., 1988**, Intergranular pressure solution in four quartzose sandstones: *Journal of Sedimentary Petrology*, v. 58/2, p. 228-246.
- Howard, J., 1992**, Influence of authigenic clay minerals on permeability: Society of Economic Palaeontologist and Mineralogist, Special Publication No. 47, p. 257-264, Tulsa.

- Hunter, R. E., and Clifton, H. E., 1982,** Cyclic deposits and hummocky cross-strata of probable storm origin in the Upper Cretaceous rocks of the Cape Sebastian area southwestern Oregon: *Journal of Sedimentary Petrology*, v. 49, p. 127-143.
- Hurst, A., and Nadeau, P. H., 1995,** Clay microporosity in reservoir sandstones: An application of quantitative electron microscopy in petrophysical evaluation: *The American Association of Petroleum Geologists Bulletin*, v. 79/4, p. 563-573.
- Ibrahim, A. E., 1996,** Petroleum geology of the Cretaceous and Pre-Cretaceous sediments in some oil fields in the Gulf of Suez area, Egypt: Ph.D. thesis, Cairo University, Cairo, Egypt, 211 p.
- Issawi, B., 1973,** Nubia sandstone: Type section: *The American Association of Petroleum Geologists Bulletin*, v. 57/4, p. 741-745.
- Issawi, B., and Jux, U., 1982,** Contributions to the stratigraphy of the Paleozoic rocks in Egypt: *Geological Survey of Egypt*, 64, 28 p.
- Issawi, B., El Hinnawi M., El Khawaga, L., Labib, S., and Anani, N., 1981,** Contributions to the geology of Wadi Feiran area, Sinai, Egypt: *Geological Survey of Egypt, Internal Belayim Exploration report*, 48 p.
- James, W. C., 1985,** Early diagenesis, Atherton Formation (Quaternary): A guide for understanding early cement distribution and grain modifications in non-marine deposits: *Journal of Sedimentary Petrology*, v. 55, p. 135-146.
- Jenkins, D. A., 1990,** North and Central Sinai: In: Said, R., ed., *The Geology of Egypt*, Chapter 19, A. A. Balkema, Rotterdam, 734 p.
- Jensen, J. L., Hinkley, D. V., and Lake, L. W., 1987,** A statistical study of reservoir permeability: distributions, correlations and averages: *SPE Formation Evaluation*, p. 461-468.

- Kantorowicz, J. D., 1985,** The petrology and diagenesis of Middle Jurassic clastic sediments, Ravenscar Group, Yorkshire: *Sedimentology*, v. 32: p. 833-853.
- Keller, W. D., 1969,** Chemistry in introductory geology, Bros. Pub., Columbia, Missouri, 108 p.
- Kessler, H., L. G., 1978,** Diagenetic sequence in ancient sandstones deposits deposited under desert climatic conditions: *Journal of Geological Society of London*, v. 135, p. 41-49.
- Ketzer, J.M., Morad, S., Nystuen, J.P., and De Ros, L. F., 2003,** The role of the Cimmerian unconformity (Early Cretaceous) in the kaolinitization and related reservoir-quality evolution in Triassic sandstones of the Snorre Field, North Sea: In: Worden, R. H., and Morad, S., eds., *Clay Mineral Cements in Sandstones*, International Association of Sedimentologists, Special Publication No. 34, p. 361-382, Blackwell Publishers, Oxford.
- Khairy, A., and Swidan, N., 1992,** The geological evolution of the development of Ras Budran field: Proceedings of Eleventh Exploration and Production Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, 1992.
- Klitzsch, E., 1990,** Paleozoic: In: Said, R., ed., *The Geology of Egypt*, Chapter 21, A. A. Balkema, Rotterdam, 734 p.
- Kora, M., 1984,** The Paleozoic outcrops of Um-Bogma area, Sinai, Egypt: Ph.D. thesis, University of Mansoura, Mansoura, Egypt, 235 p.
- Kora, M., 1989a,** Biostratigraphy of the Late Paleozoic succession in Gebel Ekma, southwestern Sinai, Egypt: *Neues Jahrbuch Geologie Palaontologie Monatshefte*, Stuttgart, v 5, p. 293-307.

- Kostandi, A. B., 1959**, Facies maps for the study of Paleozoic and Mesozoic sedimentary basins of the Egyptian region: Proceedings of First Arab Petroleum Congress, Cairo, 2, p. 54-62.
- Krauskopf, K. B., 1979**, Introduction to geochemistry: McGraw-Hill, Kogakusha Ltd., New York, 617 p.
- Krumbein, W. C., and Monk, G. D., 1942**, Permeability as a function of the size parameters of unconsolidated sands: Am. Inst. Min. and Met. Eng., Technical Publication No. 1492, p. 1-11.
- Krynine, P. D., 1946**, The Tourmaline group in sediments: Journal of Geology, v. 54, p. 65-87.
- Lamplugh, G. W., 1902**, Calcrete: Geol. Mag., v. 9, p. 575.
- Land, L. S., and Dutton, S. P., 1978**, Cementation of a Pennsylvanian deltaic sandstone: Isotopic data: Journal of Sedimentary Petrology, v. 48, p. 1167-1176.
- Land, L. S., Milliken, K. L., and McBride, E. F., 1987**, Diagenetic evolution of Cenozoic sandstones, Gulf of Mexico sedimentary Basin: Sedimentary Geology, v. 50, p. 195-225.
- Langford-Smith, T., ed., 1978**, Silcrete in Australia: Armidale, University of New England, 304 p.
- Langmuir, D., 1971**, particle size effect on the reaction goethite = hematite + water: American Journal of Science, v. 271, p. 147-156.
- Lee, C. H., 1919**, Geology and ground waters of the western part of the San Diego County, California, Washington: Water Supply Invig. Papers, 446-121 p.
- Levorsen, A. I., 1967**, Geology of petroleum: Freeman, San Francisco, 2nd ed., 724 P.

- Loucks, R. G., Dodge, M. M., and Galloway, W. E., 1984,** Regional controls on diagenesis and reservoir quality in Lower Tertiary sandstones along the Texas Gulf Coast: In: McDonald, D. A., and Surdam, R. C., eds., *Clastic Diagenesis: The American Association of Petroleum Geologists, Memoir No. 37*, p.15-45.
- Lundegard, P. D., 1992,** Sandstone porosity loss-a "big picture" view of the importance of compaction: *Journal of Sedimentary Petrology*, v. 62/2, p. 250-260.
- Mackenzie, W. S. and Guilford, G., 1986,** *Atlas of Rock-Forming Minerals in thin section: Longman House, Burnt Mill, Harlow, England, 98 p.*
- Magwood, J. P., and Pemberton, S. G., 1990,** Stratigraphic significance of Cruziana: New data concerning The Cambrian-Ordovician ichnostratigraphic paradigm: *Geology*, v.18, p. 729-732.
- Mange, M. A., and Maurer, H. F. W., 1992,** *Heavy minerals in color: Chapman and Hall, London.*
- Martinsson, A., 1965,** Aspects of a Middle Cambrian Thanatotope in Oland: *Geol. För. Stockh. Förh.*, v. 87, p. 181-230.
- Matlack, K. S., Houseknecht, D. W., and Applin, K. R., 1989,** Emplacement of clay into sand by infiltration: *Journal of Sedimentary Petrology*, v. 59, p. 77-87.
- Mason, C. C., and Folk, R. L., 1958,** Differentiation of beach, dune and aeolian flat environments by size analysis, Mustang Island, Texas: *Journal of Sedimentary Petrology*, v. 28/2, p. 211-226.
- McBride, E. F., 1984,** Rules of sandstone diagenesis related to reservoir quality (expanded abs.): *Gulf Coast Assoc. Geol. Soc. Trans.*, v. 34, p. 137-139.

- McBride, E. F., 1985**, Diagenetic processes that affect provenance determinations in sandstone: In: Zuffa , G. G., ed., Provenance of Arenites: Reidel Publication, Dordrecht, Holland, p. 95-113.
- McBride, E. F., 1987**, Diagenesis of the Maxon Sandstone (Early Cretaceous), Marathon region, Texas: A diagenetic quartzarenite: *Journal of Sedimentary Petrology*, v. 57, p. 98-107.
- McBride, E. F., 1989**, Quartz cement in sandstones: A review: *Earth-Science Reviews*, v. 26, p. 69-112.
- McBride, E. F., Diggs, T. N., and Wilson, J. C., 1991**, Compaction of Wilcox and Carrizo sandstones (Paleocene-Eocene) to 4420 m, Texas Gulf Coast: *Journal of Sedimentary Petrology*, v. 61. p. 73-85.
- McBride, E. F., Abdel-Wahab, A. A., and Salem, A. M. K. S., 1996**, The influence of diagenesis on the reservoir quality of Cambrian and Carboniferous sandstones, southwest Sinai, Egypt: *Journal of African Earth Sciences*, v. 22/3, p. 285-300.
- McKee, E. D., 1962**, Origin of the Nubian and similar sandstones: *Geol. Rdsch.*, v 52/2, SS 1-586.
- Meshref, W. M., 1990**, Tectonic framework: In: Said, R., ed., *The Geology of Egypt*, Chapter 8, A. A. Balkema, Rotterdam, 734 p.
- Milliken, K. L., 1988**, Loss of provenance information through subsurface diagenesis in Plio-Pleistocene sandstones, northern Gulf of Mexico: *Journal of Sedimentary Petrology*, v. 58, p. 992-1002.
- Milliken, K. L., and Mack, L. E., 1990**, Subsurface dissolution of heavy minerals, Frio Formation sandstones of the ancestral Rio Grand Province, South Texas: *Sedimentary Geology*, v. 68, p. 187-199.
- Milliken, K. L., McBride, E. F., and Land, L. S., 1989**, Numerical assessment of dissolution versus replacement in the subsurface

- destruction of detrital feldspar, Oligocene Frio Formation, South Texas: *Journal of Sedimentary Petrology*, v. 59, p. 740-757.
- Millot, G., 1960**, Silice, silex, silicifications et croissance des cristaux: *Bull. Serv. Carte Geol. Als. Lorr.*, v. 13, p. 129-146.
- Mills, P. C., 1983**, Genesis and diagnostic value of soft sediment deformation structures: A review: *Sedimentary Geology*, v. 35, p. 83-104.
- Milner, H. B., 1962a**, *Sedimentary Petrography: Volume I, Methods in sedimentary petrography*, 4th ed., London, George Allen and Unwin Ltd., 643 p.
- Milner, H. B., 1962b**, *Sedimentary petrography: Volume II, Principles and applications*, 4th ed., London, George Allen and Unwin Ltd., 715 p.
- Milnes, A. R., Bourman, R. P., and Fitzpatric, R. W., 1987**, Petrology and mineralogy of laterites in southern and eastern Australia and southern Africa: *Chemical Geology*, v. 60, p. 237-250.
- Molenaar, N., 1986**, The interrelation between clay infiltration, quartz cementation and compaction in Lower Givetian terrestrial sandstones, northern Ardennes, Belgium: *Journal of Sedimentary Petrology*, v. 56, p. 359-369.
- Morton, A. C., 1984**, Stability of detrital heavy minerals in Tertiary sandstones from the North Sea Basin: *Clay Mineralogy*, v. 19, p. 287-308.
- Morton, A. C., 1987**, Influences of provenance and diagenesis on detrital garnet suites in the Paleocene Forties, central North Sea: *Journal of Sedimentary Petrology*, v. 57, p. 1027-1032.

- Moustafa, A. M., 1976**, Block faulting in the Gulf of Suez: Proceedings of Fifth Exploration Conference sponsored by The Egyptian General Petroleum Corporation, Cairo, 19 p.
- Naggar, A. A., and El Hilaly, H., 1985**, Ras Budran field: Geological aspects and reservoir units: Proceedings of Eighth Production Seminar sponsored by The Egyptian General Petroleum Corporation, Cairo, Egypt, March 26-28, 1985.
- Nahon, D. B., 1991**, Evolution of iron crusts in tropical landscapes, In: Coleman, S. M., and Dethier, D. P., eds., Rates of chemical weathering: Academic Press, San Diego, California, p. 169-197.
- Nassr, M. N., 1985**, Contrasts between quartz grain shapes of the Cambrian and Carboniferous "Nubia" facies in Gabal Abu-Durba, Sinai, Egypt: Fourier grain shape analysis: Ph.D. thesis, University of South Carolina, USA, 394 p.
- Neasham, J. W., 1977**, Applications of scanning electron microscopy to the characterization of hydrocarbon bearing rocks: Scanning Electron Microscopy, v. 1, p. 1-8.
- Nelson, C. H., 1982**, Modern shallow water graded sand layers from storm surges, Bering shelf: A mimic of Bouma sequences and turbidite systems: Journal of Sedimentary Petrology, v. 52, p.537-545.
- Nelson, P. H., 1994**, Permeability-porosity relationships in sedimentary rocks: Log Anal., v. 35/3, p. 38-62.
- Nelson, P. H., 2000**, Evolution of permeability-porosity trends in sandstones: In: Transactions of the SPWLA 41st Annual Logging Symposium, Society of Professional Well Log Analyst, Houston, Texas, p. MM1-14.

- Nickel, E., 1973**, Experimental dissolution of light and heavy minerals in comparison with weathering and interstitial solution: Contribution to Sedimentology, v. 1, p. 1-68.
- Oelkers, E. H., Bjørkum, P., A., and Murphy, W. M., 1996**, A petrographic and computational investigation of quartz cementation and porosity reduction in North Sea sandstones: American Journal of Science, v. 296, p. 420-452.
- Ollier, C. D., 1991**, Aspects of silcrete formation in Australia: Z. Geomorph., N. F., v. 35, p. 151-163.
- Ollier, C. D., and Galloway, R. W., 1990**, The laterite profile, ferricrete and unconformity: Catena Verlage, Cremlingen, v. 17, p. 97-109.
- Omara, S., 1965**, A micropaleontological approach to the stratigraphy of the Carboniferous exposures of the Gulf of Suez region: Neues Jahrbuch Geologie Palaontologie Monatshefte, Stuttgart, v 6, p. 409-419.
- Omara, S., 1972**, An Early Cambrian outcrop in southwestern Sinai, Egypt: Neues Jahrbuch Geologie Palaontologie Monatshefte, Stuttgart, v 5, p. 306-314.
- Pate, C., 1989**, Assessing the relative importance of compaction processes and cementation to reduction of porosity in sandstones: Discussion: The American Association of Petroleum Geologists Bulletin, v. 73/10, p. 1270-1273.
- Pettijohn, F. J., 1975**, Sedimentary Rocks: 3rd edition, Harper and Row, New York, 628 p.
- Pettijohn, F. J., and Potter, P. E., 1964**, Atlas and glossary of primary sedimentary structures: Springer-Verlag, New York, 370 p.

- Pettijohn, F. J., Potter, P. E., and Siever, R., 1987**, Sand and sandstone: 2nd edition, Springer-Verlag, New York, 533 p.
- Pittman, E. D., 1972**, Diagenesis of quartz in sandstones as revealed by scanning electron microscopy: *Journal of Sedimentary Petrology*, v.42/3, p. 507-519.
- Pittman, E. D., 1979**, Porosity, diagenesis and production capability of sandstone reservoirs: Society of Economic Palaeontologist and Mineralogist, Tulsa, Special Publication No. 26, p. 159-174.
- Pryor, W. A., 1973**, Permeability-porosity patterns and variations in some Holocene sand bodies: *The American Association of Petroleum Geologists Bulletin*, v. 57, p. 162-189.
- Pye, K., 1983**, Depositional reddening of late Quaternary coastal dune sands, northeastern Australia: In: Wilson, R. C. L., ed., *Residual deposits, surface related-weathering processes and materials*: Geological Society of London, Special Publication No, 11, Blackwell Sci. Publ., Oxford.
- Reineck, H. E., and Singh, I. B., 1980**, Depositional sedimentary environments: 2nd edition, Springer Verlag, Berlin, 549 p.
- Retallack, G. J., 1990**, Soils of the past: An introduction to the paleopedology: Unwin Hymen, Boston, Mass., 250 p.
- Rodriguez, J., and Gutschick, R. C., 1970**, Late Devonian-Early Mississippian ichnofossils from western Montana and northern Utah: In: Crimes, T. P. and Harper, J. C., eds., *Trace Fossils*, Geological Journal, Special Issue, 3, p. 407-438.
- Rogers, J. J., and Head, W. B., 1961**, Relationship between porosity, median size and sorting coefficients of synthetic sands: *Journal of Sedimentary Petrology*, v. 31, p. 467-470.

- Said, R., 1961**, Tectonic framework of Egypt and its influence on distribution of foraminifera: American Association of Petroleum Geologists Bulletin, v. 45, p. 198-218.
- Said, R., 1962**, The Geology of Egypt: Elsevier, 377 p.
- Said, R., 1971**, Explanatory notes to accompany the geological map of Egypt: Geological Survey of Egypt, Paper No. 55, 26 p.
- Said, R., and Eissa, R. A., 1969**, Some microfossils from Upper Paleozoic rocks of western coastal plain of Gulf of Suez region, Egypt: Third Afric. Micropal. Coll. Cairo, p. 337-383.
- Salem, A. M. K., 1995**, Diagenesis and isotopic study of the Paleozoic clastic sequence (Cambrian and Carboniferous), southwest Sinai, Egypt: Unpublished Ph.D. thesis, Tanta University, Tanta, Egypt, 246 p.
- Salem, A. M. K., Abdel-Wahab, A. A., and McBride, E.F., 1998**, Diagenesis of shallowly buried cratonic sandstones, southwest Sinai, Egypt: Sedimentary Geology, v. 119, p. 311-335.
- Saner, S., and Sahin, A., 1999**, Lithological and zonal porosity-permeability distribution in the Arab-D reservoir, Uthmaniyah field, Saudi Arabia: American Association of Petroleum Geologists Bulletin, v. 38/2, p. 230-243.
- Schmidt, V., McDonald, D. A, and Platt, R. L., 1977**, Pore geometry and reservoir aspects of secondary porosity in sandstones: Bulletin of Canadian Petroleum Geology, v. 25, p. 271-290.
- Schmidt, V., and McDonald, D. A, 1979a**, The role of secondary porosity in the course of sandstone diagenesis: In: Scholle, P. A., and Schuldger, P. R., eds., Aspects of Diagenesis: Society of Economic

Palaeontologist and Mineralogist, Tulsa, Special Publication No. 26, p. 175-207.

Schmidt, V., and McDonald, D. A., 1979b, Texture and recognition of secondary porosity in sandstone diagenesis: In: Scholle, P. A., and Schuldger, P. R., eds., Aspects of Diagenesis: Society of Economic Palaeontologist and Mineralogist, Tulsa, Special Publication No. 26, p. 209-225.

Scholle, P. A., 1979, A color illustrated guide to constituents, textures, cements, and porosities of sandstones and associated rocks: The American Association of Petroleum Geologists, Memoir 28, 201 p.

Schwertmann, U., 1988, Occurrence and formation of iron oxides in various pedoenvironments: In: Stucki, J. W., Goodman, B. A., and Schwertmann, U., Iron in soils and clay minerals, Chapter 11: Reidel Publication, Dordrecht, Holland, p. 267-308.

Schwertmann, U., and Taylor, R. M., 1987, Iron Oxides: In: Dixon, J. B., and Weed, S. B., eds., Minerals in Soil Environments, Chapter 8: Soil Sci. Soc. Amer., Madison, Wisc., 2nd edition, p. 379-468.

Seilacher, A., 1964, Biogenic sedimentary structures: In: Imbrie, J., and Newell, N. D., eds., Approaches to Paleoecology, John Wiley, New York, p. 296-315.

Seilacher, A., 1967, Bathymetry of trace fossils: Marine Geology, v. 5, p. 413-425.

Seilacher, A., 1990, Paleozoic trace fossils: In: Said, R., ed., The Geology of Egypt, Chapter 32, A. A. Balkema, Rotterdam, 734 p.

Selley, R. C., 1985, Elements of petroleum geology: Freeman and Company, 449 p.

- Selley, R. C., 1988**, Applied sedimentology: Academic Press Limited, London, UK, 446 p.
- Sellwood, B. W., and Netherwood, R. E., 1984**, Facies evolution in the Gulf of Suez area: Sedimentation history as an indicator of rift initiation and development: *Modern Geology*, v. 9, p. 43-69.
- Shata, A., 1956**, Structural development of the Sinai Peninsula, Egypt: *Egyptian Desert Institute Bulletin*, v. 6, p.41-46.
- Shanmuggan, G., 1984**, Porosity in sandstones: Classification, recognition and significance: *West Texas Geological Society Bulletin*, v. 24, p. 8-12.
- Siebert, R. M., Moncure, G. K., and Lahann, R. W., 1984**, A theory of framework grain dissolution in sandstones: In: McDonald, D. A., and Surdam, R. C., eds., *Clastic Diagenesis: The American Association of Petroleum Geologists, Memoir No. 37*, p.163-175.
- Siever, R., 1959**, Petrology and geochemistry of silica cementation in some Pennsylvanian sandstones: In: Ireland, H. A., ed., *Silica in Sediments: Society of Economic Palaeontologist and Mineralogist, Special Publication No. 7*, p. 55-79.
- Smale, D., 1973**, Silcretes and associated silica diagenesis in southern Africa and Australia: *Journal of Sedimentary Petrology*: v. 43, p. 1077-1089.
- Sneider, R. M., Richardson, F. H., Paynter, D. D., Eddy, R. E., and Wyant, I. A., 1977**, Predicting reservoir rock geometry and continuity in Pennsylvanian reservoirs, Elk City field, Oklahoma: *J. Petrol. Tech.*, v. 29, p. 851-866.
- Soliman, S. M., and El-Fetouh, M. A., 1969a**, Lithostratigraphy of the Carboniferous Nubian type sandstone in west central Sinai, Egypt:

Proceedings of Sixth Arab Science Congress, Damascus, Pt. 4, p. 311-377.

Soliman, S. M., and El-Fetouh, M. A., 1969b, Petrology of the Carboniferous sandstones in west central Sinai, Egypt: *Egyptian Journal of Geology*, v. 13/2, p. 61-143.

Soliman, S. M., and El-Fetouh, M. A., 1970, Carboniferous of Egypt: Isopach and lithofacies maps: *The American Association of Petroleum Geologists Bulletin*, v. 54, p. 1918-1930.

Sommer, F., 1978, Diagenesis of Jurassic sandstones in the Viking Graben: *Journal of Geological Society of London*, v. 135, p. 63-68.

Sorby, H. C., 1880, On the structure and origin of non-calcareous stratified rocks: *Quaternary Journal of Geological Society of London*, v. 37, p. 49-92.

Strainsky, A., Bielski, M., Ecker, A., and Steinitz, G., 1983, Tracing the origin of salts in groundwater by Sr isotopic composition (the crystalline complex of the southern Sinai, Egypt): *Isot. Geosci.*, v. 1, p. 257-267.

Summerfield, M. A., 1983a, Silcrete: In: Goudie, A. S., and Pye, K., eds., *Chemical Sediments and Geomorphology*: Academic Press, London, p. 59-91.

Summerfield, M. A., 1983b, Petrography and diagenesis of silcrete from the Kalahari Basin and Cape Coastal Zone, southern Africa: *Journal of Sedimentary Petrology*: v. 53, p. 895-909.

Suttner, L. J., Basu, A. and Mack, G. H., 1981, Climate and the origin of quartzarenite: *Journal of Sedimentary Petrology*, v. 51, p. 1235-1246.

- Tardy, Y., and Nahon, D. B., 1985,** Geochemistry of laterites, stability of Al-goethite, Al-hematite and Fe³⁺-kaolinite in bauxites and ferricretes: An approach to the mechanism of concretion formation: *American Journal of Science*, v. 285, p. 865-903.
- Thiry, M., and Millot, G., 1987,** Mineralogical forms of silica and their sequence of formation in silcretes: *Journal of Sedimentary Petrology*, v. 57, p. 343-352.
- Thiry, M., and Milnes, A. R., 1991,** Pedogenic and groundwater silcretes at Stuart Creek Opal Field, South Australia: *Journal of Sedimentary Petrology*, v. 61, p. 111-127.
- Torrent, J., and Schwertmann, U., 1987,** Influence of hematite on the color of red beds: *Journal of Sedimentary Petrology*, v. 57, p. 682-686.
- Tucker, M. E., 1982,** The field description of sedimentary rocks: *Geological Society of London Handbook Series*, 2, Open University Press, Buckingham, UK, 112 p.
- Turner, P., 1980,** Continental red beds: *Developments in Sedimentology*, v. 29, Elsevier, Amsterdam, 562 p.
- van Loon, A. J., and Brodzikowski, K., 1987,** Problems and progress in the research on soft sedimentation deformations: *Sedimentary Geology*, v. 50, p. 167-194.
- von Engelhardt, W., 1960,** *Der porenraum der sedimente*: Berlin, Springer-Verlage, 207 p.
- Waldschmidt, W. A., 1941,** Cementing materials in sandstones and their influence on the migration of oil: *The American Association of Petroleum Geologists Bulletin*, v. 25, p. 1839-1879.

- Walker, T. R., 1960,** Carbonate replacement of detrital crystalline silicate minerals as a source of authigenic silica in sedimentary rocks: Geological Society of America Bulletin, v. 71, p. 145-152.
- Walker, T. R., 1967,** Formation of red beds in modern and ancient deserts: Geological Society of America Bulletin, v. 78: p. 353-368.
- Walker, T. R., 1974,** Formation of red beds in moist tropical climates: A hypothesis: Geological Society of America Bulletin, v. 85, p. 633-638.
- Walker, T. R., 1976,** Diagenetic origin of continental red beds: In: Falke, H., ed., The continental Permian in Central, West and South Europe: Reidel Publication, Dordrecht, Holland, p. 240-282.
- Walker, T. R., Waugh, B., and Crone, A. J., 1978,** Diagenesis in first-cycle desert alluvium of Cenozoic age, southwestern United states and northwestern Mexico: Geological Society of America Bulletin, v. 89, p. 19-32.
- Walker, T. R., Larson, E. E., and Hoblitt, R. P., 1981,** Nature and origin of hematite in the Moenkopi Formation (Triassic), Colorado Plateau: A contribution to the origin of magnetism in red beds: Journal of Geophysical Research, v. 86, p. 317-333.
- Waugh, B., 1970,** Formation of quartz overgrowths in the Penrith sandstone (Lower Permian) of northwest England as revealed by scanning electron microscopy: Sedimentology, v. 14, p. 309-320.
- Williams, G. E., 1970,** Origin of disturbed bedding in Torridon Group Sandstones: Scott. Journal of Geology, v. 6, p. 409-410.
- Williams, L. A., and Crerar, D. A., 1985,** Silica Diagenesis, II: General mechanism, Journal of Sedimentary Petrology, v. 55, p. 312-321.
- Willner, A. P., Ermolaeva, T., Stroink, L., Glasmacher, U. A., Giese, U., Puchkov, V. N., Kozlov, V. I., and Walter, R., 2001,** Contrasting

provenance signals in Riphean and Vendian sandstones in the SW Urals (Russia): constraints for a change from passive to active continental margin conditions in the Neoproterozoic: *Precambrian Research*, v. 110, p. 215-239.

Wilson, J. C., and McBride, E.F., 1988, Compaction and porosity evolution of Pliocene sandstones, Ventura Basin, California: *The American Association of Petroleum Geologists Bulletin*, v. 72, p. 664-681.

Wilson, M. D., and Pittman, E. D., 1977, Authigenic clays in sandstone: Recognition and influence on reservoir properties and paleoenvironment analysis: *Journal of Sedimentary Petrology*, v. 47, p. 3-31.

Worden, R. H., and Burley, S. D., 2003, Sandstone diagenesis, the evolution of sand to stone: In: Burley, S. D., and Worden, R. H., eds., *Sandstone Diagenesis, Recent and Ancient*, Reprint Series of the International Association of Sedimentologists, v. 4, p. 3-44. Blackwell Publishers, Oxford.

Worden, R. H., and Morad, S., 2003, Clay minerals in sandstones, controls in formation, distribution and evolution: In: Worden, R. H., and Morad, S., eds., *Clay Mineral Cements in Sandstones*, International Association of Sedimentologists, Special Publication No. 34, p. 3-41, Blackwell Publishers, Oxford.

Worthington, P. F., 2003, Effect of clay content upon some physical properties of sandstone reservoirs: In: Worden, R. H., and Morad, S., eds., *Clay Mineral Cements in Sandstones*, International Association of Sedimentologists, Special Publication No. 34, p. 191-211. Blackwell Publishers, Oxford.

Wright, V. P., Sloan, R. J., Garces, B. V., and Garvie, L. a. J., 1992,
Ground ferricretes from the Silurian of Ireland and Permian of the
Spanish Pyreness: *Sedimentary Geology*, v. 77, p. 37-49.

Young, S. W., 1976, Petrographic textures of detrital polycrystalline
quartz as an aid to interpreting crystalline source rocks: *Journal of
Sedimentary Petrology*, v. 46/3, p. 595-603.

APPENDIX

Abbreviations used in the Tables 4.1-4.4

Na	Samples from Naqus Formation
Ar	Samples from Araba Formation
Q	Quartz
MCQ	Monocrystalline quartz
MCQU	Undulose monocrystalline quartz
MCQN	Nonundulose monocrystalline quartz
MCQT	Total monocrystalline quartz
PCQ	Polycrystalline quartz
PCQ 2-3	Polycrystalline quartz of two to three crystals
PCQ 3-5	Polycrystalline quartz of four to five crystals
PCQ >5	Polycrystalline quartz of more than five crystals
U/N Ratio	Undulose/Nonundulose ratio
P/M Ratio	Polycrystalline/Monocrystalline ratio
Feld.	Feldspar
R.F.	Rock fragments
H.M.	Heavy Minerals
Det. Clay	Detrital clay
Qtz. Over.	Quartz overgrowth
Mic. Qtz.	Microcrystalline quartz
Kao. P.F.	Pore-filling kaolinite
Kao. Rep.	Replacement kaolinite
Kao. Total	Total kaolinite
Gyp.	Gypsum
Inter	Intergranular porosity
Intra	Intragranular porosity
OVSP	Oversized pores
Frac.	Fracture porosity
1 st Por.	Primary porosity
2 nd Por.	Secondary porosity
PCP	Pre-cement porosity
PLOSS-Comp	Porosity loss due to compaction
PLOSS-Cement	Porosity loss due to cementation
Comp Index	Compaction Index

obeyk.com

ARABIC SUMMARY

دراسات ترسيبية وعمليات ما بعد الترسيب وكفاءة الخزان
لبعض القطاعات السطحية وتحت السطحية من منطقة بلاعيم
والمناطق المجاورة لها ، جنوب غرب سيناء ، مصر.

رسالة مقدمة من
وائل سعيد على مطر

ماجستير في العلوم في الجيولوجيا - جامعة الملك فهد للبترول والمعادن - المملكة العربية السعودية

١٩٩٦

باحث مساعد بمعهد بحوث البترول

للحصول على
درجة دكتوراه الفلسفة في العلوم في الجيولوجيا

تحت إشراف

الأستاذ الدكتور حافظ شمس الدين عبد الوهاب
أستاذ المعادن والصخور - قسم الجيولوجيا - كلية العلوم - جامعة عين شمس

الأستاذ الدكتور إبراهيم محمد إبراهيم المنسى

أستاذ الجيولوجيا الرسوبية - قسم الاستكشاف

معهد بحوث البترول

الأستاذ الدكتور عنتر على عبد الوهاب

أستاذ الجيولوجيا الرسوبية - كلية التربية

جامعة كفر الشيخ

قسم الجيولوجيا - كلية العلوم

جامعة عين شمس

٢٠٠٧



In The Name Of Allah
The Most Merciful
The Most Gracious



صفحة العنوان

اسم الطالب : وائل سعيد على مطر

الدرجة العلمية : دكتوراه الفلسفة فى العلوم فى الجيولوجيا

اسم الكلية : العلوم

الجامعة : عين شمس

سنة التخرج : ١٩٨٩

سنة المنح : ٢٠٠٧

رسالة دكتوراه فلسفة فى العلوم فى الجيولوجيا

عنوان الرسالة : دراسات ترسيبية وعمليات ما بعد الترسيب وكفاءة الخزان لبعض القطاعات السطحية وتحت السطحية من منطقة بلايم والمناطق المجاورة لها ، جنوب غرب سيناء ، مصر.

اسم الطالب : وائل سعيد على مطر

لجنة الإشراف



١- الأستاذ الدكتور حافظ شمس الدين عبد الوهاب
أستاذ المعادن والصخور بقسم الجيولوجيا بكلية العلوم جامعة عين شمس



٢- الأستاذ الدكتور عنتر على عبد الوهاب
أستاذ الجيولوجيا الرسوبية بكلية التربية جامعة كفر الشيخ



٣- الأستاذ الدكتور إبراهيم محمد إبراهيم المنسى
أستاذ الجيولوجيا الرسوبية بقسم الاستكشاف معهد بحوث البترول

بتاريخ : / /

لجنة الممتحنين

- ١- ١ د. محمد حسان محمد عوض أستاذ الجيولوجيا ووكيل كلية العلوم جامعة الأزهر
٢- ١ د. جمال الدين محمد حسن أستاذ ورئيس قسم الجيولوجيا بكلية العلوم جامعة حلوان
٣- ١ د. حافظ شمس الدين عبد الوهاب أستاذ الجيولوجيا بكلية العلوم جامعة عين شمس

الدراسات العليا

أجيزت الرسالة بتاريخ / / ٢٠٠٧ ختم الإجازة

موافقة مجلس الكلية : / / ٢٠٠٧ موافقة مجلس الجامعة : / / ٢٠٠٧

شكر

اشكر السادة الأساتذة الذين قاموا بالإشراف وهم :

١- الأستاذ الدكتور حافظ شمس الدين عبد الوهاب
أستاذ المعادن والصخور بقسم الجيولوجيا بكلية العلوم جامعة عين شمس

٢- الأستاذ الدكتور عنتر على عبد الوهاب
أستاذ الجيولوجيا الرسوبية بكلية التربية جامعة كفر الشيخ

٣- الأستاذ الدكتور إبراهيم محمد إبراهيم المنسى
أستاذ الجيولوجيا الرسوبية بقسم الاستكشاف معهد بحوث البترول

وكذلك الهيئات الأتية :

١- قسم الجيولوجيا - كلية العلوم - جامعة عين شمس.

٢- قسم الاستكشاف - معهد بحوث البترول.

٣- معمل التحاليل الصخرية (إبرى كورلاب) - معهد بحوث البترول.

٤- شركة السويس للزيت (سوكو).

دراسات ترسيبية وعمليات ما بعد الترسيب وكفاءة الخزان لبعض القطاعات
السطحية وتحت السطحية من منطقة بلاعيم والمناطق المجاورة لها ،
جنوب غرب سيناء ، مصر.

رسالة مقدمة من
وائل سعيد على مطر

للحصول على
درجة دكتوراه الفلسفة في العلوم في الجيولوجيا

ملخص الرسالة باللغة العربية

تتضمن هذه الرسالة دراسات معدنية وبتروجرافية والعمليات المابعدية للصخور
الرملية لبعض القطاعات السطحية وتحت السطحية في منطقة بلاعيم (جنوب غرب
سيناء). كذلك تضم الدراسة تقييما لكفاءة الخزان ومدى ملاءمته كمكن للبتترول.
اعتمد الباحث في إجراء دراسته عن القطاعات تحت السطحية على بيانات موثقة
للخرائط وبعض السجلات والدراسات وأيضا عدد (٦٠ عينة لبية core
samples) من شركة السويس للزيت (سوكو).
وقد اختيرت مئة وخمسين عينة تمثل كل الأنماط الممكنة للحجر الرملي، درست
بالتفصيل فيها البنيات والأنسجة الصخرية التي تميز هذه القطاعات بتروجرافيا
ومعدنيا باستخدام الميكروسكوب المستقطب والميكروسكوب الإلكتروني الماسح
وحيود الأشعة السينية. و قام الباحث أيضا بتحديد المعاملات البتروفيزيقية للصخور
المدروسة، إضافة إلى التحليل الحجمي ودراسة مدى تأثير توزيع الحجم
ومعاملاتها على طبيعة هذه الصخور.

وتضم الدراسة السطحية للصخور الرملية دراسة جيولوجية لمتكونين هما: متكون
ناقوس (الكمبرى العلوى) ومتكون عربية (الكمبرى السفلى). وسجلت الدراسة
الحقلية المستفيضة للبنيات والتراكيب وجود أسطح السحنات ورواسب ملء القنوات
والتطبيق الكتلي والتطبيق المتدرج والسطائح النضدية والتطبيق المتقاطع والتطبيق
الملفوف والمتحرف كذلك تمت دراسة التمايز اللوني في الصخور المدروسة
وعروق الجبس والكالسيت التي تتداخل مع الحجر الرملي. وتم مناقشة ظروف
تكوين البنيات والتراكيب على أساس علمي منهجي وتمت أيضا دراسة البنيات التي
تكونت ميكانيكيا، وكذلك مدى تأثير الصخور المدروسة بأنواع التجوية المختلفة.
وكذلك تمت دراسة البنيات الأحيائية التي سجل وجودها بمناطق البحث. وتدل
الدراسات التي أجريت على هذه الصخور أنها أحجار رملية لها حجوم تتردد بين

دقيقة التحبب جدا إلى متوسطة التحبب. لكنها فى بعض الأحيان تكون غليظة التحبب. كذلك فهى رديئة الفرز عموماً، و تميل إلى أن تكون لها درجة فرز متوسطة، أو جيدة فى بعض الأحيان. وفى مجملها فهذه الصخور لها درجة فرز متوسطة. أما حبيبات الرمل فهى متباينة فى أشكالها بين الزاوى إلى الدائرى، أيضاً هى شبه زاوية إلى شبه دائرية بصفة عامة.

وتدل الدراسة البتروجرافية التفصيلية أن أحجار الرمل المدروسة يتكون بنيانها من الحبيبات والملاط والمسامية والوسط الفتاتى الذى يكون نسبة بسيطة للغاية يمكن إهمالها عند تقسيم هذه الصخور الرملية. وبالرغم من وجود معادن الصلصال المتغلغلة فى بعض العينات، إلا أن أصلها فى الغالب يعد ثانوياً (ما بعدى). وعلى أساس نسب المكونات المعدنية والصخرية المكونة، فإن الصخور المدروسة تصنف على أنها كوارتز أرينيت، وتتميز بصحبة معدنية ناضجة تتدرج إلى معادن فوق ناضجة. ونظراً لتأثر هذه الصخور بالعمليات المابعدية إلى حد كبير، فيمكن تسميتها " كوارتز أرينيت ما بعدى".

وقد دلت الدراسة المعدنية والبتروجرافية أن حبيبات الكوارتز قد توجد على هيئة بلورات مفردة التبلور أو عديدة التبلور وقد تكون ذات انطفاء متموج أو غير متموج. لكن أكثر أنواع الكوارتز شيوعاً هو المتميز بالانطفاء المتوازي غير المتموج.

ولوحظ ندرة معادن الفلسبار أو غيابها تماماً، وقد يكون ذلك راجعاً إلى ذوبانها بفعل المياه الأرضية فى الدورة الهيدرولوجية النشطة أو من خلال عمليات الدفن فى المراحل المابعدية التى أثرت فى صخور الحجر الرملى المدروسة. ومن الأدلة المهمة لتأكيد ندرة وجود معادن الفلسبار وجود الفجوات المتنامية الحجم سواء أكانت مملوءة بالملاط أم خالية منه. ومن أهم أنواع الملاط التى تملأ الفراغات بين حبيبات الحجر الرملى معادن الكوارتز والكالسيدونى والأوبال والصلصال والهيماتايت والجوثيت والهاليت والجبس، وكذلك معدن الكالسيت الذى يوجد فى الفجوات و الذى قد يكون من أصل فتاتى أو ملاطاً مكانى النشأة. كذلك توجد معادن الفلسبار البوتاسى ولكن بكمية ضئيلة للغاية. ويمثل الفتات الصخرى مكوناً قليل الشيوع بنسبة لا تتعدى ٠,٢٢% لمتكون ناقوس، ٠,١٨% لمتكون عربة، بينما فى القطاعات تحت السطحية تصل النسبة إلى ٠,١٤%. والفتات الصخرى فى مجمله قد يكون من أصل رسوبى أو نارى أو من أصل متحول.

وتعد معادن الصلصال ذات الأصل الفتاتى شبه نادرة الوجود فى الصخور الرملية المدروسة حيث إنها تكون ١,٠٦% من العماد الهيكلى لصخور متكون ناقوس بينما فى متكون عربة فهى تكون ٢,٨٦%. أما فى القطاعات تحت السطحية، فتوجد معادن الصلصال ذات الأصل الفتاتى بنسبة معدلها ٠,٢٣%.

وتكون المعادن الثقيلة صلبة معدنية تتمثل في معادن الزركون والتورمالين والروتيل والجارنت وسفين وايبيدوت وستوروليت وسيلمانيت ومونازيت وهورنبلند.

وتدل الدراسات التي أجريت لدراسة معادن الصلصال سواء باستخدام الميكروسكوب الإلكتروني الماسح وحيود الأشعة السينية أن الصلبة المعدنية لمعادن الصلصال تتكون من معدن كاولينيت (بصفة أساسية) مع وجود كميات قليلة من معادن الليت وسمكتيت والليت-سمكتيت؛ طبقة صلصالية مختلطة وبعضاً من معدن كلوريت.

وبالنسبة إلى دراسة مسامية الصخور الرملية المدروسة فقد قدرت على أساس حسابات ومعادلات رياضية طبقت على القطاعات الرقيقة، ووجد أن معدل مسامية الصخور في متكوني ناقوس و عربية ١٧,٥٧% ، ٣,٤٥% على الترتيب، بينما بلغت النسبة ٤,٨٩% لقطاعات تحت السطح. ودلت الدراسات التي أجريت باستخدام جهاز تعيين المسامية أن معدل مسامية الصخور في متكوني ناقوس و عربية ١٧,٨% ، ٢٣,١% على الترتيب، بينما سجلت ١٠,٤% لقطاعات تحت السطح. وبينت الدراسات التي أجريت باستخدام جهاز تعيين النفاذية أن معدل نفاذية الصخور في متكوني ناقوس و عربية ٢٨١٨ مللي دارسي ، ٥٠ مللي دارسي على الترتيب، بينما سجلت الدراسة ٩٢ مللي دارسي (نفاذية أفقية) و ١٠٠ مللي دارسي (نفاذية رأسية) لقطاعات تحت السطح.

وتدل الدراسات التفصيلية على وجود تغيرات ما بعدية كثيرة أثرت في الصخور المدروسة إلى حد كبير و تشمل هذه العمليات : الرشح الميكانيكي لمعادن الصلصال وذوبان المعادن غير الثابتة التي أدت إلى زيادة مسامية الصخر وكذلك تكوين معادن مكانية النشأة. وأيضاً تضاعف الصخر وتماسكه و ذوبان معادن الكربونات التي تكونت في مراحل سابقة ثم يلي كل هذه المراحل جميعاً تكوين الأنواع اللاحقة للملاط بأنواعه، مما أعطى صخور الحجر الرملي في مناطق الدراسة صورتها الحالية التي نطلق عليها

الحجر الرملي بوضعه الحالي The present-day sandstone