

## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
(1) Adult cardiac arrest algorithm.	7
(2) EEG shows discontinuous EEG background activity (burst-suppression pattern).	17
(3) EEG of post-cardiac arrest male patient, classified as low voltage.background, showing diffuse cerebral slowing with suppression of the background, calibration of 10 uv/cm.	17
(4) Correlation between EEG scale 1st day and GOS.	34
(5) Correlation between EEG scale 7 <sup>th</sup> day and GOS.	34
(6) Correlation between NSE 1 <sup>st</sup> day with GOS.	38
(7) Correlation between NSE 3rd day with GOS.	38
(8) ROC curve for NSE in 1 <sup>st</sup> and 3 <sup>rd</sup> days to predict bad GOS.	39
(9) Correlation between EEG scale 1 <sup>st</sup> day with NSE 1 <sup>st</sup> day.	40
(10) Correlation between EEG scale 7 <sup>th</sup> day with NSE 1 <sup>st</sup> day.	41
(11) Correlation between EEG scale 1 <sup>st</sup> day with NSE 3 <sup>rd</sup> day.	41

## LIST OF ABBREVIATIONS

<b>SCA</b>	: Sudden Cardiac Arrest
<b>CPR</b>	: Cardiopulmonary Resuscitation
<b>VF</b>	: Ventricular Fibrillation
<b>PEA</b>	: Pulseless Electrical Activity
<b>CHD</b>	: Coronary Heart Disease
<b>ACS</b>	: Acute Coronary Syndrome
<b>ACLS</b>	: Advanced Cardiac Life Support
<b>AHA</b>	: American Heart Association
<b>VT</b>	: Ventricular Tachycardia
<b>ECG</b>	: Electrocardiogram
<b>AEDs</b>	: Automated External Defibrillators
<b>ROSC</b>	: Return Ofspontaneous Circulation
<b>ETCO2</b>	: End-Tidal Carbon Dioxide
<b>GCS</b>	: Glasgow Coma Scale
<b>SSEPs</b>	: Somatosensory-Evoked Potentials
<b>EEG</b>	: Electroencephalography
<b>CT</b>	: Computed Tomography
<b>NSE</b>	: Neuron-Specific Enolase
<b>ICU</b>	: Intensive Care Unit
<b>MRS</b>	: Modified Rankin Scale
<b>GOS</b>	: Glasgow Outcome Scale