

LIST OF FIGURES

<u>Figure No.</u>	<u>Title</u>	<u>PageNo.</u>
1	QuECHERS multiresidue-method employed for the determination of the pesticide residues using LC-MSMS and GC-MSD in tomato fruits	30
2	Mycelial growth reduction (%) on the linear growth of <i>A. solani</i> in response to different concentrations of applied antioxidants on CDA medium	35
3	Mycelial growth reduction (%) on the linear growth of <i>A. solani</i> in response to different concentrations of bio-and chemical fungicides applied on CDA medium	38
4	Mycelial growth reduction (%) on the linear growth of <i>F. solani</i> in response to different concentrations of applied antioxidant on CDA medium	41
5	Mycelial growth reduction (%) on the linear growth of <i>F. solani</i> in response to different concentrations of bio-and chemical fungicides applied on CDA medium	44
6	Effect of the performed treatments with different tested compounds on the reduction percentages of root rot disease incited by <i>F. solani</i> (first season).	46
7	Effect of the performed treatments with different tested compounds on the reduction percentages of early blight disease, incited by <i>A. solani</i> (first season)	47
8	Effect of the evaluated treatments with different tested compounds on the reduction percentages of root rot disease, incited by <i>F. solani</i> (second season)	49
9	Effect of the evaluated treatments with different tested compounds on the reduction percentages of early blight disease, incited by <i>A. solani</i> (second season)	51
10	General mean and changes in peroxidase activity in tomato leaves treated with some antioxidants, bio-and chemical fungicides and inoculated with <i>A. solani</i> , at different periods following inoculation	53
11	General mean and changes in peroxidase activity in tomato leaves treated with some antioxidants, bio-and chemical fungicides and inoculated with <i>A. solani</i> , at different periods following inoculation	55
12	General mean and changes in polyphenol oxidase activity in tomato leaves treated with some antioxidants, bio-and chemical fungicides and inoculated with <i>A. solani</i> , at different periods following inoculation	57
13	General mean and changes in polyphenol oxidase activity in tomato leaves treated with some antioxidants, bio-and chemical fungicides and inoculated with <i>A. solani</i> , at different periods following inoculation	59
14	Dissipation of tested fungicides in tomato fruits at different periods post treatment	61

LIST OF ABBREVIATIONS

<i>Alternaria</i>	A.
Analysis of variance	ANOVA
and others	<i>et al.</i>
Dextrose agar	DA
Diameter	di
Disease severity	DS
Distilled water	DW
Early blight	EB
Efficacy ratio	ER
Figure	Fig.
Food and Agriculture Organization Statistics	FAOSTAT
<i>Fusarium</i>	F.
High performance liquid chromatography	HPLC
In a <u>test tube</u> or <u>Petri dish</u>	<i>In vitro</i>
Induced systemic resistance	ISR
Least significant difference	LSD
Liquid nitrogen	Liquid N ₂
Microliter	μl
Micrometer (micron)	μm
Millimeter	mm
Millimole	mM
Optical density	OD
Part per milion	ppm
Pathogenesis related proteins	PRs
Peroxidase	PO
Poly phenol oxidase	PPO
Potato dextrose agar	PDA
Reduction of growth percentage	RG (%)
Salicylic acid	SA
species (plural)	spp.
species (single)	sp.
Systemic acquired resistance	SAR