

STUDIES ON INTERACTIONS OF THE PETROLEUM  
DERIVATIVES (PROPANIL - SO<sub>2</sub>) WITH HERBICIDE

AND INSECTICIDE ON BARNYARDGRASS

AND RICE

Mehreshan, T. El-Mokadem, Zeinab, Y.M. Abou-Bakr  
and Faida, A.A. Sharara

Bot. Dept. Women's College, Ain Shams University Heliopolis,  
Cairo, Egypt. Egyptian Petroleum Research Institute.

SUMMARY

Pot experiments were carried out to estimate the efficiency of the synthesized petroleum aromatic derivative (propanil - SO<sub>2</sub>) as herbicide either separately or in combination with commercial herbicide thiobencarb and insecticides (Carbaryl, Carbofuran and Malathion) on barnyardgrass (Echinochloa crus - galli (L.) P. Beauv) and /or rice (*Oryza sativa* cv Giza 176).

The determination of the effectiveness of propanil - SO<sub>2</sub> in controlling barnyardgrass showed no significant results. After application of propanil - SO<sub>2</sub> with thiobencarb the injury symptoms appeared on the treated seedlings of either barnyardgrass or rice were nearly similar to those sprayed with thiobencarb alone.

The interaction effects between propanil - SO<sub>2</sub> and the examined insecticides showed slight reduction in rice dry weight and grain yield ranged from 1 - 3% after the application of the insecticides 1 to 4 days before and after propanil - SO<sub>2</sub> treatment respectively. The obtained data was not significant.

The vegetative growth and yield of rice plants were less injured when the time interval between insecticides and propanil treatment increased.

Rice plants were not affected by the application of the used insecticides plus propanil - SO<sub>2</sub>.

## INTRODUCTION

Chemical weed control is a miracle of our technological age and it has been at the fore front in technological achievement. It involves knowledge in the fields of chemistry and biology (Ashton and Crafts, 1981).

New and better herbicides are being continuously synthesized and developed. Yields of cereals, soybeans, cotton, sugar beet, and in any other crops have increased in some cases 100% after application of synthetic organic herbicides (Ashton & Crafts, 1981). The herbicides are classified into two major groups inorganic and organic.

Propanil is a member of amide herbicides. It is especially valuable in rice culture because of its high selectivity. Rice plants are 40 times more tolerant of propanil than barnyardgrass (Matsunaka, 1965). The resistance of rice to propanil has been attributed to the ability of these species to degrade propanil more rapidly than most weed species (Adachi, *et. al.* 1966).

Chemical weed control in direct - seeded rice demands astrict herbicides application timing because rice and weeds are of the same growth stage (De Datta and Bernasor, 1973). The ideal herbicides for weed control in rice


combines high selectivity and efficacy with safe application methods (Hassan *et al.*, 1990).

Interaction between herbicides and insecticides were reported by (Smith and Tugwell, 1975 and Mukhopadhyay and sen, 1981), It was found that the organophosphate and carbamate insecticides interact with the herbicide propanil to increase injury to rice (Bowling and Hudgins, 1966).

The aim of the present study was to test the efficiency of the synthesized propanil SO<sub>2</sub> as herbicides either separately or in combination with commercial herbicide and insecticide on the growth of barnyardgrass and rice.

### MATERIALS AND METHODS

#### Preparation of N-propanil 3,4 - dichlorobenzene sulphonamide (propanil SO<sub>2</sub>)



This compound was prepared in three steps:

- A. Mixing 1,2 dichlorobenzene with Cl SO<sub>3</sub> H in a solvent of CH<sub>2</sub>Cl<sub>2</sub> (Stewart; 1922).
- B. The produced dichlorobenzene sulphonyl chloride was treated with NH<sub>4</sub> OH according to (Somasekhara, 196B) to prepare 3,4- dichlorobenzene sulphonamide.
- C. 3,4 - dichlorobenzene sulphonamide (0.005 mol) was mixed with propionyl chloride (0.007 mal) and glacial acetic acid (5 ml) on boiling bath. The solid product was filtered, dissolved in Na HCO<sub>3</sub>.

acidified with glacial acetic acid and crystallized by ethanal to produce N-propionyl 3,4 - dichlorobenzene sulphonamide (El-Dib, 1978).

#### Commercial herbicides and insecticides used

Pesticides	Common	Chemical name	Trade name	Formulation
Herbicides	Propanil	3,4 - dichloropropionanilide S [(4 - chlorophenyl) methyl] diethyl carbamothioate.	Stam F -34	EC
	Thiobencarb		Saurn 50%	EC
Insecticides	Carbaryl	1-naphthyl methyl carbamate	Siven 85%	WP
	Carbofuran	2,3 dihydro - 2 - 2 - dimethyl - 7 - benzofuranyl methyl carbamate.	Furadan 5%	G
	Malathion	o,o- dimethyl phosphorodithioate ester of diethyl mercaptosuccinoate.	Malathion 80%	EC

Ec = Emulsifiable

WP = Witable Powder

G = Granules

#### pot experiment:

The experiments were carried out at Rice Research and Training Centre, Sakka, Kafr El-Sheikh under greenhouse conditions (average temperature 28 + 2 C during the day time and 18 C at night). seeds were obtained from the same Research Centre.

The effect of the synthesized herbicides propanil - SO<sub>2</sub> was tested either separately or in combination with desired commercial herbicides or insecticides on barnyardgrass (Echinochloa crus - galli (L.) P. Beauv) and rice (Oryza sativa)

a. Giza 176. Plants were grown in plastic pots 20 cm depth and 20 cm diameter. In each pot 4 Kg of clay loam soil was placed. Some chemical analysis of the used soil was as follows:

pH	Ec ds/m	Co <sub>3</sub> Meq/L	HCo <sub>3</sub> Meq/L	OM %	NH <sub>4</sub> ppm	No <sub>3</sub> ppm	P ppm	K ppm	Zn ppm
7.8	1.1	1.0	1.9	2.1	2.9	19.3	11	625	1.9

Fifteen seeds each of barnyardgrass and rice were planted per pot, after emergence the seedlings were thinned to 10 plants per pot. Pots were irrigated by saturating the soil several times during the early season. It was flooded first when the plants ranged from 15 to 25 cm tall. Water was maintained on the plants during the season except for draining one day before and reflooding one day after each treatment. Rice plants received 40 Kg nitrogen /fed as urea 46% at 25 days after seeding.

The synthesized propanil - SO<sub>2</sub> and all commercial herbicides and insecticides were used as Kg/fed in a total water emulsion of 200L/fed., with a CO<sub>2</sub> - pressurized back - pack sprayer at 2.11 Kg/cm<sup>2</sup> fitted with flat - fan spray nozzles # 1004. The boom was positioned 25 cm above each pot.

The Herbicides were applied at half -, one- two- and three- leaf stages of barnyardgrass and rice at different rates.

Growth stages of rice plants when treated with insecticides in relation to

propanil application were listed in the following Table:

Application time of insecticides		Rice growth stage	
10	DBP	1	Leaf stage
7	DBP	2	Leaf stage
4	DBP	2.5	Leaf stage
2	DBP	3	Leaf stage
1	DBP	3.5	Leaf stage
<hr/>			
1	DAP	3.5	Leaf stage
2	DAP	4	Leaf stage
4	DAP	4.5	Leaf stage
7	DAP	5	Leaf stage
10	DAP	5.5	Leaf stage

DBP = days before propanil application.

DAP = days after propanil application.

The insecticides carbaryl, carbofuran and malathion were applied at the rates of 1.2 Kg aifed wettable powder, 0.8 Kg aifed granules and 1.2 Kg aifed emulsifiable concentrate respectively in relation to the propanil or propanil - SO<sub>2</sub> treatment at the rate of 1.05 Kg aifed emulsifiable concentrate.

Fresh and dry weights of shoots of barnyardgrass and rice, plant height and leaf area of rice plant were recorded at 30 days after seeding. leaf area was determined using LI-3100 Area Meter. Total chlorophyll content of rice leaves were estimated at 5 days after seeding according to th procedure (Arnon, 1959).

After maturity the harvested grains were adjusted to 14% moisture (Smith, 1974). Rice grain yield in grams per pot were obtained. The total nitrogen

content of rice grains was determined (Jackson, 1967). The starch content of rice grains was estimated using the method of (Yoshida *et al.*, 1972).

The results were statistically analyzed and significance of treatment differences was estimated through L.S.D. (Snedecor and Cochran, 1969).

**Effect of propanil, propanil - SO<sub>2</sub>, thiobencarb and their combinations on barnyardgrass and rice:**

Effective weed control programmes have been developed for the most severe weeds of rice (barnyardgrass) were based primarily on the use of propanil and other herbicides (Smith *et al.*, 1977 and Eastin, 1981).

In the present investigation we tried to compare the effects of either propanil or propanil - So with thiobencarb when applied at different stages of growth for controlling barnyardgrass.

Both barnyardgrass and rice plants developed dark green leaves and stunting within 1 - 3 days after thiobencarb application, while the plants showed scorching and stunting following the application of propanil plus thiobencarb. The symptoms were obvious at higher concentrations and when the treatments were done at the early stages of growth one - and two - leaf stage. Rice plants recovered from the injury within 7 - 15 days after treatments.

Table (1) represented that either propanil or thiobencarb caused significant reduction in fresh and dry weights of barnyardgrass at one - leaf stage. Weed injury was ranged from 16 - 87% and 28 - 64% following the application of propanil and thiobencarb respectively. The barnyardgrass - control

was decreased at late stages of growth (two - and three - leaf stage). However, propanil tank-mixed with thiobencarb provided excellent control of barnyardgrass (72 - 100%) at higher concentrations of propanil plus thiobencarb 0.35 + 1.0, 0.70 + 0.5, 0.70 + 0.75, 0.70 + 1.0, 1.05 + 0.5, 1.05 + 0.75 and 1.05 + 1.0 Kg aifed when applied at one - and two - leaf stage of growth.

It was clear from the experimental results that postemergence tank - mixture of propanil + thiobencarb caused an effective control of barnyardgrass more than single treatments. The phytotoxic effects were apparent when the herbicides were applied at lower stages of growth and at higher concentrations.

The obtained results were in agreement with (Smith and Khodayari, 1985 and Khodayari et al., 1989) who reported that propanil tank - mixed with thiobencarb postemergence provided an effective control of barnyardgrass than sequential application of them.

It was reported by different investigators (Smith 1975, Smith et al., 1977 and Hassan and Mahrous, 1989) and confirmed by the present work, that postemergence tank - mixture of propanil with thiobencarb was most effective for controlling barnyardgrass at one - and two - leaf stage of growth, while larger barnyardgrass plants were frequently not controlled.

Tables (2,3) showed that the effect of propanil, thiobencarb and their combinations on plant height, fresh weight, dry weight, leaf area and total chlorophyll content of rice. At one - leaf stage of rice there was severe reduction on the growth parameters after the application of the herbicides. The reduction was clearly observed as the rate of herbicides increased. At one - leaf

stage of growth plant height, fresh weight, dry weight and leaf area were reduced by 61% , 55%, 54% and 72% respectively following the application of thiobencarb at 1.0 Kg aiffed. This injury decreased at two - and three - leaf stage of growth. Postemergence tank - mixtures of propanil plus thiobencarb enhanced the injury of rice plants greatly at lower stages of growth and at higher concentrations of the herbicides.

Propanil at 1.05 Kg aiffed, propanil + thiobencarb at 1.05 + 0.5, 1.05 + 0.75 and 1.05 + 1.0 Kg aiffed caused 20% reduction in the total chlorophyll content of rice leaves. This reduction was not significant as shown in Table (3).

Table (4) represented that rice grains harvested from plant treated with the herbicides were severely affected at the early stages of growth. When the plants were sprayed with either propanil or thiobencarb at one - leaf stage the observed reduction in yield ranged from 29 - 54%, 29 - 56% respectively. This harmful effect increased by using tank - mixture of both herbicides at the same stage of growth. however, the reduction in rice grain yield was not statistically significant when the plants were treated with the herbicides either alone or in combination at the late stage of growth (three - leaf stage).

The total nitrogen content, crude protein and starch content of rice grains were not significantly affected by the herbicidal treatments at all rates and times of application as illustrated in Table (4).

Results from this investigation showed that the application of propanil in combination with thiobencarb increased rice injury as compared with single

treatments. This injury was clearly observed at one - and two - leaf stage of growth and at higher rates of the combined herbicides.

Previous investigators (Smith et al., 1977, Richard et al., 1981 and Smith, 1981) found that mixtures of propanil with thiobencarb applied early postemergence injured rice moderately, but the rice recovered and the grain yield was developed which supports the data reported herein.

Some herbicides injured rice more than propanil but injury for all treatments was inconsistent (Smith and Khodayari, 1985).

Trials were made to examine the phytotoxicity of propanil - SO<sub>2</sub> when used in combination with thiobencarb as shown in Table (5). After application of propanil - SO<sub>2</sub> with thiobencarb, the injury symptoms appeared on the treated seedlings of either barnyardgrass or rice were nearly similar to those sprayed with thiobencarb alone. Postemergence application of propanil - SO<sub>2</sub> at 0.35, 0.70 and 1.05 Kg aifed, in combination with thiobencarb at 0.50, 0.75 and 1.0 kg aifed resulted in significant reduction in fresh and dry weights of barnyardgrass at lower stages of growth and at higher concentrations. The percent injury ranged from 74 to 77% and 41 to 44% at one - and two - leaf stage of growth respectively at higher rates of propanil - SO<sub>2</sub> + thiobencarb 0.70 + 1.0 and 1.05+1.0 kg aifed as shown in Table (5).

Propanil - SO<sub>2</sub> in combination with thiobencarb was also tested on the vegetative growth and the crop yield of rice at different rates and times of application as indicated in Tables (6,7 & 8). The phytotoxic effects increased at lower stages of growth (one - and two - leaf stage) and at higher rates of

propanil - SO<sub>2</sub> + thiobencarb 0.7 + 1.0, 1.05 + 0.50, 1.05 + 0.75 and 1.05 + 1.0 kg aifed

Rice grain yield and total nitrogen content, crude protein & starch content of rice grains as influenced by propanil - SO<sub>2</sub> and thiobencarb were illustrated in Table (8). There was significant reduction in rice grain yield ranged from 65-68% at one - leaf stage when propanil - SO<sub>2</sub> + thiobencarb applied at rates of 0.70 + 1.0 and 1.05 + 1.0 kg aifed respectively. However, the percent injury of the total nitrogen content, crude protein and starch content of rice grains was not significant.

#### Herbicides - insecticides interactions on rice:

Losses in the yield of rice crop due to weeds and insects are quite severe. Biological - control is at present perhaps impractical for large scale use to combat these pestes. Large number of investigations have been made on the use of herbicides and insecticides separately, but very few investigations have been made on the combined use of herbicides and insecticides in rice crop. Very often weeds and insects infest the rice crop at the same time (Gifford, 1973 and Smith and Seaman, 1973). Hence herbicides and insecticides are needed at about the same time and one application of these combined pesticides will help to reduce the cost of operation. Therefore, it was of great necessity to study in detail the interaction of these herbicides and insecticides combinations.

With these ideas trials were made in the present investigation to drive more knowledge about the interaction of propanil or propanil - SO<sub>2</sub> with some insecticides on rice when they were applied at various times and rates before or after herbicidal treatments.

**Propanil or propanil - SO<sub>2</sub> and insecticides interactions on rice:**

Observations indicated that leaf chlorosis and necrosis of rice plants were less as the time interval between insecticides and propanil treatment increased. The rice recovered from the interacting effects within a 5 - week period after propanil treatment. Older rice plants recovered more quickly than younger ones.

Propanil is widely used as a selective herbicide for controlling barnyardgrass in rice and usually causes no significant injury to rice plants. However, certain insecticides interact adversely with propanil result in phytotoxic injury to rice plants.

The above observations were in agreement with those reported by (Smith and Tugwell, 1975 and Khodayari et al., 1986). The application of propanil plus carbofuran caused yellowing and burning of rice leaves but rice recovered about five weeks after application (Mukhopadhyay and Sen, 1981).

Synergistic phytotoxicity was apparent on rice plants when propanil was applied one day after carbaryl treatment where most rice plants were killed. Slight injury was noticed on plants treated with propanil two weeks after carbaryl treatment (Yih et al., 1968 b).

Table (9) showed that all insecticides (carbaryl, carbofuran and malathion) interacted with propanil at all times of application. Their interactions injured rice vegetatively. The degree of injury is influenced both by the type and the application time of the used insecticides in relation to the propanil application. The reduction in rice dry weight expressed as percent proved an

excellent indication of phytotoxicity. The synergistic effects in rice resulting in dry weight loss from 39% to 69% when the insecticides were applied 4 days to 1 day before or after propanil treatment respectively. Injury was lowered from 28% to 0% with time intervals of 7 to 10 days before or after propanil application.

After recovery of the vegetatively injured rice plants, grain yield was developed. Rice grain yield was reduced as the same trend observed in rice dry weight as shown in Table (9).

The phytotoxic symptoms exhibited by rice plants following the interaction of insecticides and propanil may be attributed to the inhibiting effect of the insecticides on the activity of rice arylacylamidase enzyme which detoxifies propanil causing loss of selectivity in rice.

Results of previous researchs obtained by different investigators (Bowling and Hudgins, 1966, Bowling and Flinchum, 1968 and Yih et al., 1968 a) indicated that the degree of rice plants injury was directly correlated with the degree of propanil hydrolyzing-enzyme inhibition caused by certain insecticides. The basis for the interaction between propanil and certain insecticides appeared to be an inhibition of rice arylacylamidase enzyme which metabolizes propanil to non-toxic compounds 3,4-dichloroaniline and propionic acid (Frear and Still, 1968, Matsunaka, 1968, El-Refai and Mowafy, 1973 and Matsunaka, 1981).

A mixture of propanil at 3 lb/acre and carbaryl at 0.5 lb/acre were applied to rice 4 inches tall caused 55% injury, compared with 5 and 0% injury from either propanil or carbaryl respectively (Smith, 1968).

Table (10) represented the interaction effects between propanil SO<sub>2</sub> and the examined insecticides. Slight reduction was observed in rice dry weight and grain yield ranged from 1% to 3% after the application of the insecticides 4 days to 1 day before or after propanil SO<sub>2</sub> treatment respectively. The obtained data was not significant.

#### **CONCLUSION:**

The results from the present study revealed that the response injury of barnyardgrass and rice to tested herbicides depends on the leaf - stage of growth at which they were applied. It is preferable to apply the herbicides at one - or two - leaf stage of barnyardgrass to achieve maximum control. However, it is safe to use herbicides after three - leaf stage of rice to avoid any phytotoxic effect.

The synthesized herbicides need knowledge in the fields of chemistry and biology and at least observational experience in the responses of common weeds and crops to them. Also weed and crop ecology and appreciation of the factors determining selectivity of the new herbicides are very important.

Generally herbicides must be applied at lowest rates by which maximum control of weeds is achieved.

#### **Acknowledgment:**

The authors express their gratitude to prof. Samy Mahmoud Hassan, who gave the opportunity to do this work in the laboratory of the Agriculture Dept., Rice Research and Training Centre, Sakka, Kafr El-Sheikh, for his help, kindness during the work.

## REFERENCES

- Adachi, M.; Tonegawa, K. and Uejima, T. (1966)  
Selective herbicidal action of 3,4 - dichloropropion - anilide. Penetration into plants and detoxication by their tissues. *Noyaku Seison Gijytsu* 14: 19 - 22.
- Arnon, D.I. (1959)  
Copper enzymes in isolated chloroplasts. Polyphenol oxidase in Beta vulgaris. *Plant Physiol.* 24: 1 - 15.
- Ashton, F.M. and Crafts, A.S. (eds.) (1981)  
Mode of Action of Herbicides. John Willy and Sons. New York.
- Bowling, C.C. and Flinchum, W.T. (1968)  
Interaction of propanil with insecticides applied as seed treatments on rice. *J. Econ. Entomol.* 61: 67 - 69.
- Bowling, C.C. and Hudgins, H.R. (1966)  
The effect of insecticides on the selectivity of propanil on rice. *Weeds* 14: 94 - 95.
- De Datta, S.K. and Bernasor, P.C. (1973)  
Chemical weed control in broadcast - seeded flooded tropical rice. *Weed Res.* 13: 351 - 354.
- Eastin, E.F. (1981)  
Weed management systems for U.S. rice. Pages 539 - 547 in D. Pimentel, ed. *Handbook of Pest Management in Agriculture*, Vol. 3. CRC Press, Inc., Boca Raton, FL.

- El-Dib, F.I. (1978)  
Some bio-cides from dihalogenated benzenes Ph. D. thesis Fac. of Sci  
Al-Azhar Univ. Cairo, Egypt.
- El-Refai, A.R. and Mowafy, M. (1973)  
Interaction of propanil with insecticides observed from soil and  
translocated into rice plants. Weed Sci. 21: 246 - 248.
- Frear, D.S. and Still, G.G. (1968)  
The metabolism of 3,4 - dichloropropionanilide in plants. Partial  
purification and properties of an arylacylamidase from rice.  
Phytochemistry 7: 913 - 920.
- Gifford, J.R. (1973)  
Insects and their control, Pages 151 - 154. in Rice in the United States:  
Varieties and Production. U.S. Dep. Agr. Handb. 289. U.S. Gov. Printing  
Office, Washington, D.C.
- Hassan, S.M. and Mahrous, F.N. (1989)  
Weed management for rice in Egypt. P. 330 - 337. In. proceeding of the  
4th EWRS Symposium on Weed Problems in Mediterranean Climates.  
Vol. II Problems of weed control in fruit, horticultural crops and rice.  
Valencia, Spain.
- Hassan, S.M.; Abd El-Rahman, A.A.M.; El-Serafy, A.M. and Ghanem, S.A. (1990)  
Evolution of different herbicide spraying schemes in rice. P. 455 - 463. In :  
Vol. II., Proc. 4th Conf. Agrons, 15 - 16 Sept., Cairo, Egypt.
- Jackson, M.L. (1967)  
Soil Chemical Analysis. India : Prentice Hall.
- Khodayari, K.; Nestasi, P. and Smith, R.J., Jr. (1989)  
Fenoxaprop for grass control in dry seeded rice (*Oryza sativa*). Weed  
Technology 3 : 131 - 135.

Khodayari, K. ; Smith, R.J., Jr. and Tugwell, P. (1986)

Interaction of propanil and selected insecticides on rice (Oryza sativa).

Weed Sci. 34: 800 - 803.

Matsunaka, S. (1968)

Propanil hydrolysis : Inhibition in rice plants by insecticides. Science. 100:

1360 - 1361.

Matsunaka, S. (1981)

Evaluation of rice weed control practices and research: World Perspective. p. 5 - 17 in Proc. Conf. Weed Control in Rice. Los Banos, Manila, Philippines.

Mukhopadhyay, S.K. and Sen, A.K. (1981)

Studies on interactions of herbicides with insecticides in rice crop. p. 463 - 468. In Proc. 8th Asian Pacific Weed Sci. Soc. Conf., Bangalore, India.

Richard, E.P., J. ; Miller, T.C. and Bowman, D.H. (1981)

Control of barnyardgrass in rice with thiobencarb. (Delta Branch, MAFES, MS. USA). Res. Rep. Miss. Agric. For. Exp. Stn. 6 (2) : 4.

Smith, R.J., Jr. (1968)

Control of grass and other weeds in rice with several herbicides. Ark. Agric. Exp. Sta. Rpt. Ser. 167: 38 pp.

Smith, R.J., Jr. (1974)

Responses of rice to postemergence treatments of propanil. Weed Sci. 22: 563 - 568.

Smith, R.J., Jr. (1975)

Herbicides for control of Leptochloa panicoides in water - seeded rice. Weed Sci. 23: 36 - 39.

Smith, R.J., Jr. (1981)

Herbicide programs for weed control in rice. Agric. Res. Results (ARR-S-8), Sci. Ed. Admin., U.S. Dep. Agric. 52 pp.

Smith, R.J., Jr. and Khodayari, K. (1985)

Herbicide treatments for control of weeds in dry seeded rice (*Oryza sativa*). Weed Sci. 33 : 686 - 692.

Smith, R.J., Jr. and Seaman, D.E. (1973)

Weeds and their control. Pages 135 - 140 in rice in the United States: Varieties and Production, U.S. Dep. Agr. Handb. 289. U.S. Gov. Printing Office, Washington, D.C.

Smith, R.J., Jr. and Tugwell, N.P. (1975)

Propanil - carbofuran interactions in rice. Weed Sci. 23: 176 - 178.

Smith, R.J., Jr. ; Flinchum, W.T. and Seaman, D.E. (1977)

Weed control in U.S. rice production. U.S. Dep. Agric. Handb. 497. U.S. Gov. Printing Office, Washington, DC. 78 pp.

Snedecor, G.W. and Cochran, W.G. (1969)

" Statistical Methods " 6th, Iowa State Univ., Press, Iowa, USA.

Somasekhara, S. (1968)

Indian, J. Chem. 6 (1) 19 - 23, c.f. Ph. D. Thesis, Some Biocides from Dihalogenated Benzenes. El-Dib, F.I. (1978). Fac. of Sci. Al-Azhar Univ., Cairo.

Stewart, J. (1922)

Chem. Soc. 121: 2555, c.f. Ph. D. Thesis. Some Biocides from Dihalogenated Benzenes. El-Dib, F.I. (1978). Fac. of Sci. Al-Azhar Univ., Cairo.

Ware, G.W. (1983)

Pesticides (Theory and Application). W.H. Freeman and Company. San Francisco.

Yih, R.Y. ; Mc Rae, D.H. and Wilson, H.F. (1968 a)

Mechanism of selective action of 3,4 - dichloropropionanilide. Plant Physio. 43: 1291 - 1296.

Yih, R.H.; Mc Rae, D.H. and Wilson, H.F. (1968 b)

Metabolism of 3,4 - dichloropropionanilide : 3,4 - dichloroaniline - lignin complex in rice plants. Science 161 : 376 - 377.

Yoschida, S. ; Forno, D.A. and Cock, J.H. (1972)

Laboratory manual for physiological studies of rice. The International Rice Res. Institute, Second Edition.

Table 1: Effect of propanil, thiobencarb and their combinations on the fresh and dry weights of barnyardgrass (*Echinochloa crus-galli*) at 30 days after seeding.

Herbicides	Rate of application Kg ai / fed	Fresh weight (g / pot)						Dry weight (g / pot)					
		Leaf stage & % reduction						Leaf stage & % reduction					
		1 lf	%	2 lf	%	3 lf	%	1 lf	%	2 lf	%	3 lf	%
Control (untreated)	-	21	0	21	0	21	0	3.67	0	3.51	0	3.69	0
Propanil	0.35	17.6	16	19.4	8	19.7	6	3.1	16	3.24	8	3.47	6
	0.70	4.6	78	7.8	63	12.5	41	0.8	78	1.3	63	2.13	42
	1.05	2.8	87	3.1	85	7.5	64	0.4	89	0.56	84	1.37	63
Thiobencarb	0.5	15.2	28	20.4	3	21.	0	2.69	27	3.40	3	3.69	0
	0.75	11.5	45	19.7	6	20.2	4	2.0	46	3.32	5	3.55	4
	1.0	7.5	64	16.8	20	18	14	1.3	65	2.8	20	3.15	15
Propanil + thiobencarb	0.35 + 0.5	14.5	31	19.8	6	20.8	1	2.53	31	3.38	4	3.60	2
	0.35 + 0.75	11.2	47	19.1	9	18.3	13	1.95	47	3.20	9	3.22	13
	0.35 + 1.0	5.9	72	14.8	30	15	29	1.03	72	2.45	30	2.63	29
Propanil + thiobencarb	0.70 + 0.5	3.5	83	7.0	67	11.7	44	0.60	84	1.17	67	2.05	44
	0.70 + 0.75	2.8	87	6.6	69	11.5	45	0.53	86	1.10	69	2.02	45
	0.70 + 1.0	2.3	89	5.9	72	11.3	46	0.40	89	1.03	71	2.0	46
Propanil + thiobencarb	1.05 + 0.5	2.0	91	6.4	70	8.9	58	0.34	91	1.08	69	1.59	57
	1.05 + 0.75	1.9	91	6.2	71	7.6	64	0.31	92	1.05	70	1.28	65
	1.05 + 1.0	0.0	100	0.0	100	6.6	69	0.0	100	0.0	100	1.2	68
L.S.D <sub>0.05</sub>		1.9		1.3		1.7		0.32		0.14		0.23	

Table : 2 Effect of propanil , thiobencarb and their combinations on plant height, fresh weight and dry weight of rice (*Oryza sativa*) cv Giza 176 at 30 days after seeding.

Herbicides application Kg ai/fed	Plant height (cm / plant)						Fresh weight (g / pot)						Dry weight (g / pot)					
	Leaf stage & % reduction						Leaf stage & % reduction						Leaf stage & % reduction					
	1IF	%	2IF	%	3IF	%	1IF	%	2IF	%	3IF	%	1IF	%	2IF	%	3IF	%
Control (untreated)	28	0	28	0	28	0	13	0	13	0	13	0	2.35	0	2.29	0	2.30	0
Propanil 0.35 0.70 1.05	24	14	28	0	28	0	11.6	11	12.9	1	13	0	2.1	11	2.31	0	2.35	0
	21	25	26.7	5	28	0	9.2	29	12.2	6	12.9	1	1.7	28	2.17	5	2.31	0
	13	54	23	18	26.9	4	5.2	60	10.9	16	12.5	4	0.95	60	1.9	17	2.22	4
Thiobencarb 0.5 0.75 1.0	18	36	28.1	0	28	0	9.5	27	13.5	0	13.7	0	1.7	28	2.34	0	2.36	0
	13	54	27.8	1	27.9	0	7.8	40	13	0	13.1	0	1.4	40	2.32	0	2.31	0
	11	61	27.3	3	27.8	1	5.9	55	12	8	12.9	1	1.08	54	2.11	9	2.28	1
Propanil + thiobencarb 0.35 + 0.5 0.35 + 0.75 0.35 + 1.0	17	39	27.9	0	28	0	7.0	46	12.9	1	13	0	1.3	45	2.29	0	2.31	0
	12	57	27.5	2	27.7	1	6.1	53	12.5	4	13	0	1.1	53	2.23	3	2.30	0
	10	64	26.9	4	27.4	2	4.9	62	12.1	7	12.9	1	0.9	62	2.10	8	2.28	1
Propanil + thiobencarb 0.70 + 0.5 0.70 + 0.75 0.70 + 1.0	15	46	26.5	5	27.6	1	5.9	55	12.6	3	13.2	0	1.08	54	2.23	3	2.32	0
	11	61	26.4	6	27.3	3	4.9	62	12.4	5	12.9	0	0.95	60	2.20	4	2.30	0
	9	67	25.2	10	26.5	5	3.8	71	10.4	20	11.4	12	0.68	71	1.8	21	2.0	13
Propanil + thiobencarb 1.05 + 0.5 1.05 + 0.75 1.05 + 1.0	10	64	23.1	18	27	4	2.0	85	9.21	29	11.4	12	0.33	86	1.6	30	2.0	13
	8	71	22.9	18	26.4	5	1.8	86	9.0	31	11.1	15	0.30	87	1.57	31	1.95	15
	7	75	22.2	21	26.2	6	1.38	89	8.5	35	10.9	16	0.28	88	1.48	37	1.90	17

Table 3 : Effect of propanil , thiobencarb and their combinations on leaf area and total chlorophyll content of rice (*Oryza sativa*) cv Giza 176.

Herbicides	Rate of application Kg ai / fed	Leaf area (cm <sup>2</sup> / pot ) at 30 days after seeding						Total * chlorophyll mg/g fresh wt.	
		Leaf stage & % reduction						2lf	% red
		1 lf	%	2 lf	%	3 lf	%		
Control (untreated)	-	96	0	96	0	96	0	5	0
Propanil	0.35	83	14	96	0	97	0	6	0
	0.70	61	37	89	7	96	0	5	0
	1.05	34	65	69	28	95	1	4	20
Thiobencarb	0.5	68	29	92	4	98	0	6	0
	0.75	44	54	90	6	97	0	6	0
	1.0	27	72	89	7	95	1	5	0
Propanil + thiobencarb	0.35 + 0.5	63	34	90	6	96	0	6	0
	0.35 + 0.75	41	57	89	7	95	1	5	0
	0.35 + 1.0	21	78	87	9	94	2	5	0
Propanil + thiobencarb	0.70 + 0.5	57	41	89	7	95	1	5	0
	0.70 + 0.75	39	59	89	7	94	2	5	0
	0.70 + 1.0	20	79	82	15	93	3	4	20
Propanil + thiobencarb	1.05 + 0.5	23	76	66	31	93	3	4	20
	1.05 + 0.75	20	79	64	33	91	5	4	20
	1.05 + 1.0	10	90	63	34	90	6	4	20
L.S.D <sub>0.05</sub>		11		8		N.S		N.S	

\* At 45 days after seeding

Table 4 : Effect of propanil , thiobencarb and their combinations on grain yield and total nitrogen , crude protein & starch content of rice-grains (*Oryza sativa*) cv Giza 176.

Herbicides	Rate of application Kg ai / fed	Grain yield ( g / pot )						Total nitrogen		Crude Protein		Starch content	
		Leaf stage & % reduction						mg / g grain					
		1 lf	%	2 lf	%	3 lf	%	2 lf	% red	2 lf	% red	2lf	%red
Control (untreated)	-	41	0	40	0	41	0	21	0	125	0	43	0
Propanil	0.35	29	29	39	3	41	0	20	5	125	0	42	2
	0.70	28	32	38	5	39	5	18	14	113	10	40	7
	1.05	19	54	37	8	38	7	18	14	113	10	40	7
Thiobencarb	0.5	29	29	40	0	42	0	19	10	119	5	42	2
	0.75	25	39	39	3	41	0	18	14	113	10	40	7
	1.0	18	56	39	3	39	5	17	19	106	15	40	7
Propanil + thiobencarb	0.35 + 0.5	18	56	38	5	42	0	19	10	119	5	41	5
	0.35 + 0.75	9	78	37	8	41	0	18	14	113	10	40	7
	0.35 + 1.0	7	83	36	10	39	5	17	19	106	15	40	7
Propanil + thiobencarb	0.70 + 0.5	8	81	37	8	41	0	18	14	113	10	40	7
	0.70 + 0.75	7	83	36	10	40	2	18	14	113	10	40	7
	0.70 + 1.0	6	85	34	15	39	5	17	19	106	15	39	9
Propanil + thiobencarb	1.05 + 0.5	4	90	32	20	40	2	18	14	113	10	40	7
	1.05 + 0.75	3	93	31	23	39	5	17	19	106	15	39	9
	1.05 + 1.0	2	95	30	25	39	5	17	19	106	15	39	9
L.S.D <sub>0.05</sub>		3		6		N.S		N.S		N.S		N.S	

Table 5 : Effect of propanil - SO<sub>2</sub>, thiobencarb and their combinations on the fresh and dry weights of barnyardgrass (*Echinochloa crus-galli*) at 30 days after seeding.

Herbicides	Rate of application Kg ai / fed	Fresh weight (g/pot)						Dry weight (g/pot)					
		Leaf stage & % reduction						Leaf stage & % reduction					
		1 lf	%	2 lf	%	3 lf	%	1 lf	%	2 lf	%	3 lf	%
Control (untreated)	-	23	0	23	0	23	0	3.89	0	3.97	0	3.91	0
Propanil SO <sub>2</sub>	0.35	23.1	0	23.2	0	23.6	0	3.90	0	3.99	0	3.93	0
	0.70	22.8	1	22.7	1	22.9	0	3.84	1	3.93	1	3.90	0
	1.05	21.5	7	21.9	5	22.5	2	3.65	6	3.79	5	3.85	12
Thiobencarb	0.5	15.9	31	21.5	7	22.9	0	2.7	31	3.65	8	3.90	0
	0.75	13.3	42	21	9	22.1	4	2.3	41	3.60	9	3.81	3
	1.0	7.1	69	18.5	20	21.5	7	1.19	69	3.15	21	3.69	6
Propanil-SO <sub>2</sub> +thiobencarb	0.35 + 0.5	15	35	21.2	8	22.5	2	2.55	34	3.60	9	3.83	2
	0.35 + 0.75	12.7	45	20.5	11	22.1	4	2.17	44	3.54	11	3.71	5
	0.35 + 1.0	6.9	70	17.6	23	21	9	1.15	70	3.07	23	3.57	9
Propanil-SO <sub>2</sub> +thiobencarb	0.70 + 0.5	14	39	21	9	22.1	4	2.4	38	3.63	9	3.74	4
	0.70 + 0.75	12.2	47	20.4	11	21.5	7	2.05	47	3.52	11	3.64	7
	0.70 + 1.0	5.9	74	13.5	41	20.5	11	1.0	74	2.34	41	3.40	13
Propanil-SO <sub>2</sub> +thiobencarb	1.05 + 0.5	13.3	42	20.5	11	22.1	4	2.3	41	3.54	11	3.71	5
	1.05 + 0.75	11.5	50	18.7	19	21.5	7	1.95	50	3.20	19	3.59	8
	1.05 + 1.0	5.3	77	12.9	44	20.3	12	0.9	77	2.22	44	3.33	15
L.S.D <sub>0.05</sub>		2.91		3.15		N.S		0.31		0.47		N.S	

Table : 6 Effect of propanil-SO<sub>2</sub>, thiobencarb and their combinations on plant height, Fresh weight and dry weight of rice (*Oryza sativa*) cv Giza 176 at 30 days after seeding.

Herbicides application Kg ai / fed.	Plant height (cm / plant)						Fresh weight (g / pot)						Dry weight (g / pot)						
	Leaf stage & % reduction			Leaf stage & % reduction			Leaf stage & % reduction			Leaf stage & % reduction			Leaf stage & % reduction			Leaf stage & % reduction			
	1lf	%	2lf	%	3lf	%	1lf	%	2lf	%	3lf	%	1lf	%	2lf	%	3lf	%	
Control (untreated)	29	0	29	0	29	0	17	0	17	0	17	0	2.87	0	2.89	0	2.89	0	
Propanil-SO <sub>2</sub>	0.35	29	0	29.1	0	28.9	0	17.1	0	17.1	0	17.3	0	2.90	0	2.88	0	2.89	0
	0.70	28.4	2	28.6	1	28.8	1	16.5	3	16.6	2	16.8	1	2.81	3	2.83	2	2.86	1
	1.05	28	3	28.4	2	28.5	2	16.1	5	16.3	4	16.5	3	2.74	5	2.79	4	2.81	3
Thiobencarb	0.5	16.8	42	29	0	29.1	0	12.3	28	16.4	4	17	0	2.06	28	2.77	4	2.89	0
	0.75	13.4	54	28.1	3	28.8	1	9.9	42	16.0	6	16.6	2	1.7	41	2.73	6	2.83	2
	1.0	10.7	63	28.1	3	28.4	2	7.0	59	15.5	9	16.3	4	1.18	59	2.63	9	2.79	4
Propanil-SO <sub>2</sub> +thiobencarb	0.35 + 0.5	16.2	44	28.9	0	29	0	12.0	29	16.3	4	16.8	1	2.03	29	2.78	4	2.86	1
	0.35 + 0.75	13.0	55	28.2	3	28.8	1	9.7	43	16.0	6	16.6	2	1.65	43	2.73	6	2.83	2
	0.35 + 1.0	10.1	65	28	3	28.3	2	6.8	60	15.3	10	16.1	5	1.15	60	2.63	9	2.74	5
Propanil-SO <sub>2</sub> +thiobencarb	0.70 + 0.5	15.9	45	28.8	1	29.3	0	11.5	32	16.0	6	16.5	3	1.98	31	2.73	6	2.81	3
	0.70 + 0.75	12.2	58	28.2	3	28.5	2	9.5	44	15.9	7	16.1	5	1.65	43	2.69	7	2.74	5
	0.70 + 1.0	9.5	67	27.9	4	28.4	2	6.5	62	15.3	10	15.8	7	1.10	62	2.59	10	2.70	7
Propanil-SO <sub>2</sub> +thiobencarb	1.05 + 0.5	15.0	48	28.4	2	29	0	11.0	35	15.8	7	16.3	4	1.89	34	2.70	7	2.78	4
	1.05 + 0.75	11.8	59	28	3	28.5	2	8.7	49	15.6	8	16.1	5	1.48	48	2.66	8	2.75	5
	1.05 + 1.0	9.2	68	27.8	4	28	3	6.0	65	15.2	11	15.9	7	1.01	65	2.59	10	2.69	7
L.S.D <sub>0.05</sub>	3.4		N.S		N.S		2.71		2.3		N.S		0.42		0.37		N.S		

Table :7 Effect of propanil-SO<sub>2</sub> , thiobencarb and their combinations on leaf area and total chlorophyll content of rice ( *Oryza sativa* ) cv Giza 176.

Herbicides	Rate of application Kg. ai / fed	Leaf area (cm <sup>2</sup> / pot) at 30 days after seeding						Total * chlorophyll mg/g fresh wt.	
		Leaf stage & % reduction						2lf	% red
		1 lf	%	2 lf	%	3 lf	%		
Control (untreated)	-	103	0	103	0	103	0	5	0
Propanil-SO <sub>2</sub>	0.35	104	0	103	0	103	0	6	0
	0.70	102	1	102	1	102	1	6	0
	1.05	100	3	101	2	101	2	5	0
Thiobencarb	0.5	61	41	98	5	103	0	6	0
	0.75	35	66	97	6	102	1	5	0
	1.0	27	74	95	8	101	2	5	0
Propanil-SO <sub>2</sub> +thiobencarb	0.35 + 0.5	57	45	97	6	102	1	6	0
	0.35 + 0.75	32	69	95	8	102	1	5	0
	0.35 + 1.0	26	75	94	9	101	2	5	0
Propanil-SO <sub>2</sub> +thiobencarb	0.70 + 0.5	56	46	96	7	102	1	6	0
	0.70 + 0.75	30	71	95	8	101	2	5	0
	0.70 + 1.0	25	76	93	10	100	3	4	20
Propanil-SO <sub>2</sub> +thiobencarb	1.05 + 0.5	55	47	96	7	101	2	5	0
	1.05 + 0.75	27	74	94	9	101	2	5	0
	1.05 + 1.0	24	77	93	10	100	3	4	20
L.S.D <sub>0.05</sub>		9		N.S		N.S		N.S	

\* At 45 days after seeding

Table 8 : Effect of propanil - SO<sub>2</sub>, thiobencarb and their combinations on grain yield and total nitrogen, crude protein & starch content of rice grains (*Oryza sativa*) cv Giza 176.

Herbicides	Rate of application Kg ai / fed	Grain yield (g / pot)						Total nitrogen		Crude Protein		Starch content	
		Leaf stage & % reduction						mg / g grain					
		1 lf	%	2 lf	%	3 lf	%	2 lf	% red	2 lf	% red	2 lf	% red
Control (untreated)	-	37	0	36	0	36	0	21	0	125	0	43	0
Propanil-SO <sub>2</sub>	0.35	37	0	38	0	38	0	20	5	125	0	43	0
	0.70	36	3	35	3	37	0	20	5	125	0	43	0
	1.05	35	5	35	3	35	3	19	10	119	5	42	2
Thiobencarb	0.5	26	30	37	0	38	0	19	10	119	5	41	5
	0.75	22	41	35	3	36	0	18	14	113	10	40	7
	1.0	15	60	34	6	35	3	17	19	106	15	40	7
Propanil-SO <sub>2</sub> + thiobencarb	0.35 + 0.5	25	32	36	0	37	0	19	10	119	5	41	5
	0.35 + 0.75	21	43	35	3	35	3	18	14	113	10	40	7
	0.35 + 1.0	14	62	34	6	35	3	17	19	106	15	40	7
Propanil-SO <sub>2</sub> + thiobencarb	0.70 + 0.5	23	38	35	3	36	0	18	14	113	10	41	5
	0.70 + 0.75	19	49	35	3	35	3	18	14	113	10	40	7
	0.70 + 1.0	13	65	34	6	35	3	17	19	106	15	39	9
Propanil-SO <sub>2</sub> + thiobencarb	1.05 + 0.5	21	43	35	3	35	3	18	14	113	10	40	7
	1.05 + 0.75	18	51	34	6	35	3	18	14	113	10	40	7
	1.05 + 1.0	12	68	34	6	34	6	17	19	106	15	39	9
L.S.D <sub>0.05</sub>		5		N.S		N.S		N.S		N.S		N.S	

Table: 9 Rice dry weight ( 30 days after seeding ) and grain yield as influenced by propanil and insecticides interactions .

Time of insecticide application *	Carbaryl				Carbofuran				Malathion			
	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	% red.	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	% red.	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	% red.
Control (untreated)	4.25	0	58	0	4.25	0	58	0	4.25	0	58	0
10 DBP	3.9	8	54	7	4.05	5	55	5	4.16	2	57	2
7 DBP	3.08	28	51	12	3.29	23	52	10	3.59	16	55	5
4 DBP	2.0	53	32	45	2.2	48	34	41	2.6	39	40	31
2 DBP	1.5	65	29	50	1.6	62	31	47	1.78	58	37	36
1 DBP	1.3	69	22	62	1.5	65	25	57	1.7	60	31	47
1 DAP	1.65	61	33	43	2.14	50	38	35	2.14	50	41	29
2 DAP	1.74	59	35	40	2.11	50	40	31	2.42	43	44	24
4 DAP	2.28	46	39	33	2.6	39	45	22	2.48	42	49	16
7 DAP	3.8	11	52	10	3.9	8	51	12	3.83	10	55	5
10 DAP	3.9	8	55	5	4.1	4	54	7	4.25	0	58	0
L.S.D 0.05	4	4	3	4	4	4	3	4	4	4	3	3

\*DBP = days before propanil application  
 DAP = days after propanil application

Table 10 Rice dry weight (30 days after seeding) and grain yield as influenced by propanil-SO<sub>2</sub> and insecticides interactions.

Time of insecticide application *	Carbaryl			Carbofuran			Malathion			
	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	Dry wt. (g/pot)	% red.	Grain yield (g/pot)	
Control (untreated)	3.19	0	50	3.17	0	50	3.17	0	50	
10 DBP	3.2	0	52	3.18	0	52	3.18	0	51	
7 DBP	3.16	1	51	3.16	0	51	3.13	2	50	
4 DBP	3.15	1	50	3.15	1	50	3.11	2	49	
2 DBP	3.12	2	49	3.11	2	49	3.09	3	49	
1 DBP	3.08	3	48	3.07	3	49	3.07	3	48	
1 DAP	3.1	3	48	3.08	3	48	3.08	3	49	
2 DAP	3.14	2	49	3.12	2	49	3.12	2	49	
4 DAP	3.16	1	49	3.15	1	50	3.1	2	50	
7 DAP	3.19	0	50	3.15	1	51	3.14	1	50	
10 DAP	3.20	0	51	3.17	0	52	3.17	0	51	
L.S.D 0.05	N.S		N.S		N.S		N.S		N.S	

\* DBP = days before propanil - SO<sub>2</sub> application  
 DAP = days after propanil - SO<sub>2</sub> application