

Mitotic and Meiotic Effects of
Trifluralin on *Vicia faba*

By

HODA HAKIM and ANAEL SHEHAB
Botany Dept. Univ. Coll. for Girls
El Shams University

C A I R O

(141)

(1)

I N T R O D U C T I O N

The herbicide trifluralin (a, a, a, trifluoro -2, b - dinitro - N - N - dipropyl toluidine) is used as a weed control in cotton. It is a soil active compound. Growth and physiological effects of the herbicide were studied by Kempen (1965), Standifer and Thomas (1965), Ne^eg^e et al (1968) and Schultz^e et al (1968). Mitotic effects were studied by Bayer et al (1967) on cotton, and Hackaylo and Amato (1968), on corn and cotton.

In this study mitotic and meiotic effects of trifluralin on Vicia faba were experimented.

MATERIAL and METHODS

Seeds of Vicia faba (Var. Giza I) were used in this test. For mitotic studies roots 4 days old were dipped in well aerated experimental solutions (0.1, 1, 10, 100 and 500 ppm.) for 24 and 72 hours. Tap water was used in control experiments. All observations were made from Feulgon permanent root tip squash preparations. For meiotic studies two types of treatments were carried on:-

1. Flower buds were directly treated with an aqueous solution of trifluralin (500 ppm.) using moisted piece of cotton for 3 hours, buds were gathered 24 hours after treatment. Moisted piece of cotton with tap water was used for controls.
2. Soil treatment, seeds were sown in pots and as shoot sprouts, they were irrigated regularly with the herbicide solution (10 ppm.) until flowering when flower buds were collected. Tap water was used for irrigating control plants. Pollen mother cells were examined from permanent preparations using aceto-carmin

smear method.

Stainability of pollen grains with aceto-carmin was taken as an index for determining pollen sterility.

Length of pollen grains was accurately measured by ocular micrometer.

RESULTS and DISCUSSION

The inhibiting mitotic effects of trifluralin (Table I) agree with those observed by Tablert (1965), ¹⁹⁶⁶ Bayer et al (1967) and Hačlavý and Amato (1968).

After 24 hours treatments the percentage of metaphases increased gradually on the expense of prophase, while after 72 hours treatments the percentage of metaphases increased markedly on the expense of both prophase and ana-telophases (table I). Generally the total percentage of abnormal cells increased with increase of time of treatment and concentration (table I). Stickness was a common abnormality in 72 hours treatment. Accumulation of metaphases may be attributed to different degrees of spindle disturbances shown in the presence of prophase-

metaphases, ball and star-shaped metaphases, lagging chromosomes and scattering of all the chromosomes all over the cell (Fig. 1-2-3-4). Prophase-metaphases has been observed by many others after treatment of colchicine (barber and callan 1942), ethylene glycol (Damato 1948) chromosomes remain nearly in their arrangement as they were during prophase due to inhibition of spindle formation. Amer (1965) considered star-metaphase type as a fore-step of complete disturbance of the spindle. Barthelmeas (1977) attributed the phenomenon of lagging chromosomes to the adhesion of the centromeres of one or more chromosomes to the outer layer of plasma while other chromosomes continue moving to the equatorial plate.

Anaphase abnormalities were: sticky anaphases with a bridge, c-anaphases and tetripolar anaphases (Fig. 6, 7) which ensure evidences of spindle disturbances. The polyploidising action of the herbicide is indicated by the presence of tetraploid and polyploid cells resulting in giant nuclei and restitution nuclei (Fig. 7, 8). Multipolarity resulted in polynucleate cells, (Fig. 5) Most of the above observations agree with those of Bayer et al (1967) on the effect of trifluralin on cotton. These abnormalities ensured that

114/16
-146-

(55)

Kinese is connected here with stathmokinese, that
multipolarity represented in multipolar anaphases
multinucleate cells are found together with c-meta
anaphases, polyploid cells and restitution nuclei.

The most common meiotic abnormalities were stick-
and sticky bridges (Fig. 9). After direct
treatment these abnormalities are obviously higher in
first division than the second division, Table II.

temporary effect of trifluralin on meiosis agree
the assumption of Amer and Ali (1968) that meiotic
kness induced by external agents seems to be temporary.
Table II shows also that in case of soil treatment the
percentage of abnormalities in the first and second
division did not show much variation. This may lead
one to believe that the induced abnormalities after soil
treatment may be the result of changes in the ionic
environment and metabolic activities thus leading to
secondary effects such as those observed.

Table II shows also that direct treatment of buds
(0 ppm.) had no effect on either fertility or length
of pollen grains, while soil treatment (10 ppm) induced
sterility and ^{abnormal} ~~abnormal~~ effect on pollen grains

22/2

shown in the increase in their length. These results strengthen the assumption that meiotic treatments with external agents exert temporary effects.

Comparing mitotic and meiotic effects of influralin it is observed that most of the mero-statokinetic abnormalities of the herbicide in mitosis disappeared during plant growth and were not traced during meiosis.

* * * * *




TABLE (I)

MITOTIC INDEX, TOTAL PERCENTAGE OF ABNORMALITIES AND PERCENTAGE OF DIFFERENT PHASES IN TREATED VICIA ROOT CELLS

Concentration	Mitotic Index		Total % of abnor.	Prophase %		Metaphase %		Ana-telophase %	
	24 h	72 h		24 h	72h	24h	72h	24h	72h
Control	82±5.3	79±5.1	-	32.8	32.9	24.4	25.2	42.6	41.7
0.1 ppm.	64±12	64±11.2	4.7	25	35.9	31.1	31.2	42.2	34.3
1 ppm.	64±9.4	65±4.4	12.3	23.4	21.6	39.1	53.9	37.3	24.6
10 ppm.	66±4.8	61±15.2	10.6	19.7	19.6	39.4	65.5	40.9	14.7
100 ppm.	65±13.2	54±15.5	12.2	24.6	24.1	30.7	53.7	41.5	22.2
500 ppm.	48±9	Toxic	29.1	22.9	Tox.	39.5	Tox.	37.5	Tox.

TABLE (II)
 PERCENTAGE OF ABNORMALITIES, POLLEN STERILITY AND LENGTH OF POLLEN
 GRAINS, OF TREATED VICIA PLANTS AFTER SOIL AND DIRECT TREATMENT

Type of	No. of counted cells	Total% abnorm	1st Division		2nd Division			Pollen Steri- lity	Length of p.-g. in micr.
			Blak- met.	Ara- teloph.	Total% of abnorm.	met.	ana-telo phase		
Control	2129	3.8	3.5	2.5	2.9	2.2	3.01	2.4	38.5
Soil	1101	19.6	38.5	15.1	26.1	25.7	11.5	19.3	55.65
Direct	1574	20.2	24.8	28.4	29.1	13.4	5.9	8.01	39.60

Literature Cited

- Amer, S. 1965. Mitotic effects of N-methyl -1-naphthyl carbamate "Sevin". *Cytologia* 30: 175-181
- Amer, S. and Ali, H.M. 1968. Meiotic effects of some phenols, *cytologia* 33 : 21 - 33
- Barber, H. N. and Callan, H. G. 1942; The effects of cold and colchicine on mitosis in the newt. *Proc. Roy. Soc.* 131 : 258-271
- Barthelmeß, A. 1957. Chemisch induzierte multipolare Mitosen. *Protoplasma* 48 : 546-561
- Bayer, D. E., Foy, C.L., Mallary, T.E. and Cutter, E.G. 1967 Morphological and histological effects of trifluralin on root development. *Amer. J. Bot.* 54. 945
- D'Amato, F. 1948. The effects of colchicine and ethylene glycol on sticky chromosomes in *Allium cepa*. *Hereditas* 34 : 83 - 103.
- Hacskeylo, J. and Amato, V.A. 1968. Effect of trifluralin on roots of corn and cotton. *Weed Sci.* 16 : 4