



أولاً: المراجع العربية

- ١ - ٢٠٠٣ حجازى ، ع.م. ، الإدارة المتكاملة للآفات - الطريق إلى بيئة نظيفة و غذاء آمن - ٨٥٠ صفحة - مكتبة المعارف الحديثة - الاسكندرية .
- ٢ - ٢٠٠٣ حجازى ، ع.م. و خفاجى ، و.ا. الحشرات النافعة - ٢٠٠ صفحة - مكتبة المعارف الحديثة - الاسكندرية
- ٣ - ١٩٩٣ حجازى ، ع.م. و البارونى ، م. ، المكافحة الحيوية - الجزء الأول - الحشرات الأكلة للحشرات - ٥٤٧ صفحة - منشورات جامعة عمر المختار - البيضاء - مطبعة قار يونس - ليبيا .

ثانياً : المراجع الأجنبية

- 4-Abbasher, A.A. (1994). Microorganisms associated with *Striga hermonthica* and possibilities of their utilization as biological control agents. Plits 12 (1), Margraf Verlag, Weikersheim, Germany.
- 5-Abbasher, A.A.; Hess, D.E.; Sauerborn, J. and Kroschel, J. (1996). Effect of different *Fusarium* spp. on seed germination of *Striga hermonthica* (sorghum and millet strains), *S. asiatica* and *S. gesnerioides*. In: M.T. Moreno, J.I. Cubero, D. Berner; D. Joel, L.J. Musselman, C. Parker (eds.), Advances in Parasitic Plant Research. Sixth International Parasitic Weed Symposium, Cordoba, Spain, 880-887.

- 6-Abbasher, A.A.; Kroschel, J. and Sauerborn, J. (1995). Microorganisms of *Striga hermonthica* in Northern Ghana with potential as biocontrol agents. *Biocontrol Science and Technology*, 5: 157-161.
- 7-Abbasher, A.A. and Sauerborn, J. (1992). *Fusarium nygamai*: a potential bioherbicide for *Striga hermonthica* control in sorghum. *Biological Control*, 2: 291-296.
- 8-Abbasi, S.A. and Nipanay, P.C. (1986). Infestations by aquatic weeds of the fern genus *Salvinia*: Its status and control. *Environmental conservation*, 13: 235-241.
- 9-Al-Azawi, A.F. (1967). Agromyzid leafminers and their parasites in Iraq. *Bulletin of Entomological Research*, 57: 285-287.
- 10-Al-Menoufi, O.A. (1986). Studies on *Orobanch* spp. 2. Fungi associated with *Orobanch crenata* Forsk. *Alex. J. Agric. Res.*, 31: 297-310.
- 11-Al-Menoufi, O.A. (1991). Crop rotation as a control measure of *Orobanch crenata* in *Vicia faba* fields. In: K. Wegmann, L.J. Musselman (eds.), *Progress in Orobanch research*. Eberhard-Karis-Universität, Tübingen, Germany, 241-247.
- 12-Al-Menoufi, O.A. (1994). The *Orobanch* problem and management in Egypt. In: A.H. Pieterse, J.A.C. Verkleij, S.J. Ter Borg (eds.), *Biology and Management of Orobanch*. Proceedings of the 3rd International Workshop on *Orobanch* and related *Striga* research, Amsterdam, The Netherlands, Royal Tropical Institute, 663-671.
- 13-Amsellem, Z.; Barghouthi, S.; Cohen, B.; Goldwasser, Y.; Gressel, J.; Hornok, L.; Kerényi, Z.; Kleifeld, Y.; Klein, O.; Kroschel, J.; Sauerborn, J.; Müller-Stöver, D.; Thomas, H.; Vurro, M. and Zonno, M.C. (2001). Recent advances in biocontrol of *Orobanch* (Broomrape) species. *Biocontrol*, 46: 211-228.
- 14-Andres, L.A. (1977). The biological control of weeds. In: *Integrated control of weeds*, J.D. Fryer and S. Matsunaka (eds.), Univ. of Tokyo Press, Tokyo, pp. 153-176.

- 15-Anonymous (1986). Plants eaten by goats. Agfact A7.5.1. Department of Agriculture, New South Wales, 3p.
- 16-Atkinson, I.A.E. (1964). Relations between feral goats and vegetation in New Zealand. Proceedings of the New Zealand Ecological Society 7: 39-44.
- 17-Aycock, R. (1966). Stem rot and other diseases caused by *Sclerotium rolfsii*. North Carolina Agricultural Experimental Station Technical Bulletin no. 174.
- 18-Bashir, M.O. and Musselman, L.J. (1984). Some natural enemies of *Striga hermonthica* in the Sudan. Tropical Pest Management 30, 211.
- 19-Bedi, J.S and Donchev, N. (1991). Results on mycoherbicides' control of sunflower broomrape (*Orobanche cumana* Wall.) under glasshouse and field conditions. In: Proceedings 5th International Symposium on Parasitic Weeds, Nairobi, Kenya (eds. J.K. Ransom, L.J. Musselman, A.D. Worsham and C. Parker), pp.76-82.
- 20-Borg, S.J. ter (1986). Effects of environmental factors on *Orobanche*-host relationships: a review and some recent results. In: S.J. ter Borg (ed.), Biology and control of *Orobanche*, Proceedings of a Workshop on Biology and Control of *Orobanche*. Wageningen, The Netherlands. LH/VPO, pp. 57-69.
- 21-Bourdôt, G.W. and Harvey, I.C. (1996). The potential of the fungus *Sclerotinia sclerotiorum* as a biological herbicide for controlling thistles in pasture. Plant Protection Quarterly, 11: 259-262.
- 22-Bourdôt, G.W.; Harvey, I.C.; Hurrell, G.A. and Saville, D.J. (1995). Demographic and biomass production consequences of inundative treatment of *Cirsium arvense* with *Sclerotinia sclerotiorum*. Biocontrol Science and Technology, 5: 11-25.
- 23-Briese, D.T. (1996). Potential impact of the stem-boring weevil *Lixus cardui* on the growth and reproductive capacity of *Onopordum* thistles. Biocontrol Science and Technology, 6: 251-261.

- 24-Briese, D.T. (1996). Life history of the *Onopordum capitulum* weevil *Larinus latus* (Coleoptera : Curculionidae). *Oecologia*, 105: 454-463.
- 25-Briese, D.T.; Sheppard, A.W. and Reifenberg, J.M. (1995). Open-field host-specificity testing for potential biological control agents of *Onopordum* thistles. *Biological Control*, 5: 158-166.
- 26-Briese, D.T.; Sheppard, A.W.; Zwölfer, H. and Boldt, E. (1994). Structure of the phytophagous insect fauna of *Onopordum* thistles in the northern Mediterranean basin. *Biological Journal of the Linnean Society*, 53: 231-253.
- 27-Calder, A.A. and Sands, D.P.A. (1985). A new Brazilian *Cyrtobagous* Hustache (Coleoptera : Curculionidae) introduced to Australia to control salvinia. *Journal of the Australian Entomological Society*, 24: 57-64.
- 28-Cary, P.R. and Weerts, P.G.J. (1984). Growth of *Salvinia molesta* as affected by water temperature and nutrition. III. Nitrogen and phosphorus interactions and effect of pH. *Aquatic Botany*, 19: 171-182.
- 29-Chaboudez, P.; Burdon, J.J. and Groves, R.H. (1993). Application for release of two strains of *Puccinia cardui-pycnocephali*, a rust fungus for the classical biological control of the noxious slender thistle weeds, *Carduus pycnocephalus* and *C. tenuiflorus*. CSIRO Division of Plant Industry, Internal Report, 13 pp.
- 30-Chaboudez, P.; Groves, R.H. and Burdon, J.J. (1993). *Puccinia cardui-pycnocephali*, a potential agent for the biological control of slender thistles in Australia. *Proceedings of the 10th Australian Weeds Conference*, Volume 1, PP. 84-8.
- 31-Clark, D.A.; Lambert, M.G.; Rolston, M.P. and Dymock, N. (1982). Diet selection by goats and sheep on hill country. *Proceedings of the New Zealand Society of Animal Production* 42: 155-7.
- 32-Crawley, M.J. (1989). Insect herbivores and plant population dynamics. *Annual Review of Entomology*, 34: 531-564.

- 33-CSIRO (1990). The host-specificity of *Aculus hyperici* (Liro) (Acarina: Eriophyidae) in relation to different species in the genus *Hypericum*. CSIRO Division of Entomology Report to AOIS. pp. 1-15.
- 34-Cubero, J.I. (1983). Parasitic diseases in *Vicia faba* L. with special reference to broomrape (*Orobanche crenata* Forsk.). In: P.D. Hebblethwaite (ed.), The faba bean (*Vicia faba* L.). London, UK, Butterworth, pp. 493-521.
- 35-Cullen, J.M. and Hasan, S. (1988). Pathogens for the control of weeds. Phil. Trans. R. Soc. Lond. B., 318: 213-224.
- 36-Davidson, A. (1963). Insects attacking *Striga* in Kenya. Nature 197, 923.
- 37-De Bach, P. (1970). Biological control of insect pests and weeds. Chapman and Hall Ltd.
- 38-Diatloff, G.; Lee, A.N. and Anderson, T.M. (1979). A new approach for *Salvinia* control. Journal of Aquatic Plant Management, 17: 24-27.
- 39-Dillon, S. P. and Forcella, F. (1985). Fluctuating temperatures break seed dormancy of catclaw mimosa (*Mimosa pigra*). Weed Science, 33: 196-198.
- 40-Dodd, A.P. (1940). The biological campaign against prickly pear. A.H. Tucker, Government Printer.
- 41-El-Helaly, A.F.; El-Arosi, H.; Hassouna, M.G. and Al-Menoufi, O.A. (1973). *Orobanche* spp. in Egypt. Egyptian Journal of Phytopathology, 5: 1-8.
- 42-Evans, H.C. (1995). Report on a survey for fungal pathogens of *Carthamus lanatus* (saffron thistle) in Greece. Report of Project CAR010, International Institute of Biological Control, Ascot, UK, pp. 1-4.
- 43-Flanagan, G.J.; Wilson, C. G. and Gillett, J. D. (1990). The abundance of native insects on the introduced weed *Mimosa pigra* in Northern Australia. Journal of Tropical Ecology, 6:219-230.

- 44-Forno, I.W. (1983). Native distribution of *Salvinia auriculata* complex and keys to species identification. *Aquatic Botany*, 17: 71-83
- 45-Forno, I.W. (1985). How quickly can insects control salvinia in the tropics? In Proceedings of the 10th Asian-Pacific Weed Science Society Conference, Chiang-mai, 1985, 1: 271-276.
- 46-Forno, I.W. (1987). Biological control of the floating fern *Salvinia molesta* in north-eastern Australia: plant-herbivore interactions. *Bulletin of Entomological Research*, 77: 9-17.
- 47-Forno, I.W. and Bourne, A.S. (1984). Studies in South America of arthropods on the *Salvinia auriculata* complex of floating ferns and their effects on *Salvinia molesta*. *Bulletin of Entomological Research*, 74: 609-621.
- 48-Forno, I.W. and Bourne, A.S. (1985). Feeding by adult *Cyrtobagous salviniae* on *Salvinia molesta* under different regimes of temperature and nitrogen content and the effects on plant growth. *Entomophaga*, 30: 279-286.
- 49-Forno, I.W.; Sands, D.P.A. and Sexton, W. (1983). Distribution, biology and host specificity of *Cyrtobagous singularis* Hustache (Coleoptera : Curculionidae) for the biological control of *Salvinia molesta*. *Bull. Ent. Res.*, 73: 85-95.
- 50-Garcia-Torres, L. and Lopez-Granados, F. (1991). Control of Broomrape (*Orobanche crenata* Forsk.) in broad bean (*Vicia faba* L.) with imidazolinones and other herbicides. *Weed Research*, 31: 227-235.
- 51-Greathead, D.J. (1984). The natural enemies of *Striga* spp. and their utilization as biological control agents. In: E.S. Ayensu, H. Doggett, R.D. Keynes, J. Marton-Lefevre, L.J. Musselman, C. Parker, A. Pickering (eds.), *Striga* Biology and Control. IRL Press Ltd. Oxford, UK, pp. 133-160.
- 52-Greathead, D.J. and Milner, J.E.D. (1971). A survey of *Striga* spp. (Scrophulariaceae) and their natural enemies in East Africa with a discussion on the possibilities of biological control. *Tropical Agriculture (Trinidad)*, 48: 111-124.

- 53-Habeck, D.H. and Thompson, C.R. (1994). Host specificity and biology of *Spodoptera pectinicornis* (Lepidoptera : Noctuidae). A biological control agent of water lettuce (*Pistia stratiotes*). *Biological Control*, 4: 263-268.
- 54-Hammad, S.; El-Arosi, H. and Al-Menoufi, O. (1967). *Phytomyza orobanchia* Kalt feeding on *Orobanche crenata* Forsk. in Egypt (Diptera : Agromyzidae). *Bulletin de la Société Entomologique d'Egypte*, 51: 141-144.
- 55-Harlan, J.R. (1982). Relationships between weeds and crops. W. Holzner and N. Numata (eds.), In: *Biology and ecology of weeds*, Dr. W. Junk Publishers, The Hague. ISBN9061936829, pp. 91-95.
- 56-Harley, K.L.S. (1992). A guide to management of *Mimosa pigra*. Csiro Canberra.
- 57-Harley, K.L.S. and Forno, I.W. (1992). *Biological control of weeds. A Handbook for Practitioners and Students*. Inkata Press.
- 58-Harman, H.M.; Syrett, P.; Hill, R.L. and Jessep, C.T. (1996). Arthropod introductions for biological control of weeds in New Zealand, 1929-1995. *New Zealand Entomologist*, 19: 71-80.
- 59-Harris, P. (1991). Classical biocontrol of weeds: its definition, selection of effective agents and administrative-political problems. *Canadian Entomologist*, 123: 827-849.
- 60-Harris, J.A. and Gill, A.M. (1997). History of the introduction and spread of St. John's wort (*Hyppicum perforatum* L.) in Australia. *Plant Protection Quarterly*, 12: 52-56.
- 61-Hasan, S. (1988). Biocontrol of weeds with microbes. In *Biocontrol of plant diseases*, vol. 1 (ed. K.G. Mukerji & K.L. Karg). Boca Raton, Florida: CRC Press.
- 62-Hassan, S. and Wapshere, A.J. (1973). The biology of *Puccinia chondrillina* a potential biological control agent of skeleton weed. *Ann. Appl. Biol.*, 74: 325-332.

- 63-Hegazi, E.M.; Kolaib, M.O.; El-Said, AM. and El-Nabawy, A. (1980). *Phytomyza orobanchia* Kalt. controlling the root parasite *Orobanche crenata* Forsk. in Egypt. Proc. 1st Conf. Pl. Prot. Res. Ins., Vol. III.
- 64-Hezewijk, MJ. van; A.H. Pieterse, A.H.; Saxena, M.C. and Borg, S.J. ter. (1987). Relationship between sowing date and the development of *Orobanche crenata* on faba bean and lentil in Syria. In: H.C. Weber and W. Forstreuter (eds.), Parasitic flowering plants, Proceedings of the 4th International Symposium on Parasitic Flowering Plants, Marburg, Germany, Philipps University, pp. 377-390.
- 65-Holst, P.J. (1980). Goats for weed control. Proceedings of the Post Graduate Committee of Veterinary Science, University of Sydney, 52: 187-189.
- 66-Holst, P.J. (1993). The role of goats in controlling weeds of table and pastures. Delfosse, E.S. (ed.). Pest of pastures: weed, invertebrate and disease pests of Australian sheep pastures. X + 434 pp. CSIRO Information Services, Melbourne.
- 67-Holst, P.J. and Campbell, M.H. (1985). The role of goats in the control of weeds of pastures. Australian Wool Corporation Research Review Conference, Leura, New South Wales.
- 68-Holzner, W. and Numata, M. (1982). Biology and ecology of weeds. Dr. Junk Publishers, The Hague-Boston-London.
- 69-Hosking, J.R.; Mc Fadyen, R.E. and Murray, N.D. (1988). Distribution and biological control of *Cactus species* in eastern Australia. Plant Protection Quarterly, 3: 115-123.
- 70-Howard, F.G. (1991). Environmental impacts of classical biological control. Annual Review of Entomology, 36: 485-509.
- 71-Hutchinson, J. and Dalziel, J.M. (1958). The Flora of West Tropical Africa 2nd edn. (revised by R.W.J. Keay). Crown Agents, London.

- 72-Jacobsohn, R. and Kelman, Y. (1980). Effectiveness of glyphosate in broomrape (*Orobanche* spp.) control in four crops. *Weed Science*, 28: 692-699.
- 73-Jong, J. de and Kok, T. (1978). The purification of waste water and effluents using marsh vegetations and soils. *Proc. EWRS 5th Symp. Aquatic Weeds*, Amsterdam: 135-142.
- 74-Kemmerling, W. (1978). Mechanische Bekämpfung unerwünschter Pflanzen in und an Gewässern. *Proc. EWRS 5th Symp. Aquatic Weeds*, Amsterdam.
- 75-Julien, M.H. (Ed.) (1992) *Biological Control of Weeds: a world catalogue of agents and their target weeds*. CAB International, Wallingford, 3rd edn.
- 76-Julien, M.H. (1993). Control of *Salvinia molesta* in Kakadu National Park. Final Report for the 1991 to 1993 Consultancy. Project 97. Unpublished report prepared for ANPWS, dated 15 April 1993, 28 pp.
- 77-Julien, M.H. and Bourne, A.S. (1988). Effects of leaf-feeding by larvae of the moth *Jamea multiplicalis* Guen. (Lep.; Pyralidae) on the floating weed *Salvinia molesta*. *Journal of Applied Entomology*, 106: 518-526.
- 78-Julien, M.H.; Bourne, A.S. and Chan, R.R. (1987). Effects of adult and larval *Cyrtobagous salviniae* on the floating weed *Salvinia molesta*. *Journal of Applied Ecology*, 24: 935-944.
- 79-Julien, M.H. and Griffiths, M.W. (1999). *Biological control of weeds, A world catalogue of agents and their target weeds*. Fourth edition, Antony Rowe, Chippenham.
- 80-Jupp, P.W. (1993). The biological control of St. John's wort in Australia. Project CS 113. Final Report to the Meat Research Corporation.
- 81-Jupp, P.W. and Cullen, J.M. (1996). Expected and observed effects of the mite *Aculus hyperici* on St. John's wort, *Hypericum perforatum*, in Australia. *Proceedings of the IX International Symposium on Biological Control of Weeds*, pp. 365-370.

- 82-Kassulke, R.C.; Harley, K.L.S. and Maynard G.V. (1990). Host specificity of *Acanthoscelides quadridentatus* and *A. puniceus* (Col.: Bruchidae) for biological control of *Mimosa pigra*. *Entomophaga*, 35: 85-96.
- 83-Khalaf, K.A. (1994). Intercropping fenugreek with faba bean or Egyptian clover: prospects for *Orobanche crenata* control. In: A.H. Pieterse, J.A.C. Verkleij, S.J. ter Borg (eds.), *Biology and management of Orobanche*. Proceedings of the 3rd International Workshop on *Orobanche* and related *Striga* Research, Amsterdam, The Netherlands, Royal Tropical Institute, 663-671.
- 84-Klein, O.; Kroschel, J. and Sauerborn, J. (1995). Efficacite de *Phytomyza orobanchia* Kalt. (Diptera : Agromyzidae) pour la lutte biologique contre L'orobanche au maroc. In: J. Kroschel, M. Abderabihi, H. Betz (eds.), *Advances in parasitic weed control at on-farm level. vol.II. joint action to control Orobanche in the WANA Region*. Margraf Velag, Weikersheim, Germany, 161-171.
- 85-Kok, L.T. and Surles, W.W. (1975). Successful biocontrol of musk thistle by an introduced weevil *Rhinocyllus conicus*. *Environ Ent.* 4: 1025-1027.
- 86-Kok, L.T.; Ward, R.H. and Grills, C.C. (1975). Biological studies of *Ceuthorrhynchidius horridus* Panzer, an introduced weevil for thistle control. *Ann. Ent. Soc. Am.*, 68: 503-505.
- 87-Korashi, A.A.; El-Borollosy, M.M.; Hassan, E.A.; Abo El-Suoud, M.R.; Zain El-Deen and Koraim, A. (1996). Hosts of *Orobanche* spp. and yield losses in Delta and Upper Egypt. In: M.T. Moreno, J.I. Cubero, D. Berner, D. Joel, L.J. Musselman, C. Parker (eds.), *Advances in parasitic plant research. Proceedings of the 6th International Symposium on Parasitic Weeds* Cordoba, Spain, 487-493.
- 88-Kovalev, O.V. (1977). Biological control of Weeds. *Zaschita rastenii* 4: 12-14. Review only in *Review Appl. Entomol. Series A*. 66: 171.

- 89-Kropff, M.J. and Schippers, P. (1986). Simulation of the growth of faba bean (*Vicia faba* L.) infested with broomrapes (*Orobancha crenata* Forsk.). In: S.J. ter Borg (ed.), Biology and control of *Orobancha*. Proceedings of a Workshop on Biology and Control of *Orobancha*. LH/VPO, Wageningen, The Netherlands, pp. 77-79.
- 90-Kroschel, J.; Abbasher, A.A. and Sauerborn, J. (1995). Herbivores of *Striga hermonthica* in northern Ghana and approaches to their use as biocontrol agents. *Biocontrol Science and Technology*, 5: 163-164.
- 91-Kroschel, J.; Hundt, A.; Abbasher, A.A. and Sauerborn, J. (1996). Pathogenicity of fungi collected in northern Ghana to *Striga hermonthica*. *Weed Research*, 36: 515-520.
- 92-Kroschel, J.; Jost, A. and Sauerborn, J. (1995). Insects for *Striga* control – possibilities and constraints. In: J. Kroschel, H. Mercer-Quarshie, J. Sauerborn (eds.), *Advances in Parasitic Weed Control at on-Farm Level. vol. I. joint action to control Orobancha in WANA Region*. Margraf Verlag, Weikersheim, Germany, 117-132.
- 93-Kroschel, J. and Klein, O. (1999). Biological control of *Orobancha* spp. with *Phytomyza orobanchia* Kalt., a review. In: J. Kroschel, M. Abderabihi, H. Betz (eds.), *Advances in Parasitic Weed Control at on-Farm Level. vol. II. joint action to control Striga in Africa*. Margraf Verlag, Weikersheim, Germany, 135-159.
- 94-Kroschel, J.; Sauerborn, J.; Kachelriess, S.; Hoffmann, V. and Mercer-Quarshie, H. (1996). Possibilities and constraints in implementing *Striga* control methods in African agriculture. *Proceedings First International Symposium on Food Security and Innovations: Successes and Lessons Learned*. University of Hohenheim, Germany.

- 95-Kukula, S.; Haddad, A. and Masri, H. (1985). Weed control in lentils, faba beans and chickpeas. In: M.C. Saxena and S. Varma (eds.), Faba beans, kabuli chickpeas, and lentils in the 1980s, Proceedings of the International Workshop held at ICARDA in 1983. Aleppo, Syria, ICARDA, pp. 169-177.
- 96-Legner, E.F.; Fisher, T.W. and Medved, R.A. (1973). Biological control of aquatic weeds in the lower Colorado river basin. Reprinted from Proceedings and Papers of Forty-First Annual Conference of the California Mosquito Control Association, INC, January 28-31.
- 97-Lekic, M. (1971). The role of the dipteran *Phytomyza orobanchia* Kalt. Agromyzidae in reducing parasitic phanerogam populations of the *Orobanche* genus in Vojvodina. *Savremena poljo-privreda*, 18: 627-637.
- 98-Linke, K.H.; Schnell, H. and Saxena, M.C. (1991). Factors affecting the seed bank of *Orobanche crenata* in fields under lentil based cropping systems in Northern Syria. In: J.K. Ransom, L.J. Musselman, A.D. Worsham and C. Parker (eds.), Proceedings of the 5th International Symposium on Parasitic Weeds. Nairobi, Kenya, CIMMYT, pp. 321-327.
- 99-Linke, K.H.; Vorlander, C. and Saxena, M.C. (1990). Occurrence and impact of *Phytomyza orobanchia* (Diptera : Agromyzidae) on *Orobanche crenata* (Orobanchaceae) in Syria. *Entomophaga*, 35 (4): 633-639.
- 100-Little, E.C.S. (1979). Handbook of utilization of aquatic plants. FAO, Rome, 176 pp.
- 101-Lonsdale, W.M. (1988). Litterfall in an Australian population of *Mimosa pigra*, an invasive tropical shrub. *Journal of Tropical Ecology*, 4: 381-392.
- 102-Lonsdale, W.M. and Abrecht, D.G. (1989). Seedling mortality in *Mimosa pigra*, an invasive tropical shrub. *Journal of Ecology*, 77: 371-385.

- 103-Lonsdale, W.M.; Harley, K.L.S. and Gillett, J.D. (1988). Seed bank dynamics in *Mimosa pigra*, an invasive tropical shrub. *Journal of Applied Ecology*, 25: 963-976.
- 104-Lopez-Granados, F. and Garcia-Torres, L. (1991). Estimation of *Orobanche crenata* seed production. In: K. Wegmann and L.J. Musselman (eds.), *Progress in Orobanche Research*. Tübingen, Germany, Eberhard-Karls University, pp. 92-98.
- 105-Lutzeyer, H.J.; Kroschel, J. and Sauerborn, J. (1994). *Orobanche crenata* in legume cropping: Farmers, perception difficulties and prospects of control; a case study in Morocco. In: *Biology and management of Orobanche*, Proceedings of the Third International Workshop on *Orobanche* and related *Striga* Research. A.H. Pieterse, J.A.C. Verkleij and S.J. ter Borg (eds.). Amsterdam, The Netherlands, Royal Tropical Institute, pp. 432-441.
- 106-Mahmoud, S.A. and Khalil, S.A. (1996). Overview of faba bean production in Egypt. In: W. Bertenbreiter, M. Sadiki (eds.), *Rehabilitation of faba bean*. Proc. Première séminaire du réseau maghrébin de recherche sur fève (remaveve), Rabat, Morocco, 39-42.
- 107-McFadyen, R.E. (1979). The cactus mealy-bug *Hypogeococcus festerianus* (Hem.: Pseudococcidae) an agent for the biological control of *Eriocereus martinii* (Cactaceae) in Australia. *Entomophaga*, 24: 281-287.
- 108-McFadyen, R.E. (1980). A *Cactoblastis* (Lep.: Phycitidae) for the biological control of *Eriocereus martinii* (Cactaceae) in Australia. *Entomophaga*, 25: 37-42.
- 109-McFadyen, R.E. (1984). Alien weeds. In: "The Brigalow Belt of Australia". Royal Society of Queensland Symposium, 23 October 1982, ed. A. Bailey, pp.177-182. (Department of Primary Industries: Brisbane).
- 110-McFadyen, R.E. (1985). Larval characteristics of *Cactoblastis* spp. (Lepidoptera : Pyralidae) and the selection of species for biological control of prickly pears.

- 111-McFadyen, R.E.C. (1998). Biological control of weeds. *Annu. Rev. Entomol.*, 43: 369-393.
- 112-Miller, I.L. (1992). Features of herbicides for the control of *Mimosa pigra*. Department of Primary Industry and Fisheries, Technical Bulletin No. 182.
- 113-Miller, I.L. and Forno, I.W. (1991). Focus on catclaw mimosa in Australia. *Aquatics*, 13 (4): 16-21.
- 114-Mitchell, D.S. (1974). Aquatic vegetation and its use and control. Unesco, Paris, 135 pp.
- 115-Moran, V.C. and Zimmermann, H.G. (1984). The biological control of cactus weeds: achievements and prospects. *Biocontrol News and Informations*, 5: 297-320.
- 116-Müller-Stöver, D.; Adam, M.A.; Al-Menoufi, O.A. and Kroschel, J. (1994). Importance of *Orobanche* spp. in two regions of Egypt—farmer's perceptions and difficulties, and prospects of control. In: J. Kroschel, M. Abderabihi, H. Betz (eds.), *Advances in Parasitic Weed Control at on-Farm Level*. vol. II. joint action to control *Orobanche* in WANA Region. Margraf Verlag, Weikersheim, Germany, pp. 37-52.
- 117-Murray, N.D. (1982). Ecology and evolution of the *Opuntia-Cactoblastis* ecosystem in Australia. In "Ecological Genetics and Evolution", eds. J.S.F. Barker and W.T. Starmer, pp. 17-30. (Academic Press, Sydney)
- 118-Murthy, D.V. (1960). Insects attacking *Striga*. *Nature*, 187, 959.
- 119-National Academy of Science (1968). Principles of plant and animal pest control. Vol. 2. Weed control. Chapter 6. The biological control of weeds. National Academy of Science Publication 1597: 86-119.
- 120-National Academy of Sciences (1976). Making aquatic weeds useful: Some perspectives for developing countries. Washington, 175pp.

- 121-Norambuena, H.: Kroschel, J. and Klen, O. (1995). Introduction of *Phytomyza orobanchia* for biocontrol of *Orobanche* spp. in Chile. In: J. Kroschel, M. Abderabihi, H. Betez (eds.), *Advances in Parasitic Weed Control at on-Farm Level*, vol. II. joint action to control *Orobanche* in WANA Region. Margraf Verlag, Weikersheim, Germany, pp. 197-204.
- 122-Pantone, D.J.; Williams, W.A. and Maggenti, A.R. (1989). An alternative approach for evaluating the efficacy of potential biological control agents of weeds. 1. Inverse linear model. *Weed science* 37: 771-777.
- 123-Philip, B.A.; Winks, C.J.; Joynt, P.W. and Sutherland, O.R.W. (1988). Current status of biological control of alligator weed in New Zealand. *Proceedings of the 41st New Zealand Weed and Pest Control Conference*, pp. 61-65.
- 124-Pitlo, R.H. (1978). Regulation of aquatic vegetation by interception of daylight. *Proc. EWRS 5th Symp. Aquatic Weeds*, Amsterdam, pp. 91-99.
- 125-Ridout, J. (1980). Water weed problems, potential utilization and control. *Miscellaneous Report No. 52*, London W8 55J.
- 126-Roberts, H.A. (1982). *Weed control handbook: Principles*. Seventh edition – Blackwell Scientific Publications, pp. 268-291.
- 127-Robson, T.O. (1978). The present status of chemical aquatic weed control. *Proc. EWRS 5th Symp. Aquatic Weeds*. Amsterdam, pp. 17-25.
- 128-Rolston, M.P.; Lambert, M.G.; Clark, D.A. and Devantier, B.P. (1981). Control of rushes and thistles in pastures by goat and sheep grazing. *Proceedings of the 34th New Zealand Weed and Pest Control Conference* pp. 117-121.
- 129-Room, P.M. (1986). *Salvinia molesta* - A floating weed and its biological control. In *The Ecology of Exotic Animals and Plants*, ed. R.L. Kitching, John Wiley and Sons, Brisbane, 164-186.

- 130-Room, P.M. and Thomas, P.A. (1986a). Nitrogen, phosphorus and potassium in *Salvinia molesta* Mitchell, in the field: effects of weather, insect damage, fertilizers and age. *Aquatic Botany* 24: 213-232.
- 131-Room, P.M. and Thomas, P.A. (1986b). Population growth of the floating weed *Salvinia molesta*: field observations and a global model based on temperature and nitrogen. *Journal of Applied Ecology*, 23: 1013-1028.
- 132-Room, P.M.; Forno, I.W. and Taylor, M.F.J. (1984). Establishment in Australia of two insects for biological control of the floating weed *Salvinia molesta*. *Bulletin of Entomological Research*, 74: 505-516.
- 133-Sands, D.P.A. and Kassulke, R.C. (1986). Assessment of *Paulinia acuminata* (de Greer) (Orthoptera : Acrididae) for the biological control of *Salvinia molesta* in Australia. *Entomophaga*, 31: 11-17.
- 134-Sands, D.P.A.; Schotz, M. and Bourne, A.S. (1983). The feeding characteristics and development of larvae of a salvinia weevil *Cyrtobagous* sp. *Entomologia Experimentalis et Applicata*, 34: 291-296.
- 135-Sauerborn, J. (1991). *Parasitic flowering plants: ecology and management*. Margraf Verlag, Weikersheim, Germany.
- 136-Sauerborn, J. (1991). The economic importance of the phytoparasites *Orobanche* and *Striga*. *Proceedings 5th International Symposium on Parasitic Weeds, Nairobi, Kenya* (eds. J.K. Ransom, L.J. Musselman, A.D. Worsham and C. Parker), 137-143.
- 137-Sauerborn, J. (1992). *Parasitic Flowering Plants: Ecology and Management*. Margraf Verlag, Weikersheim, Germany.

- 138-Sauerborn, J., Abbasher, A.A. and Kroschel, J. (1994). Biological control of parasitic weeds by phytopathogenic fungi. In: Biology and management of *Orobanchae*. Proceedings of third International Workshop on *Orobanchae* and related *Striga* Research. A.H. Pieterse, J.A.C. Verkleij and S.J. ter Borg (eds.). Amsterdam, The Netherlands. Royal Tropical Institute pp. 545-549
- 139-Sauerborn, J., Dorr, I., Abbasher, A.A., Thomas H and Kroschel J. (1996). Electron microscopic analysis of the penetration process of *Fusarium nygamai* a hyperparasite of *Striga hermonthica*. *Biological Control*, 7: 53-59.
- 140-Sauerborn, J.; Linke, K.H., Saxena, M.C. and Koch, W. (1989). Solarization: a physical control method for weeds and parasitic plants (*Orobanchae* spp.) in Mediterranean agriculture. *Weed Research*, 29: 391-397
- 141-Sauerborn, J., Wyrwal, G. and Linke, K.H. (1991). Soil sampling for broomrape seeds in naturally infested fields. In K. Wegmann and L.J. Musselman (eds.), *Progress in Orobanchae Research*. Tübingen, Germany. Eberhard-Karls University pp. 35-41
- 142-Saxena, M.C., Linke, K.H. and Sauerborn, J. (1994). Integrated control of *Orobanchae* in cool-season food legumes. In: *Biology and management of Orobanchae*. Proceedings of third International Workshop on *Orobanchae* and related *Striga* Research. A.H. Pieterse, J.A.C. Verkleij and S.J. ter Borg (eds.). Amsterdam, The Netherlands. Royal Tropical Institute pp. 419-430.
- 143-Scheepens, P.C. and van Zon, H.C.J. (1982). Microbial herbicides. In: *Microbial and viral pesticides* (ed. E. Kurstak), pp. 623-641. New York: Marcel Dekker.
- 144-Schmitz, D.D. (1990). The invasion of exotic aquatic and wetland plants in Florida: history and efforts to prevent new introductions. *Aquatics*, 12 (2): 6-13, 24.

- 145-Schnell, H.; Sauerborn, J. and Linke, K.H. (1994). Crop rotation in Syria and its impact on *Orobanche crenata* seed bank. In: Biology and management of *Orobanche*, Proceedings of third International Workshop on *Orobanche* and related *Striga* Research. A.H. Pieterse, J.A.C. Verkleij and S.T. ter Borg (eds.). Amsterdam, The Netherlands, Royal Tropical Institute.
- 146-Sheppard, A.W.; Aeschlimann, J.P.; Sagliocco, J.L. and Vitou, J. (1991). Natural enemies and population stability of the winter-annual *Carduus pycnocephalus* L. in Mediterranean Europe. *Acta Ecologica*, 12: 707-726.
- 147-Sheppard, A.W.; Aeschlimann, J.P.; Sagliocco, J.L. and Vitou, J. (1995). Below-ground herbivory in *Carduus nutans* (Asteraceae) and the potential for biological control. *Biocontrol Science and Technology*, 5: 261-270.
- 148-Sheppard, A.W.; Briese, D.T. and Michalakis, I. (1992). Host choice in field in *Genus larinus* (Coleoptera : Curculionidae) attacking *Onopordum* and *Cynara* (Asteraceae). Proceedings of the Eighth International Symposium on Biological Control of Weeds, 2-7 February, Lincoln University, Canterbury, New Zealand. Delfosse, E.S. and R.R. Scott (eds.). DSIR/CSIRO, Melbourne, pp. 605-614 (1994).
- 149-Shappard, A.W.; Cullen, J.M.; Aeschlimann, J.P.; Sagliocco, J.L. and Vitou, J. (1988). The importance of insect herbivores relative to other limiting factors on weed population dynamics: a case study of *Carduus nutans*. Proc. VII Int. Sym. Biol. Contr. Weeds, 6-11 March, Rome, Italy. Delfosse, E.S. (ed.). 1st Sperm. Patol. Veg. (MAF), pp. 211-219 (1989).
- 150-Sheppard, A.W.; Gullen, J.M. and Aeschlimann, J.P. (1994). Predispersal seed predation on *Carduus nutans* (Asteraceae) in southern Europe. *Acta Ecologica*, 15: 529-541.
- 151-Smith, M.C. and Webb, M. (1996). Estimation of the seed bank of *Striga* spp. (Scrophulariaceae) in Malian fields and the implications for a model of biocontrol of *S. hermonthica*. *Weed Research*, 36: 85-92.

- 152-Speight, R.I. and Whittaker, J.B. (1987). Interactions between the chrysomelid beetle *Gastrophysa viridula*, the weed *Rumex obtusifolius* and the herbicide asulam. *Journal of Applied Ecology*, 24: 119-130.
- 153-Storrs, M.J. and Julien, D.H. (1996). *Salvinia*, a handbook for the integrated control of *Salvinia molesta* in Kakadu national park. NT Uniprint, Darwin, NT.
- 154-Swarbrick, J.T. (1983). The introduction of *Mimosa pigra* into northern Australia. *Australian Weeds*, 2, 124.
- 155-Swarbrick, J.T.; Willson, B.W. and Hannan-Jones, M.A. (1995). The biology of Australian weeds 25 *Lantana camara* L. *Plant Protection Quarterly*, 10: 82-95.
- 156-Syrett, P. (1997). Is biological control of weeds working in New Zealand?. Evaluation of introduced biological control agents. *Bulletin of the Entomological Society of New Zealand*, 10: 91-99.
- 157-Syrett, P. and Emberson, R.M. (1997). The natural host range of beetle species feeding on broom, *Cytisus scoparius* (L.) Link (Fabaceae), in Southwest Europe. *Biocontrol Science and Technology*, 7: 309-326.
- 158-Syrett, P. and Smith, L.A. (1998). The insect fauna of four weedy *Hieracium* (Asteraceae) species in New Zealand. *New Zealand Journal of Zoology*, 25: 73-83.
- 159-Tawfik, M.F.S.; Awadallah, K.T. and Shalaby, F.F. (1976). Biology of *Phytomyza orobanchiae* Kalt. (Diptera ; Agromyzidae). *Bulletin de la Société Entomologique d'Égypte*, 60: 53-64.
- 160-TeBeest, D.O. and Templeton, G.E. (1985). Mycoherbicide: progress in the biological control of weeds. *Pl. Dis.*, 69: 6-10.
- 161-Templeton, G.E.; Smith, R.J. and Klomparens, W. (1980). Commercialisation of fungi and bacteria for biological control. *Biocontrol News Inf.*, 1: 291-294.
- 162-Thomas, P.A. (1985). Management of *Salvinia molesta* in Papua, New Guinea. *FAO Plant Prot. Bull.*, 33: 50-56.

- 163-Torrano, L.; Holst, P.J. and Stanley, D.F. (1999). The effect of herbicide and goats on survival and seed production of Illyrian thistle (*Onopordum illyricum* L.). *Plant Protection Quarterly*, 14: 13-15.
- 164-Traoré, D. (1995). Bionomics of *Smicronyx guineanus* Voss and *S. umbrinus* Hustache (Coleoptera : Curculionidae), potential agents of *Striga hermonthica* (Del.) Benth. Ph.D. thesis, Macdonald Campus, McGill University, Quebec, Canada, (unpublished).
- 165-Traoré, D.; Vincent, C. and Stewart, R.K. (1995). Life history of *Smicronyx guineanus* and *S. umbrinus* (Col.: Curculionidae) on *Striga hermonthica* (Scrophulariaceae). *Entomophaga*, 40 (2): 211-221.
- 166-Vere; D.T. and Holst, P.J. (1979). Using goats to control blackberries and briars. *Agricultural Gazette of New South Wales*, 90: 11-13.
- 167-Wapshere, A.J. (1973). A comparison of strategies for screening biological control organisms for weeds. *Proc. 2nd Int. Symp. Biol. Control of Weeds*, Rome 1971. CIBC Misc. Publ. no. 6: 95-102, C.A.B.
- 168-Wapshere, A.J. (1974a). A strategy for evaluating the safety of organisms for biological weed control. *Ann. Appl. Biol.* 77 (2): 201-211.
- 169-Wapshere, A.J. (1974b). Host specificity of phytophagous organisms and the evolutionary centres of plant genera or sub genera. *Entomophaga*, 19 (3): 301-309.
- 170-Wapshere, A.J. (1974c). Towards a science of biological control of weeds. *Proc. 3rd Int. Symp. Biol. Control Weeds*, Montpellier 1973. CIBC Misc. Publ. no. 8: 3-12.
- 171-Wapshere, A.J. (1975). A protocol for programmes for biological control of weeds. *PANS*, 21 (3): 295-303.

- 172-Wapshere, A.J. (1982). Biological control of weeds. W. Holzner and N. Numata (eds.). in *Biology and Ecology of Weeds*. Dr. W. Junk Publisher, The Hague. ISBN 9061936829, 47-56.
- 173-Wapshere, A.J. (1989). A testing sequence for reducing rejection of potential biological control agents for weeds. *Annals of Applied Biology*, 114: 515-526.
- 174-Wapshere, A.J.; Hasan, S.; Wahba, W.K. and Caresche, L. (1974). The ecology of *Chondrilla juncea* in the western Mediterranean. *J. Appl. Ecol.*, 11: 783-800.
- 175-Watson, A.K. (1985). Host specificity of plant pathogens in biological weed control. In: *Proceedings of the VIth International Symposium on the Biological Control of Weeds*, Vancouver, Canada, 1984 (ed. E.S. Delfosse), pp. 577-586. Agriculture Canada.
- 176-Watson, A.K. and Alkhoury, I. (1981). Response of safflower cultivars to *Puccinia jaceae* collected from diffuse knapweed in eastern Europe. In: *Proceedings of the VIth International Symposium on the Biological Control of Weeds*, Brisbane, Australia, 1980. (ed. E.S. Delfosse), pp. 301-305. Melbourne: C.S.I.R.O.
- 177-Weed Research Organization (W.R.O.) (1970). Selected references on broomrape, *Orobancha* spp. 1964-1970. *Annot. Bibl.*, 23: 1-15.
- 178-Weed Research Organization (W.R.O.) (1971). Research on herbivorous fish for weed control, 1957-1971. *Annot. Bibl.*, 31: 1-13.
- 179-Whiteman, J.B. and Room, P.M. (1991). Temperatures lethal to *Salvinia molesta* Mitchell. *Aquatic Botany*, 40: 27-35.
- 180-Whittaker, J.B. and Warrington, S. (1985). An experimental field study of different levels of insect herbivory induced by *Formica rufa* predation on sycamore (*Acer pseudoplatanus*) II. Lepidoptera larvae. *Journal of Applied Ecology*, 22: 787-796.

- 181-Wilson, C.G.; Flanagan, G.J. and Gillett, J.D. (1990). The phytophagous insect fauna of the introduced shrub, *Mimosa pigra* in Northern Australia and its relevance to biological control. *Environmental Entomology*, 19: 776-784.
- 182-Woodburn, T.L. (1993). Host specificity testing, Release and establishment of *Urophora solstitialis* (L.) (Diptera : Tephritidae), a potential biological control agent for *Carduus nutans* L., in Australia. *Biocontrol Science and Technology*, 3: 419-426.
- 183-Woodburn, T.L. (1996). Interspecific competition between *Rhinocyllus conicus* and *Urophora solstitialis*, two biocontrol agents released in Australia against *Carduus nutans*. *Proceedings of IX International Symposium on Biological Control of Weeds*. pp. 409-115. V.C. Moran and J.H. Hoffmann (eds.), 19-26 January. Stellenbosch, South Africa, University of Capetown.
- 184-Woodburn, T.L. (1997). Establishment in Australia of *Trichosirocalus horrius* a biological control agent for *Carduus nutans*, and preliminary assessment of its impact on plant growth and reproductive potential. *Biocontrol Science and Technology*, 7: 645-656.
- 185-Woodburn, T.L. and Cullen, J.M. (1993). Effectiveness of *Rhinocyllus conicus* as a biological control agent for nodding thistle *Carduus nutans*, in Australia. *Proceedings I 10TH Australian Weeds Conference*. Brisbane, Australia, 6-10 September.
- 186-Woodburn, T.L. and Cullen, J. (1996). Impact of *Rhinocyllus conicus* and *Urophora solstitialis* on set achene in *Carduus nutans* in Australia. In: P.D.S. Caligari and D.J.N. Hind (eds.). *Compositae: Biology and Utilization*. *Proceedings of the International Compositae Conference*, Kew, 1994 (D.J.N. Hind, Editor-in-Chief), vol. 2, pp. 307-319. Royal Botanic Gardens, Kew.
- 187-Wright, R. (1927). Goats and noxious-weed control. *New Zealand Journal of Agriculture*, 35: 295-297.

- 188-Zermane, N.; Kroschel, J., Salle, G. and Bouznad, Z. (1994). Prospects for biological control of the parasitic weed *Orobanchae* in Algeria. In: J. Kroschel, M. Abderabihi, H. Betz (eds.). Advances in parasitic weed control at on-farm level. vol. II. joint action to control *Orobanchae* in the WANA Region. Margraf Verlag, Weikersheim, Germany. 135-159.
- 189-Zon, J.C.J. van, Zweerde, W. van der and Hoogers, B.J. (1976). The grass carp, its effects and side effects. Proc. IV Int. Symp. Biol. Contr. Weeds, Gainesville, 251-256.
- 190-Zwölfer, H. (1969). *Urophora affinis* (HG.) (Dipt.: Trypetidae), a potential insect for the biological control of *Centaurea solstitialis* L. in California, CIBC Technical Bulletin, 11: 105-155.

